

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification 7 : C07D 405/12, A61K 31/445, C07D 211/66, 211/46, 309/08, 401/14, 409/12, 405/14, 491/10, A61K 31/453, A61P 19/00</p>	<p>A1</p>	<p>(11) International Publication Number: WO 00/46221 (43) International Publication Date: 10 August 2000 (10.08.00)</p>
<p>(21) International Application Number: PCT/US00/03061 (22) International Filing Date: 7 February 2000 (07.02.00) (30) Priority Data: 60/119,181 8 February 1999 (08.02.99) US (71) Applicant (for all designated States except US): G.D. SEARLE & CO. [US/US]; Corporate Patent Dept., P.O. Box 5110, Chicago, IL 60680-5110 (US). (72) Inventors; and (73) Inventors/Applicants (for US only): DE CRESCENZO, Gary, A. [US/US]; 7345 Spruce Hill Court, St. Charles, MO 63304 (US). RICO, Joseph, G. [US/US]; 524 Wetherby Ter- race Drive, St. Louis, MO 63021 (US). BOEHM, Terri, L. [US/US]; 928 Cleta Drive, Ballwin, MO 63021 (US). CAR- ROLL, Jeffery, N. [US/US]; 2424 South 18th, St. Louis, MO 63104 (US). KASSAB, Darren, J. [US/US]; 16510 Meadview, Chesterfield, MO 63005 (US). MISCHKE, Deb- orah, A. [US/US]; 25 White River Lane, Defiance, MO 63341 (US).</p>	<p>(74) Agents: WARNER, James, M. et al.; G.D. Searle & Co., Corpo- rate Patent Dept., P.O. Box 5110, Chicago, IL 60680-5110 (US). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>	
<p>(54) Title: SULFAMATO HYDROXAMIC ACID METALLOPROTEASE INHIBITOR</p>		
<p>(57) Abstract</p> <p>A sulfamato hydroxamic acid compound that <i>inter alia</i> inhibits matrix metalloprotease activity is disclosed as are a process for preparing the same, intermediate compounds useful in those syntheses, and a treatment process that comprises administering a contemplated sulfamato hydroxamic acid compound in a MMP enzyme-inhibiting effective amount to a host having a condition associated with pathological matrix metalloprotease activity.</p>		

Atty. Docket No. 01136/1/US
Serial No. 10/603,441
Daniel P. Becker, et al.
Reference 31 of 67

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

FOR THE PURPOSES OF INFORMATION ONLY

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BK	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China			PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SF	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

-1-

SULFAMATO HYDROXAMIC ACID
METALLOPROTEASE INHIBITOR

Description

5

Technical Field

This invention is directed to proteinase (protease) inhibitors, and more particularly to the use of sulfamato hydroxamic acid compounds that, *inter alia*, are selective inhibitors of matrix metalloproteinases in a process for treating conditions associated with pathological matrix metalloproteinase activity, the selective inhibitors themselves, compositions of proteinase inhibitors, intermediates for the syntheses of proteinase inhibitors, and processes for the preparation of proteinase inhibitors.

Background of the Invention

Connective tissue, extracellular matrix constituents and basement membranes are required components of all mammals. These components are the biological materials that provide rigidity, differentiation, attachments and, in some cases, elasticity to biological systems including human beings and other mammals. Connective tissues components include, for example, collagen, elastin, proteoglycans, fibronectin and laminin. These biochemicals makeup, or are components of structures, such as skin, bone, teeth, tendon, cartilage, basement membrane, blood vessels, cornea and vitreous humor.

-2-

Under normal conditions, connective tissue turnover and/or repair processes are controlled and in equilibrium. The loss of this balance for whatever reason leads to a number of disease states.

5 Inhibition of the enzymes responsible loss of equilibrium provides a control mechanism for this tissue decomposition and, therefore, a treatment for these diseases.

Degradation of connective tissue or
10 connective tissue components is carried out by the action of proteinase enzymes released from resident tissue cells and/or invading inflammatory or tumor cells. A major class of enzymes involved in this function are the zinc metalloproteinases
15 (metalloproteases).

The metalloprotease enzymes are divided into classes with some members having several different names in common use. Examples are:
collagenase I (MMP-1, fibroblast collagenase; EC
20 3.4.24.3); collagenase II (MMP-8, neutrophil collagenase; EC 3.4.24.34), collagenase III (MMP-13), stromelysin 1 (MMP-3; EC 3.4.24.17), stromelysin 2 (MMP-10; EC 3.4.24.22), proteoglycanase, matrilysin (MMP-7), gelatinase A (MMP-2, 72 kDa gelatinase,
25 basement membrane collagenase; EC 3.4.24.24), gelatinase B (MMP-9, 92 kDa gelatinase; EC 3.4.24.35), stromelysin 3 (MMP-11), metalloelastase (MMP-12, HME, human macrophage elastase) and membrane MMP (MMP-14). MMP is an abbreviation or acronym
30 representing the term Matrix Metalloprotease with the attached numerals providing differentiation between specific members of the MMP group.

The uncontrolled breakdown of connective tissue by metalloproteases is a feature of many pathological conditions. Examples include rheumatoid arthritis, osteoarthritis, septic arthritis; corneal, epidermal or gastric ulceration; tumor metastasis, invasion or angiogenesis; periodontal disease; proteinuria; Alzheimers Disease; coronary thrombosis and bone disease. Defective injury repair processes also occur. This can produce improper wound healing leading to weak repairs, adhesions and scarring. These latter defects can lead to disfigurement and/or permanent disabilities as with post-surgical adhesions.

Metalloproteases are also involved in the biosynthesis of tumor necrosis factor (TNF), and inhibition of the production or action of TNF and related compounds is an important clinical disease treatment mechanism. TNF- α , for example, is a cytokine that at present is thought to be produced initially as a 28 kD cell-associated molecule. It is released as an active, 17 kD form that can mediate a large number of deleterious effects *in vitro* and *in vivo*. For example, TNF can cause and/or contribute to the effects of inflammation, rheumatoid arthritis, autoimmune disease, multiple sclerosis, graft rejection, fibrotic disease, cancer, infectious diseases, malaria, mycobacterial infection, meningitis, fever, psoriasis, cardiovascular/pulmonary effects such as post-ischemic reperfusion injury, congestive heart failure, hemorrhage, coagulation, hyperoxic alveolar injury, radiation damage and acute phase responses like those seen with infections and sepsis and during shock such as septic

-4-

shock and hemodynamic shock. Chronic release of active TNF can cause cachexia and anorexia. TNF can be lethal, and TNF can help control the growth of tumor cells.

5 TNF- α convertase is a metalloprotease involved in the formation of soluble TNF- α . Inhibition of TNF- α convertase (TACE) inhibits production of active TNF- α . Compounds that inhibit both MMPs activity and TNF- α production have been
10 disclosed in WIPO International Publication Nos. WO 94/24140, WO 94/02466 and WO 97/20824. Compounds that inhibit MMPs such as collagenase, stromelysin and gelatinase have been shown to inhibit the release of TNF (Gearing et al. *Nature* 376, 555-557 (1994),
15 McGeehan et al., *Nature* 376, 558-561 (1994)). There remains a need for effective MMP inhibitors. There also remains a need for effective TNF- α convertase inhibiting agents.

 MMPs are involved in other biochemical
20 processes in mammals as well. Included is the control of ovulation, post-partum uterine involution, possibly implantation, cleavage of APP (β -Amyloid Precursor Protein) to the amyloid plaque and inactivation of α_1 -protease inhibitor (α_1 -PI).
25 Inhibition of these metalloproteases permits the control of fertility and the treatment or prevention of Alzheimers Disease. In addition, increasing and maintaining the levels of an endogenous or administered serine protease inhibitor drug or
30 biochemical such as α_1 -PI supports the treatment and prevention of diseases such as emphysema, pulmonary

diseases, inflammatory diseases and diseases of aging such as loss of skin or organ stretch and resiliency.

Inhibition of selected MMPs can also be desirable in other instances. Treatment of cancer and/or inhibition of metastasis and/or inhibition of angiogenesis are examples of approaches to the treatment of diseases wherein the selective inhibition of stromelysin, gelatinase A or B, or collagenase III appear to be the relatively most important enzyme or enzymes to inhibit especially when compared with collagenase I (MMP-1). A drug that does not inhibit collagenase I can have a superior therapeutic profile. Osteoarthritis, another prevalent disease wherein it is believed that cartilage degradation of inflamed joints is at least partially caused by MMP-13 released from cells such as stimulated chondrocytes, may be best treated by administration of drugs one of whose modes of action is inhibition of MMP-13. See, for example, Mitchell et al., *J. Clin. Invest.*, 97:761-768 (1996) and Reboul et al., *J. Clin. Invest.*, 97:2011-2019 (1996).

Inhibitors of metalloproteases are known. Examples include natural biochemicals such as tissue inhibitors of metalloproteinases (TIMPs), α_2 -macroglobulin and their analogs or derivatives. These endogenous inhibitors are high molecular weight protein molecules that form inactive complexes with metalloproteases. A number of smaller peptide-like compounds that inhibit metalloproteases have been described. Mercaptoamide peptidyl derivatives have shown ACE inhibition *in vitro* and *in vivo*. Angiotensin converting enzyme (ACE) aids in the

production of angiotensin II, a potent pressor substance in mammals and inhibition of this enzyme leads to the lowering of blood pressure.

Thiol group-containing amide or peptidyl
5 amide-based metalloprotease (MMP) inhibitors are known as is shown in, for example, WO95/12389, WO96/11209 and U.S. 4,595,700. Hydroxamate group-containing MMP inhibitors are disclosed in a number of published patent applications such as WO 95/29892,
10 WO 97/24117, WO 97/49679 and EP 0 780 386 that disclose carbon back-boned compounds, and WO 90/05719, WO 93/20047, WO 95/09841 and WO 96/06074 that disclose hydroxamates that have a peptidyl back-bones or peptidomimetic back-bones, as does the
15 article by Schwartz et al., *Progr. Med. Chem.*, 29:271-334(1992) and those of Rasmussen et al., *Pharmacol. Ther.*, 75(1): 69-75 (1997) and Denis et al., *Invest. New Drugs*, 15(3): 175-185 (1997). In addition, application EP 0757 984 A1 discloses
20 aromatic sulfonamide hydroxamates in which the sulfonamido sulfonyl group is bonded on one side to a phenyl ring and the sulfonamido nitrogen is bonded to the hydroxamate group via a chain of one to four carbon atoms.

25 One possible problem associated with known MMP inhibitors is that such compounds often exhibit the same or similar inhibitory effects against each of the MMP enzymes. For example, the peptidomimetic hydroxamate known as batimastat is reported to
30 exhibit IC₅₀ values of about 1 to about 20 nanomolar (nM) against each of MMP-1, MMP-2, MMP-3, MMP-7, and MMP-9. Marimastat, another peptidomimetic

-7-

hydroxamate was reported to be another broad-spectrum MMP inhibitor with an enzyme inhibitory spectrum very similar to batimastat, except that marimastat exhibited an IC_{50} value against MMP-3 of 230 nM.

- 5 Rasmussen et al., *Pharmacol. Ther.*, 75(1): 69-75 (1997).

Meta analysis of data from Phase I/II studies using marimastat in patients with advanced, rapidly progressive, treatment-refractory solid tumor
10 cancers (colorectal, pancreatic, ovarian, prostate) indicated a dose-related reduction in the rise of cancer-specific antigens used as surrogate markers for biological activity. Although marimastat exhibited some measure of efficacy via these markers,
15 toxic side effects were noted. The most common drug-related toxicity of marimastat in those clinical trials was musculoskeletal pain and stiffness, often commencing in the small joints in the hands, spreading to the arms and shoulder. A short dosing
20 holiday of 1-3 weeks followed by dosage reduction permits treatment to continue. Rasmussen et al., *Pharmacol. Ther.*, 75(1): 69-75 (1997). It is thought that the lack of specificity of inhibitory effect among the MMPs may be the cause of that effect.

- 25 International application WO 98/38163, published on September 3, 1998 disclose a large group of hydroxamate inhibitors of MMPs and TACE. The compounds of WO 98/38163 contain one or two substituents adjacent to the hydroxamate
30 functionality and a substituent that can be an

aromatic sulfonyl group adjacent to those one or two substituents.

International application WO 98/37877, published on September 3, 1998 discloses compounds
5 that contain a 5- to 7-membered heterocyclic ring adjacent to the hydroxamate functionality and can contain an aromatic sulfonyl group adjacent to the heterocyclic ring.

More recently, WO 99/24399, published on 20
10 may, 1999, teaches hydroxamate compounds said to have activity in inhibiting MMP and TNF. Those inhibitors are exemplified by compounds having a three carbon atom chain linked to a sulfonamido group. The hydroxamate carbon is linked to a carbon that can be
15 substituted and that carbon is linked to a methylene. The methylene is linked to a sulfonyl that is bonded to a nitrogen that is further substituted. This disclosure also lacks disclosure as to possible specificity of activity among the substrate enzymes.

20 Another recent disclosure is that of WO 99/29667, published on 17 June, 1999, that discloses two carbon hydroxamate containing a sulfonamido group whose nitrogen atom is in a ring that is typically bonded directly to another one or two ring group
25 without the intermediacy of another atom. This publication suggests that some of its compounds are selective inhibitors, but provides scant data for only seven compounds.

Although many of the known MMP inhibitors
30 such as batimastat, marimastat and the hydroxamates of WO 98/37877 and WO 98/38163 exhibit a broad spectrum of activity against MMPs, those compounds

- 9 -

are not particularly selective in their inhibitory activity. This lack of selectivity may be the cause of the musculoskeletal pain and stiffness observed with their use. In addition, it can be

5 therapeutically advantageous to utilize a medicament that is selective in its activity as compared to a generally active material so that treatment can be more closely tailored to the pathological condition presented by the host mammal. The disclosure that

10 follows describes a process for treating a host mammal having a condition associated with pathological matrix metalloprotease activity that utilizes a compound that selectively inhibits one or more MMPs, while exhibiting less activity against at

15 least MMP-1.

Summary of the Invention

The present invention is directed to a treatment process that comprises administering a

20 contemplated sulfamato hydroxamic acid metalloprotease inhibitor in an effective amount to a host mammal having a condition associated with pathological metalloprotease activity. A contemplated molecule, *inter alia*, exhibits excellent

25 inhibitory activity of one or more matrix metalloprotease (MMP) enzymes, such as MMP-2 and MMP-13, while exhibiting substantially less inhibition at least of MMP-1. By "substantially less" it is meant that a contemplated compound exhibits an IC₅₀ value


30 ratio against one or both of MMP-2 or MMP-13 as compared to its IC₅₀ value against MMP-1, e.g., IC₅₀

.

-10-

MMP-2:IC₅₀ MMP-1, that is less than about 1:10, preferably less than about 1:100, and most preferably less than about 1:1000 in the *in vitro* inhibition assay utilized hereinafter. The invention also
5 contemplates particular compounds that selectively inhibit the activity of MMP-2 to a greater extent than MMP-13, as well as a composition containing such a MMP inhibitor as active ingredient and a process for using the same. A contemplated compound also
10 exhibits inhibition of the activity of the adamalysin family of enzymes, exemplified by the enzyme ADAM 10. The invention further contemplates intermediates in the preparation of a contemplated sulfamato hydroxamic acid molecule and a process for preparing
15 a sulfamato hydroxamic acid molecule.

Briefly, one embodiment of the present invention is directed to a treatment process that comprises administering a contemplated sulfamato hydroxamic acid metalloprotease inhibitor that
20 selectively inhibits matrix metalloprotease and adamalysin activity as above in an effective amount to a host mammal having a condition associated with pathological metalloprotease activity. The administered enzyme inhibitor sulfamato hydroxamic
25 acid (hydroxamate) corresponds in structure to formula I, below, or a pharmaceutically acceptable salt thereof:



I

wherein

R^1 and R^2 are preferably taken together with the carbon to which they are bonded form a
5 cycloalkyl or more preferably a heterocyclo group either of which is optionally substituted by one, two or three R^X substituents, or R^1 and R^2 are independently selected from the group consisting of:

hydrido,

10 an alkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

an alkyloxyalkyl group, optionally substituted with one, two or three groups independently selected
15 from R^X substituents,

an alkylthioalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

an alkenyl group, optionally substituted with
20 one, two or three groups independently selected from R^X substituents,

an alkynyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

25 an aryl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

an arylalkyl group, optionally substituted with one, two or three groups independently selected from
30 R^X substituents,

- an arylalkyloxyalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,
- an aryloxyalkyl group, optionally substituted
5 with one, two or three groups independently selected from R^X substituents,
- an arylthioalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,
- 10 an arylalkylthioalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,
- a cycloalkyl or bicycloalkyl group, optionally substituted with one, two or three groups
15 independently selected from R^X substituents,
- a cycloalkenyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,
- a cycloalkylalkyl or bicycloalkylalkyl group,
20 optionally substituted with one, two or three groups independently selected from R^X substituents,
- a cycloalkyloxyalkyl or bicycloalkyloxyalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,
- 25 a cycloalkylalkyloxyalkyl or bicycloalkyloxyalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,
- a cycloalkylthioalkyl or bicycloalkylthioalkyl
30 group, optionally substituted with one, two or three groups independently selected from R^X;

cycloalkylalkylthioalkyl or
bicycloalkylalkylthioalkyl, optionally substituted
with one, two or three groups independently selected
from R^X substituents,

5 a heterocyclo group, optionally substituted with
one, two or three groups independently selected from
R^X substituents,

 a heterocycloalkyl group, optionally substituted
with one, two or three groups independently selected
10 from R^X substituents

 a heteroaryl group, optionally substituted with
one, two or three groups independently selected from
R^X substituents,

 a biarylalkyl group, optionally substituted with
15 one, two or three groups independently selected from
R^X substituents,

 an arylalkenyl group, optionally substituted
with one, two or three groups independently selected
from R^X substituents,

20 an arylalkynyl group, optionally substituted
with one, two or three groups independently selected
from R^X substituents,

 a heterocycloalkylthio group, optionally
substituted with one, two or three groups selected
25 independently from R^X substituents,

 a heterocycloalkyloxyalkyl group, optionally
substituted with one, two or three groups selected
independently from R^X substituents,

 a heteroarylalkenyl group, optionally
30 substituted with one, two or three groups
independently selected from R^X substituents, and

a heteroarylalkyloxyalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents;

wherein an R^X substituent is selected from the

5 group consisting of a hydrido, aryl, heteroaryl, heterocyclo, aroyl, alkanoyl, heteroaroyl, halogen (F, Cl, Br, I), cyano, aldehydo (CHO, formyl), hydroxy, R^{CR^d} -amino ($-NR^{CR^d}$), R^{CR^d} -aminoalkyl, nitro, nitroso, alkyl, alkenyl, alkynyl, cycloalkyl,

10 cycloalkenyl, alkoxy, aryloxy, heteroaryloxy, alkenyloxy, alkynyloxy, alkoxyaryl, alkoxyheteroaryl, alkoxyalkyl, R^C -oxyalkyl, alkoxyalkyl, alkylenedioxy, aryloxyalkyl, perfluoroalkyl, trifluoroalkyl, alkylthio, arylthio, alkylloxycarbonyl,

15 alkylloxycarbonyloxy, aryloxycarbonyl, arylalkylloxycarbonyl, alkylloxycarbonyl- R^C -amino, arylalkylloxycarbonyl- R^C -amino, aryloxycarbonyloxy, carboxy, R^{CR^d} -aminocarbonyloxy, R^{CR^d} -aminocarbonyl, R^{CR^d} -aminoalkanoyl, hydroxy- R^C -aminocarbonyl, R^{CR^d} -

20 aminosulfonyl, arylsulfonyl(R^C)amino, R^{CR^d} -aminoalkoxy, R^{CR^d} -aminocarbonyl(R^C)amino, trifluoromethylsulfonyl(R^C)amino, heteroarylsulfonyl- (R^C)amino, alkylsulfonyl, arylsulfonyl(R^C)amino, arylsulfonyl(R^C)aminocarbonyl, alkylsulfonyl-

25 (R^C)amino, arylcarbonyl(R^C)-aminosulfonyl, and an alkylsulfonyl(R^C)aminocarbonyl substituent;

wherein R^C and R^d are independently selected from the group consisting of a hydrido, alkanoyl, arylalkyl, aroyl, bisalkoxyalkyl, alkyl, haloalkyl,

30 perfluoroalkyl, trifluoromethylalkyl,

- perfluoroalkoxyalkyl, alkoxyalkyl, cycloalkyl,
heterocycloalkyl, heterocycloalkylcarbonyl, aryl,
heterocyclo, heteroaryl, cycloalkylalkyl,
aryloxyalkyl, heteroaryloxyalkyl,
5 heteroarylalkoxyalkyl, heteroarylthioalkyl,
arylsulfonyl, alkylsulfonyl, heteroarylsulfonyl,
carboxyalkyl, alkoxycarbonylalkyl, aminocarbonyl,
alkyliminocarbonyl, aryliminocarbonyl,
heterocycloiminocarbonyl, arylthioalkyl,
10 alkylthioalkyl, arylthioalkenyl, alkylthioalkenyl,
heteroarylalkyl, haloalkanoyl, hydroxyalkanoyl,
thiolalkanoyl, alkenyl, alkynyl, alkoxyalkyl,
alkoxycarbonyl, aryloxycarbonyl, aminoalkylcarbonyl,
hydroxyalkyl, aminoalkyl, aminoalkylsulfonyl,
15 aminosulfonyl wherein the amino nitrogen is (i)
unsubstituted or (ii) independently substituted with
one or two R^Y radicals, or the substituents on the
amino group taken together with the amino nitrogen
form a saturated or partially unsaturated heterocyclo
20 group optionally substituted with one, two or three
groups independently selected from R^W substituents or
a heteroaryl group optionally substituted with one,
two or three groups independently selected from R^V
substituents;
25 wherein R^Y is selected from the group consisting
of an arylalkyl, aryl, heteroaryl, heterocyclo,
alkyl, alkynyl, alkenyl, alkoxyalkyl, alkoxyalkyl,
substituted or unsubstituted aminoalkyl,
alkyloxycarbonyl, arylalkyloxycarbonyl, carboxyalkyl,
30 haloalkyl, alkanoyl, aroyl, substituted or
unsubstituted aminoalkanoyl, halo alkanoyl and a
hydroxyalkyl group, each of which groups is

optionally substituted by one or two groups independently selected from R^U substituents as are the substituents of the substituted aminoalkyl and substituted aminoalkanoyl groups;

5 wherein R^V is selected from the group consisting of a hydrido, aryl, heteroaryl, heterocyclo, aroyl, alkanoyl, heteroaroyl, halogen (F, Cl, Br, I), cyano, aldehydo (CHO, formyl), hydroxy, amino, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, alkoxy, 10 aryloxy, heteroaryloxy, alkenyloxy, alkynyloxy, alkoxyaryl, alkoxyheteroaryl, $R^Y R^Z$ -amino, alkoxyalkyl, alkylenedioxy, aryloxyalkyl, perfluoroalkyl, trifluoroalkyl, alkylthio, arylthio, alkyloxycarbonyl, alkyloxycarbonyloxy, 15 aryloxycarbonyl, arylalkyloxycarbonyl, arylalkyloxycarbonylamino, aryloxycarbonyloxy, carboxy, $R^Y R^Z$ -aminocarbonyloxy, $R^Y R^Z$ -aminocarbonyl, $R^Y R^Z$ -aminoalkanoyl, hydroxyaminocarbonyl, $R^Y R^Z$ -aminosulfonyl, $R^Y R^Z$ -aminocarbonyl(RY)amino, 20 trifluoromethylsulfonyl(RY)amino, heteroarylsulfonyl-(RY)amino, arylsulfonyl(RY)amino, arylsulfonyl(RY)-aminocarbonyl, alkylsulfonyl(RY)amino, arylcarbonyl-(RY)aminosulfonyl, and an alkylsulfonyl(RY)-aminocarbonyl substituent;

25 wherein R^W is selected from the group consisting of a hydrido, aryl, heteroaryl, heterocyclo, aroyl, alkanoyl, heteroaroyl, halogen (F, Cl, Br, I), cyano, aldehydo (CHO, formyl), hydroxy, amino, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, alkoxy, 30 aryloxy, heteroaryloxy, alkenyloxy, alkynyloxy, alkoxyaryl, alkoxyheteroaryl, $R^Y R^Z$ -amino,

alkoxyalkyl, alkylenedioxy, aryloxyalkyl,
perfluoroalkyl, trifluoroalkyl, alkylthio, arylthio,
alkyloxycarbonyl, alkyloxycarbonyloxy,
aryloxycarbonyl, arylalkyloxycarbonyl,
5 arylalkyloxycarbonylamino, aryloxycarbonyloxy,
carboxy, RYR^Z-aminocarbonyloxy, RYR^Z-aminocarbonyl,
RYR^Z-aminoalkanoyl, hydroxyaminocarbonyl, RYR^Z-
aminosulfonyl, RYR^Z-aminocarbonyl (RY) amino,
trifluoromethylsulfonyl (RY) amino, heteroarylsulfonyl-
10 (RY) amino, arylsulfonyl (RY) amino, arylsulfonyl (RY) -
aminocarbonyl, alkylsulfonyl (RY) amino, arylcarbonyl-
(RY) aminosulfonyl, and an alkylsulfonyl (RY) -
aminocarbonyl substituent;

R^Z is selected from the group consisting of an
15 arylalkyl, aryl, heteroaryl, heterocyclo, alkyl,
alkynyl, alkenyl, alkoxyalkyl, alkoxyalkylalkyl,
substituted or unsubstituted aminoalkyl,
alkyloxycarbonyl, arylalkyloxycarbonyl, carboxyalkyl,
haloalkyl, alkanoyl, aroyl, substituted or
20 unsubstituted aminoalkanoyl, halo alkanoyl and a
hydroxyalkyl group, each of which groups are
optionally substituted by one or two R^U substituents;

wherein R^U is selected from the group consisting
of an arylalkyl, aryl, heteroaryl, heterocyclo,
25 alkyl, alkynyl, alkenyl, alkoxyalkyl, alkoxyalkyl,
substituted or unsubstituted aminoalkyl,
alkyloxycarbonyl, arylalkyloxycarbonyl, carboxyalkyl,
haloalkyl, alkanoyl, aroyl, substituted or
unsubstituted aminoalkanoyl, halo alkanoyl and a
30 hydroxyalkyl group, wherein the substituents of the
substituted aminoalkyl and substituted aminoalkanoyl

-18-

groups are selected from the group consisting of an alkyl, alkenyl, alkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, aryloxycarbonyl and an alkyloxycarbonyl group;

5 R^{3a} and R^{3b} are independently selected from the group consisting of a hydrido, alkyl, alkenyl, alkynyl, aryl, heterocyclo, heteroaryl, cycloalkyl, and an alkoxyalkyl group, each of which groups is optionally substituted by an -AREY substituent;

10 in that AREY substituent, A is selected from the group consisting of

- (1) -O-;
- (2) -S-;
- (3) -N(R^e)-;
- 15 (4) -CO-N(R^e) or -N(R^e)-CO-;
- (5) -CO-O- or -O-CO-;
- (6) -O-CO-O-;
- (7) -HC=CH-;
- (8) -NH-CO-NH-;
- 20 (9) -C≡C-;
- (10) -NH-CO-O- or -O-CO-NH-;
- (11) -N=N-;
- (12) -NH-NH-;
- (13) -CS-N(R^e)- or -N(R^e)-CS-;
- 25 (14) -CH₂-;
- (15) -O-[(CH₂)₁₋₈]- or -[(CH₂)₁₋₈]O-; and
- (16) -S-CH₂- or -CH₂-S-; or
- (17) A is absent and R is directly
- 30 connected to R^{3a} or R^{3b}, or both R^{3a} and R^{3b};

the moiety R is selected from the group consisting of alkyl, alkoxyalkyl, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaralkyl, heterocycloalkyl, cycloalkylalkyl, cycloalkoxyalkyl, 5 heterocycloalkoxyalkyl, aryloxyalkyl, heteroaryloxyalkyl, arylthioalkyl, heteroarylthioalkyl, cycloalkylthioalkyl, and a heterocycloalkylthioalkyl group wherein the aryl, heteroaryl, cycloalkyl or heterocycloalkyl 10 substituent is (i) unsubstituted or (ii) substituted with one or two radicals selected from the group consisting of a halo, alkyl, perfluoroalkyl, perfluoroalkoxy, perfluoroalkylthio, trifluoromethylalkyl, amino, alkoxycarbonylalkyl, 15 alkoxy, C₁-C₂-alkylene-dioxy, hydroxycarbonylalkyl, hydroxycarbonylalkylamino, nitro, hydroxy, hydroxyalkyl, alkanoylamino, and a alkoxycarbonyl group;

the group E is selected from the group 20 consisting of

- (1) -CO(R^W)- or -(R^W)CO-;
- (2) -CON(R^E)- or -(R^E)NCO-;
- (3) -CO-;
- (4) -SO₂-R^W or -R^WSO₂-;
- 25 (5) -SO₂-;
- (6) -N(R^E)-SO₂- or -SO₂-N(R^E)-; or
- (7) E is absent and R is bonded directly to Y; and

Y is absent or is selected from the group 30 consisting of a hydrido, alkyl, alkoxy, haloalkyl, aryl, aralkyl, cycloalkyl, heteroaryl, hydroxy,

-20-

- nitride, nitro, aryloxy, aralkoxy, heteroaryloxy,
heteroaralkyl, R^Coxyalkyl, perfluoroalkoxy,
perfluoroalkylthio, trifluoromethylalkyl, alkenyl,
heterocycloalkyl, cycloalkyl, trifluoromethyl,
5 alkoxy carbonyl, and a aminoalkyl group, wherein the
aryl, heteroaryl or heterocycloalkyl group is (i)
unsubstituted or (ii) substituted with one or two
radicals independently selected from the group
consisting of an alkanoyl, halo, nitro, nitride,
10 haloalkyl, alkyl, aralkyl, aryl, alkoxy, and an amino
group wherein the amino nitrogen is (i) unsubstituted
or (ii) substituted with one or two groups
independently selected from hydrido, alkyl, and an
aralkyl group;
- 15 wherein R^e is selected from hydrido, alkyl,
alkenyl, alkenyl, aryl, arylalkyl, heteroaryl,
heteroarylalkyl, aryloxy carbonyl, alkyloxy carbonyl,
R^CR^damino carbonyl, R^CR^daminosulfonyl,
R^CR^daminoalkanoyl and R^CR^daminoalkylsulfonyl, and R^C,
20 R^d and R^w are as defined before; or
- R^{3a} and R^{3b} taken together with the nitrogen atom
to which they are bonded form a group -GAREY wherein
G is a N-heterocyclo group;
the substituent A is selected from the
25 group consisting of
- (1) -O-;
 - (2) -S-;
 - (3) -NR^e-;
 - (4) -CO-N(R^e) or -N(R^e)-CO-;
 - 30 (5) -CO-O- or -O-CO-;
 - (6) -O-CO-O-;

-21-

- (7) -HC=CH- ;
(8) -NH-CO-NH- ;
(9) $\text{-C}\equiv\text{C-}$;
(10) -NH-CO-O- or -O-CO-NH- ;
5 (11) -N=N- ;
(12) -NH-NH- ;
(13) $\text{-CS-N(R}^{\text{e}}\text{)-}$ or $\text{-N(R}^{\text{e}}\text{)-CS-}$;
(14) $\text{-CH}_2\text{-}$;
(15) $\text{-O-[(CH}_2\text{)}_{1-8}\text{-}$ or $\text{-[(CH}_2\text{)}_{1-8}\text{]O-}$; and
10 (16) $\text{-S-CH}_2\text{-}$ or $\text{-CH}_2\text{-S-}$; or
(17) A is absent and R is directly connected to G;

The moiety R is selected from the group consisting of alkyl, alkoxyalkyl, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaralkyl, heterocycloalkyl, cycloalkylalkyl, cycloalkoxyalkyl, heterocycloalkoxyalkyl, aryloxyalkyl, heteroaryloxyalkyl, arylthioalkyl, heteroarylthioalkyl, cycloalkylthioalkyl, and a
15 heterocycloalkylthioalkyl group wherein the aryl or heteroaryl or cycloalkyl or heterocycloalkyl substituent is (i) unsubstituted or (ii) substituted with one or two radicals selected from the group consisting of a halo, alkyl, perfluoroalkyl, perfluoroalkoxy, perfluoroalkylthio,
20 trifluoromethylalkyl, amino, alkoxycarbonylalkyl, alkoxy, $\text{C}_1\text{-C}_2\text{-alkylene-dioxy}$, hydroxycarbonylalkyl, hydroxycarbonylalkylamino, nitro, hydroxy, hydroxyalkyl, alkanoylamino, and a alkoxycarbonyl
25 group;
30

The moiety E is selected from the group consisting of


-22-

- (1) $-\text{CO}(\text{R}^{\text{W}})-$ or $-(\text{R}^{\text{W}})\text{CO}-$;
- (2) $-\text{CONH}-$ or $-\text{HNCO}-$;
- (3) $-\text{CO}-$;
- (4) $-\text{SO}_2-\text{R}^{\text{W}}-$ or $-\text{R}^{\text{W}}-\text{SO}_2-$;
- 5 (5) $-\text{SO}_2-$;
- (6) $-\text{NH}-\text{SO}_2-$ or $-\text{SO}_2-\text{NH}-$; or
- (7) E is absent and Y is bonded directly to R; and

The moiety Y is absent or is selected from the group consisting of a hydrido, alkyl, alkoxy, haloalkyl, aryl, aralkyl, cycloalkyl, heteroaryl, hydroxy, aryloxy, aralkoxy, heteroaryloxy, heteroaralkyl, perfluoroalkoxy, perfluoroalkylthio, trifluoromethylalkyl, alkenyl, heterocycloalkyl, cycloalkyl, trifluoromethyl, alkoxycarbonyl, and a
10 aminoalkyl group, wherein the aryl or heteroaryl or heterocycloalkyl group is (i) unsubstituted or (ii) substituted with one or two radicals independently selected from the group consisting of an alkanoyl, halo, nitro, aralkyl, aryl, alkoxy, and an amino
15 group wherein the amino nitrogen is (i) unsubstituted or (ii) substituted with one or two groups independently selected from hydrido, alkyl, and an aralkyl group.

25 More generally, a contemplated compound includes an inhibitor utilized as discussed above, as well as a pro-drug form of such a compound and also an intermediate used in the synthesis of a hydroxamate or hydroxamate pro-drug. Such a more
30 general compound corresponds in structure to formula II, below,

-23-



wherein R^1 , R^2 , R^{3a} and R^{3b} are as before described, and

- 5 R^{20} is (a) $-O-R^{21}$, where R^{21} is selected from the group consisting of a hydrido, C_1 - C_6 -alkyl, aryl, ar- C_1 - C_6 -alkyl group and a pharmaceutically acceptable cation, (b) $-NH-O-R^{22}$, wherein R^{22} is a selectively removable protecting group such as a 2-
- 10 tetrahydropyranyl, benzyl, p-methoxybenzyl (MOZ) carbonyl- C_1 - C_6 -alkoxy, trisubstituted silyl group or o-nitrophenyl group, peptide synthesis resin and the like, wherein trisubstituted silyl group is substituted with C_1 - C_6 -alkyl, aryl, or ar- C_1 - C_6 -
- 15 alkyl, or (c) $-NH-O-R^{14}$, where R^{14} is hydrido, a pharmaceutically acceptable cation or $C(W)R^{15}$ where W is O or S and R^{15} is selected from the group consisting of an C_1 - C_6 -alkyl, aryl, C_1 - C_6 -alkoxy, heteroaryl- C_1 - C_6 -alkyl, C_3 - C_8 -cycloalkyl- C_1 - C_6 -alkyl,
- 20 aryloxy, ar- C_1 - C_6 -alkoxy, ar- C_1 - C_6 -alkyl, heteroaryl and amino C_1 - C_6 -alkyl group wherein the aminoalkyl nitrogen is (i) unsubstituted or (ii) substituted with one or two substituents independently selected from the group consisting of an C_1 - C_6 -alkyl, aryl,
- 25 ar- C_1 - C_6 -alkyl, C_3 - C_8 -cycloalkyl- C_1 - C_6 -alkyl, ar- C_1 - C_6 -alkoxycarbonyl, C_1 - C_6 -alkoxycarbonyl, and C_1 - C_6 -alkanoyl radical, or (iii) wherein the amino C_1 - C_6 -

alkyl nitrogen and two substituents attached thereto form a 5- to 8-membered heterocyclo or heteroaryl ring.

Among the several benefits and advantages
5 of the present invention are the provision of compounds and compositions effective as inhibitors of matrix metalloproteinase activity, the provision of such compounds and compositions that are effective for the inhibition of metalloproteinases implicated
10 in diseases and disorders involving uncontrolled breakdown of connective tissue.

More particularly, a benefit of this invention is the provision of a compound and composition effective for selectively inhibiting
15 certain metalloproteinases, such as one or both of MMP-2 and MMP-13, associated with pathological conditions such as, for example, rheumatoid arthritis, osteoarthritis, septic arthritis, corneal, epidermal or gastric ulceration, tumor metastasis,
20 invasion or angiogenesis, periodontal disease, proteinuria, Alzheimer's Disease, coronary thrombosis and bone disease.

An advantage of the invention is the provision of compounds, compositions and methods
25 effective for treating such pathological conditions by selective inhibition of a metalloproteinase such as MMP-2 or MMP-13 associated with such conditions with minimal side effects resulting from inhibition of other metalloproteinases, such as MMP-1, whose
30 activity is necessary or desirable for normal body function.

-25-

A still further benefit of the invention is that a contemplated compound exhibits greater inhibition of MMP-2 than MMP-13.

A still further advantage of the invention
5 is that a contemplated compound exhibits inhibitory activity against the adamalysin family of enzymes.

Yet another advantage of the invention is the provision of a process for preparing such compounds.

10 Another benefit is the provision of a method for treating a pathological condition associated with abnormal matrix metalloproteinase activity.

A further advantage of the invention is the
15 provision of a process for preparing such compositions.

Still further benefits and advantages of the invention will be apparent to the skilled worker from the disclosure that follows.

20

Detailed Description of the Invention

In accordance with the present invention, it has been discovered that certain sulfamato hydroxamic acids (hydroxamates) are effective for inhibition of
25 matrix metalloproteinases ("MMPs") believed to be associated with uncontrolled or otherwise pathological breakdown of connective tissue. In particular, it has been found that these certain sulfamato hydroxamates are effective for inhibition
30 of one or both of MMP-2 and MMP-13, which can be particularly destructive to tissue if present or generated in abnormal quantities or concentrations, and thus exhibit a pathological activity. Included

in that pathological activity is the assistance of tumors and tumor cells in the process of penetrating basement membrane, and developing a new or improved blood supply; i.e., angiogenesis.

5 Moreover, it has been discovered that these sulfamato hydroxamates are selective in the inhibition of one or both of MMP-2 and MMP-13 without excessive inhibition of other collagenases essential to normal bodily function such as tissue turnover and
10 repair. More particularly, it has been found that a contemplated sulfamato hydroxamate of the invention, or a pharmaceutically acceptable salt thereof, is particularly active in inhibiting of one or both of MMP-2 and MMP-13 in an *in vitro* assay that is
15 predictive of *in vivo* activity. In addition, while being selective for one or both of MMP-2 and MMP-13, a contemplated sulfamato hydroxamate, or its salt, has a limited or minimal *in vitro* inhibitory effect on MMP-1. This point is illustrated in the
20 Inhibition Table hereinafter. Put differently, a contemplated compound can inhibit the activity of MMP-2 or MMP-13 compared to MMP-1.

 The advantages of the selectivity of a contemplated compound can be appreciated, without
25 wishing to be bound by theory, by considering the therapeutic uses the compounds. For example, inhibition of MMP-1 is suggested to be undesirable due to its role as a housekeeping enzyme, helping to maintain normal connective tissue turnover and
30 repair. Inhibition of MMP-1 can lead to toxicities or side effects such as such as joint or connective tissue deterioration and pain. On the other hand, MMP-13 has been suggested to be intimately involved

-27-

in the destruction of joint components in diseases such as osteoarthritis. Thus, potent and selective inhibition of MMP-13 compared with inhibition MMP-1 is highly desirable because a MMP-13 inhibitor can
5 have a positive effect on disease progression in a patient in addition to having an anti-inflammatory effect.

Inhibition of MMP-2 can be desirable for inhibition of tumor growth, metastasis, invasion
10 and/or angiogenesis. A profile of selective inhibition of MMP-2 relative to MMP-1 can provide a therapeutic advantage. A contemplated compound not only is substantially more active in inhibiting MMP-2 than MMP-1, a contemplated compound also exhibits
15 greater inhibition of MMP-2 than MMP-13.


A further advantage to MMP inhibitors with selective inhibition profiles is their suitability for use in combination with other types of medicaments. For example, a patient can be treated
20 with an MMP inhibitor compound for the inhibition of angiogenesis in combination with a second, third or fourth drug of the traditional anti-tumor type, such as taxol, cis-platinum or doxorubicin. A further advantage is that the administration of a MMP
25 inhibitor with a selective inhibition profile can permit the reduction in dosage of the drugs being administered to the patient. This is an especially important advantage given both the toxicities and dosing limits of traditional anti-tumor drugs.

30 A contemplated selective MMP inhibitor compound useful in a contemplated process can be administered to by various routes and provide adequate therapeutic blood levels of enzymatically active inhibitor. A

compound can be administered, for example, by the oral (IG, PO) or intravenous (IV) routes. Oral administration is advantageous if the patient is ambulatory, not hospitalized, physically able and sufficiently responsible to take drug at the required intervals. This is true even if the person is being treated with more than one drug for one or more diseases. On the other hand, IV drug administration is an advantage in a hospital setting wherein the dose and thus the blood levels can well controlled. A contemplated inhibitor can also be formulated for IM administration if desired. This route of administration can be desirable for the administration of prodrugs or regular drug delivery to patients that are either physically weak or have a poor compliance record or require constant drug blood levels.

Thus, in one embodiment, the present invention is directed to a treatment process that comprises administering a contemplated sulfamato hydroxamic acid metalloprotease inhibitor, or a pharmaceutically acceptable salt thereof, in an effective amount to a host mammal having a condition associated with pathological matrix metalloprotease activity. A contemplated sulfamato hydroxamate inhibitor compound useful in such a process inhibits the activity of one or both of MMP-2 and MMP-13, and exhibits substantially less inhibitory activity against at least MMP-1 in the *in vitro* assay noted above and discussed in detail hereinafter. A sulfamato hydroxamate inhibitor compound for use in a contemplated process corresponds in structure to

formula I, below, or a pharmaceutically acceptable salt thereof:



5

wherein

R^1 and R^2 are preferably taken together with the carbon to which they are bonded form a
 10 cycloalkyl or more preferably a heterocyclo group either of which is optionally substituted by one, two or three R^X substituents, or R^1 and R^2 are independently selected from the group consisting of:

hydrido,

15 an alkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

an alkyloxyalkyl group, optionally substituted with one, two or three groups independently selected
 20 from R^X substituents,

an alkylthioalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

an alkenyl group, optionally substituted with
 25 one, two or three groups independently selected from R^X substituents,

an alkynyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

an aryl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

an arylalkyl group, optionally substituted with
5 one, two or three groups independently selected from R^X substituents,

an arylalkyloxyalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

10 an aryloxyalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

an arylthioalkyl group, optionally substituted with one, two or three groups independently selected
15 from R^X substituents,

an arylalkylthioalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

a cycloalkyl or bicycloalkyl group, optionally
20 substituted with one, two or three groups independently selected from R^X substituents,

a cycloalkenyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

25 a cycloalkylalkyl or bicycloalkylalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

a cycloalkyloxyalkyl or bicycloalkyloxyalkyl group, optionally substituted with one, two or three
30 groups independently selected from R^X substituents,

- a cycloalkylalkyloxyalkyl or bicycloalkyloxyalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,
- 5 a cycloalkylthioalkyl or bicycloalkylthioalkyl group, optionally substituted with one, two or three groups independently selected from R^X;
- cycloalkylalkylthioalkyl or bicycloalkylalkylthioalkyl, optionally substituted
- 10 with one, two or three groups independently selected from R^X substituents,
- a heterocyclo group, optionally substituted with one, two or three groups independently selected from R^X substituents,
- 15 a heterocycloalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents
- a heteroaryl group, optionally substituted with one, two or three groups independently selected from
- 20 R^X substituents,
- a biarylalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,
- an arylalkenyl group, optionally substituted
- 25 with one, two or three groups independently selected from R^X substituents,
- an arylalkynyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

a heterocycloalkylthio group, optionally substituted with one, two or three groups selected independently from R^X substituents,

5 a heterocycloalkyloxyalkyl group, optionally substituted with one, two or three groups selected independently from R^X substituents,

a heteroarylalkenyl group, optionally substituted with one, two or three groups independently selected from R^X substituents, and

10 a heteroarylalkyloxyalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents;

wherein an R^X substituent is selected from the group consisting of a hydrido, aryl, heteroaryl, heterocyclo, aroyl, alkanoyl, heteroaroyl, halogen (F, Cl, Br, I), cyano, aldehydo (CHO, formyl), hydroxy, R^{CR^d} -amino ($-NR^{CR^d}$), R^{CR^d} -aminoalkyl, nitro, nitroso, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, alkoxy, aryloxy, heteroaryloxy, alkenyloxy, alkynyloxy, alkoxyaryl, alkoxyheteroaryl, alkoxyalkyl, R^C -oxyalkyl, alkoxyalkyl, alkylenedioxy, aryloxyalkyl, perfluoroalkyl, trifluoroalkyl, alkylthio, arylthio, alkyloxycarbonyl, alkyloxycarbonyloxy, aryloxycarbonyl, 25 arylalkyloxycarbonyl, alkyloxycarbonyl- R^C -amino, arylalkyloxycarbonyl- R^C -amino, aryloxycarbonyloxy, carboxy, R^{CR^d} -aminocarbonyloxy, R^{CR^d} -aminocarbonyl, R^{CR^d} -aminoalkanoyl, hydroxy- R^C -aminocarbonyl, R^{CR^d} -aminosulfonyl, arylsulfonyl (R^C) amino, R^{CR^d} - 30 aminoalkoxy, R^{CR^d} -aminocarbonyl (R^C) amino, trifluoromethylsulfonyl (R^C) amino, heteroarylsulfonyl-

(R^C)amino, alkylsulfonyl, arylsulfonyl(R^C)amino, arylsulfonyl(R^C)aminocarbonyl, alkylsulfonyl-(R^C)amino, arylcarbonyl(R^C)-aminosulfonyl, and an alkylsulfonyl(R^C)aminocarbonyl substituent;

- 5 wherein R^C and R^d are independently selected from the group consisting of a hydrido, alkanoyl, arylalkyl, aroyl, bisalkoxyalkyl, alkyl, haloalkyl, perfluoroalkyl, trifluoromethylalkyl, perfluoroalkoxyalkyl, alkoxyalkyl, cycloalkyl, 10 heterocycloalkyl, heterocycloalkylcarbonyl, aryl, heterocyclo, heteroaryl, cycloalkylalkyl, aryloxyalkyl, heteroaryloxyalkyl, heteroarylalkoxyalkyl, heteroarylthioalkyl, arylsulfonyl, alkylsulfonyl, heteroarylsulfonyl, 15 carboxyalkyl, alkoxycarbonylalkyl, aminocarbonyl, alkyliminocarbonyl, aryliminocarbonyl, heterocycloiminocarbonyl, arylthioalkyl, alkylthioalkyl, arylthioalkenyl, alkylthioalkenyl, heteroarylalkyl, haloalkanoyl, hydroxyalkanoyl, 20 thiolalkanoyl, alkenyl, alkynyl, alkoxyalkyl, alkoxycarbonyl, aryloxycarbonyl, aminoalkylcarbonyl, hydroxyalkyl, aminoalkyl, aminoalkylsulfonyl, aminosulfonyl wherein the amino nitrogen is (i) unsubstituted or (ii) independently substituted with 25 one or two R^Y radicals, or the substituents on the amino group taken together with the amino nitrogen form a saturated or partially unsaturated heterocyclo group optionally substituted with one, two or three groups independently selected from R^W substituents or 30 a heteroaryl group optionally substituted with one,

two or three groups independently selected from R^V substituents;

wherein R^Y is selected from the group consisting of an arylalkyl, aryl, heteroaryl, heterocyclo,
5 alkyl, alkynyl, alkenyl, alkoxyalkyl, alkoxyalkyl, substituted or unsubstituted aminoalkyl, alkyloxycarbonyl, arylalkyloxycarbonyl, carboxyalkyl, haloalkyl, alkanoyl, aroyl, substituted or unsubstituted aminoalkanoyl, halo alkanoyl and a
10 hydroxyalkyl group, each of which groups is optionally substituted by one or two groups independently selected from R^U substituents as are the substituents of the substituted aminoalkyl and substituted aminoalkanoyl groups;

15 wherein R^V is selected from the group consisting of a hydrido, aryl, heteroaryl, heterocyclo, aroyl, alkanoyl, heteroaroyl, halogen (F, Cl, Br, I); cyano, aldehydo (CHO, formyl), hydroxy, amino, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, alkoxy,
20 aryloxy, heteroaryloxy, alkenyloxy, alkynyloxy, alkoxyaryl, alkoxyheteroaryl, R^YR^Z-amino, alkoxyalkyl, alkylenedioxy, aryloxyalkyl, perfluoroalkyl, trifluoroalkyl, alkylthio, arylthio, alkyloxycarbonyl, alkyloxycarbonyloxy,
25 aryloxycarbonyl, arylalkyloxycarbonyl, arylalkyloxycarbonylamino, aryloxycarbonyloxy, carboxy, R^YR^Z-aminocarbonyloxy, R^YR^Z-aminocarbonyl, R^YR^Z-aminoalkanoyl, hydroxyaminocarbonyl, R^YR^Z-aminosulfonyl, R^YR^Z-aminocarbonyl (R^Y) amino,
30 trifluoromethylsulfonyl (R^Y) amino, heteroarylsulfonyl- (R^Y) amino, arylsulfonyl (R^Y) amino, arylsulfonyl (R^Y) -

aminocarbonyl, alkylsulfonyl(RY)amino, arylcarbonyl-(RY)aminosulfonyl, and an alkylsulfonyl(RY)-aminocarbonyl substituent;

wherein R^W is selected from the group consisting
5 of a hydrido, aryl, heteroaryl, heterocyclo, aroyl, alkanoyl, heteroaroyl, halogen (F, Cl, Br, I), cyano, aldehydo (CHO, formyl), hydroxy, amino, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, alkoxy, aryloxy, heteroaryloxy, alkenyloxy, alkynyloxy,
10 alkoxyaryl, alkoxyheteroaryl, RYR^Z-amino, alkoxyalkyl, alkylenedioxy, aryloxyalkyl, perfluoroalkyl, trifluoroalkyl, alkylthio, arylthio, alkyloxycarbonyl, alkyloxycarbonyloxy, aryloxycarbonyl, arylalkyloxycarbonyl,
15 arylalkyloxycarbonylamino, aryloxycarbonyloxy, carboxy, RYR^Z-aminocarbonyloxy, RYR^Z-aminocarbonyl, RYR^Z-aminoalkanoyl, hydroxyaminocarbonyl, RYR^Z-aminosulfonyl, RYR^Z-aminocarbonyl(RY)amino, trifluoromethylsulfonyl(RY)amino, heteroarylsulfonyl-(RY)amino, arylsulfonyl(RY)amino, arylsulfonyl(RY)-aminocarbonyl, alkylsulfonyl(RY)amino, arylcarbonyl-(RY)aminosulfonyl, and an alkylsulfonyl(RY)-aminocarbonyl substituent;

R^Z is selected from the group consisting of an
25 arylalkyl, aryl, heteroaryl, heterocyclo, alkyl, alkynyl, alkenyl, alkoxyalkyl, alkoxyalkylalkyl, substituted or unsubstituted aminoalkyl, alkyloxycarbonyl, arylalkyloxycarbonyl, carboxyalkyl, haloalkyl, alkanoyl, aroyl, substituted or
30 unsubstituted aminoalkanoyl, halo alkanoyl and a

hydroxyalkyl group, each of which groups are optionally substituted by one or two R^u substituents;

wherein R^u is selected from the group consisting of an arylalkyl, aryl, heteroaryl, heterocyclo,
5 alkyl, alkynyl, alkenyl, alkoxyalkyl, alkoxyalkyl, substituted or unsubstituted aminoalkyl, alkyloxycarbonyl, arylalkyloxycarbonyl, carboxyalkyl, haloalkyl, alkanoyl, aroyl, substituted or unsubstituted aminoalkanoyl, halo alkanoyl and a
10 hydroxyalkyl group, wherein the substituents of the substituted aminoalkyl and substituted aminoalkanoyl groups are selected from the group consisting of an alkyl, alkenyl, alkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, aryloxycarbonyl and an
15 alkyloxycarbonyl group;

R^{3a} and R^{3b} are independently selected from the group consisting of a hydrido, alkyl, alkenyl, alkynyl, aryl, heterocyclo, heteroaryl, cycloalkyl, and an alkoxyalkyl group, each of which groups is
20 optionally substituted by an -AREY substituent; in that AREY substituent, A is selected from the group consisting of

- (1) -O-;
- (2) -S-;
- 25 (3) -N(R^e)-;
- (4) -CO-N(R^e) or -N(R^e)-CO-;
- (5) -CO-O- or -O-CO-;
- (6) -O-CO-O-;
- (7) -HC=CH-;
- 30 (8) -NH-CO-NH-;
- (9) -C \equiv C-;
- (10) -NH-CO-O- or -O-CO-NH-;

-37-

- (11) -N=N-;
- (12) -NH-NH-;
- (13) -CS-N(R^e)- or -N(R^e)-CS-;
- (14) -CH₂-;
- 5 (15) -O-[(CH₂)₁₋₈]- or -[(CH₂)₁₋₈]O-; and
- (16) -S-CH₂- or -CH₂-S-; or
- (17) A is absent and R is directly
connected to R^{3a} or R^{3b}, or both R^{3a} and
R^{3b};
- 10 the moiety R is selected from the group
consisting of alkyl, alkoxyalkyl, aryl, heteroaryl,
cycloalkyl, heterocycloalkyl, aralkyl, heteroaralkyl,
heterocycloalkyl, cycloalkylalkyl, cycloalkoxyalkyl,
heterocycloalkoxyalkyl, aryloxyalkyl, heteroaryl-
15 oxyalkyl, arylthioalkyl, heteroarylthioalkyl,
cycloalkylthioalkyl, and a heterocycloalkylthioalkyl
group wherein the aryl, heteroaryl, cycloalkyl or
heterocycloalkyl substituent is (i) unsubstituted or
(ii) substituted with one or two radicals selected
20 from the group consisting of a halo, alkyl,
perfluoroalkyl, perfluoroalkoxy, perfluoroalkylthio,
trifluoromethylalkyl, amino, alkoxycarbonylalkyl,
alkoxy, C₁-C₂-alkylene-dioxy, hydroxycarbonylalkyl,
hydroxycarbonylalkylamino, nitro, hydroxy,
25 hydroxyalkyl, alkanoylamino, and a alkoxycarbonyl
group;
- the group E is selected from the group
consisting of
- (1) -CO(R^w)- or -(R^w)CO-;
- 30 (2) -CON(R^e)- or -(R^e)NCO-;
- (3) -CO-;

-38-

(4) $-\text{SO}_2-\text{R}^{\text{W}}$ or $-\text{R}^{\text{W}}\text{SO}_2-$;(5) $-\text{SO}_2-$;(6) $-\text{N}(\text{R}^{\text{e}})-\text{SO}_2-$ or $-\text{SO}_2-\text{N}(\text{R}^{\text{e}})-$; or

5 (7) E is absent and R is bonded directly
to Y; and

Y is absent or is selected from the group
consisting of a hydrido, alkyl, alkoxy, haloalkyl,
aryl, aralkyl, cycloalkyl, heteroaryl, hydroxy,
nitrile, nitro, aryloxy, aralkoxy, heteroaryloxy,
10 heteroaralkyl, R^{C} oxyalkyl, perfluoroalkoxy,
perfluoroalkylthio, trifluoromethylalkyl, alkenyl,
heterocycloalkyl, cycloalkyl, trifluoromethyl,
alkoxycarbonyl, and a aminoalkyl group, wherein the
aryl, heteroaryl or heterocycloalkyl group is (i)
15 unsubstituted or (ii) substituted with one or two
radicals independently selected from the group
consisting of an alkanoyl, halo, nitro, nitrile,
haloalkyl, alkyl, aralkyl, aryl, alkoxy, and an amino
group wherein the amino nitrogen is (i) unsubstituted
20 or (ii) substituted with one or two groups
independently selected from hydrido, alkyl, and an
aralkyl group;

wherein R^{e} is selected from hydrido, alkyl,
alkenyl, alkenyl, aryl, arylalkyl, heteroaryl,
25 heteroarylalkyl, aryloxycarbonyl, alkyloxycarbonyl,
 $\text{R}^{\text{C}}\text{R}^{\text{d}}$ amino carbonyl, $\text{R}^{\text{C}}\text{R}^{\text{d}}$ aminosulfonyl,
 $\text{R}^{\text{C}}\text{R}^{\text{d}}$ aminoalkanoyl and $\text{R}^{\text{C}}\text{R}^{\text{d}}$ aminoalkysulfonyl, and R^{C} ,
 R^{d} and R^{W} are as defined before; or

$\text{R}^{3\text{a}}$ and $\text{R}^{3\text{b}}$ taken together with the nitrogen atom
30 to which they are bonded form a group -GAREY (R^3)
wherein

-39-

G is a N-heterocyclo group;

the substituent A is selected from the group consisting of

- (1) -O-;
- 5 (2) -S-;
- (3) -NR^e-;
- (4) -CO-N(R^e) or -N(R^e)-CO-;
- (5) -CO-O- or -O-CO-;
- (6) -O-CO-O-;
- 10 (7) -HC=CH-;
- (8) -NH-CO-NH-;
- (9) -C≡C-;
- (10) -NH-CO-O- or -O-CO-NH-;
- (11) -N=N-;
- 15 (12) -NH-NH-;
- (13) -CS-N(R^e)- or -N(R^e)-CS-;
- (14) -CH₂-;
- (15) -O-[(CH₂)₁₋₈]- or -[(CH₂)₁₋₈]O-; and
- (16) -S-CH₂- or -CH₂-S-; or
- 20 (17) A is absent and R is directly connected to G;

the moiety R is selected from the group consisting of alkyl, alkoxyalkyl, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaralkyl, 25 heterocycloalkyl, cycloalkylalkyl, cycloalkoxyalkyl, heterocycloalkoxyalkyl, aryloxyalkyl, heteroaryl-oxyalkyl, arylthioalkyl, heteroarylthioalkyl, cycloalkylthioalkyl, and a heterocycloalkylthioalkyl group wherein the aryl or heteroaryl or cycloalkyl or 30 heterocycloalkyl substituent is (i) unsubstituted or (ii) substituted with one or two radicals selected

from the group consisting of a halo, alkyl, perfluoroalkyl, perfluoroalkoxy, perfluoroalkylthio, trifluoromethylalkyl, amino, alkoxycarbonylalkyl, alkoxy, C₁-C₂-alkylenedioxy, hydroxycarbonylalkyl, hydroxycarbonylalkylamino, nitro, hydroxy, hydroxyalkyl, alkanoylamino, and a alkoxycarbonyl group;


The moiety E is selected from the group consisting of

- 10 (1) -CO(R^W)- or -(R^W)CO-;
- (2) -CONH- or -HNCO-;
- (3) -CO-;
- (4) -SO₂-R^W- or -R^W-SO₂-;
- (5) -SO₂-;
- 15 (6) -NH-SO₂- or -SO₂-NH-; or
- (7) E is absent and R is bonded directly to Y; and

The moiety Y is absent or is selected from the group consisting of a hydrido, alkyl, alkoxy, haloalkyl, aryl, aralkyl, cycloalkyl, heteroaryl, hydroxy, aryloxy, aralkoxy, heteroaryloxy, heteroaralkyl, perfluoroalkoxy, perfluoroalkylthio, trifluoromethylalkyl, alkenyl, heterocycloalkyl, cycloalkyl, trifluoromethyl, alkoxycarbonyl, and a aminoalkyl group, wherein the aryl or heteroaryl or heterocycloalkyl group is (i) unsubstituted or (ii) substituted with one or two radicals independently selected from the group consisting of an alkanoyl, halo, nitro, aralkyl, aryl, alkoxy, and an amino group wherein the amino nitrogen is (i) unsubstituted or (ii) substituted with one or two groups

independently selected from hydrido, alkyl, and an aralkyl group.

More generally, a contemplated compound includes an inhibitor utilized as discussed above, as well as a pro-drug form of such a compound and also an intermediate used in the synthesis of a hydroxamate or hydroxamate pro-drug. Such a more general compound corresponds in structure to formula II, below,



wherein R^1 , R^2 , R^{3a} and R^{3b} are as before described with the above preferences, and


15 R^{20} is (a) $-O-R^{21}$, where R^{21} is selected from the group consisting of a hydrido, C_1 - C_6 -alkyl, aryl, ar- C_1 - C_6 -alkyl group and a pharmaceutically acceptable cation, (b) $-NH-O-R^{22}$, wherein R^{22} is a selectively removable protecting group such as a 2-
20 tetrahydropyranyl, benzyl, p-methoxybenzyl (MOZ) carbonyl- C_1 - C_6 -alkoxy, trisubstituted silyl group or o-nitrophenyl group, peptide synthesis resin and the like, wherein trisubstituted silyl group is substituted with C_1 - C_6 -alkyl, aryl, or ar- C_1 - C_6 -
25 alkyl, or (c) $-NH-O-R^{14}$, where R^{14} is hydrido, a pharmaceutically acceptable cation or $C(W)R^{15}$ where W is O or S and R^{15} is selected from the group consisting of an C_1 - C_6 -alkyl, aryl, C_1 - C_6 -alkoxy,

-42-

heteroaryl-C₁-C₆-alkyl, C₃-C₈-cycloalkyl-C₁-C₆-alkyl, aryloxy, ar-C₁-C₆-alkoxy, ar-C₁-C₆-alkyl, heteroaryl and amino C₁-C₆-alkyl group wherein the aminoalkyl nitrogen is (i) unsubstituted or (ii) substituted with one or two substituents independently selected from the group consisting of an C₁-C₆-alkyl, aryl, ar-C₁-C₆-alkyl, C₃-C₈-cycloalkyl-C₁-C₆-alkyl, ar-C₁-C₆-alkoxycarbonyl, C₁-C₆-alkoxycarbonyl, and C₁-C₆-alkanoyl radical, or (iii) wherein the amino C₁-C₆-alkyl nitrogen and two substituents attached thereto form a 5- to 8-membered heterocyclo or heteroaryl ring.

The substituent -NR^{3a}R^{3b} can also be referred to as a R³ group. One exemplary R³ group is -N(CH₃)₂, whereas another is the before-discussed substituent group -GAREY that is present in more preferred compounds as is discussed hereinbelow.


One group of more preferred compounds correspond ion structure to formula III, formula IIIA or a pharmaceutically acceptable salt thereof:



wherein substituents R¹, R², R²⁰ and -GAREY are as discussed before, with the before-described preferences.


Yet another more preferred group of contemplated compounds correspond ion structure to formula IV,

formula IVA or a pharmaceutically acceptable salt thereof:




5 wherein substituents R¹, R², R²⁰ and -AREY are as discussed before, with the before-described preferences.

A still more preferred group of contemplated compounds correspond ion structure to formula V,
10 formula VA or a pharmaceutically acceptable salt thereof:



15 wherein substituents R²⁰ and -AREY are as discussed before, with the before-described preferences.

Another more preferred group of contemplated compounds correspond ion structure to formula VI,
20 formula VIA or a pharmaceutically acceptable salt thereof:




25 wherein substituents R¹, R², R²⁰ and -EY are as discussed before, with the before-described

- 44 -


preferences, and A is -CH₂-, -O-CH₂-, -CH₂-O-, -S-CH₂- or -CH₂-S-.

A still more preferred group of contemplated compounds correspond ion structure to formula VII,
 5 formula VIIA or a pharmaceutically acceptable salt thereof:



wherein substituents R²⁰ and -EY are as
 10 discussed before as part of an -AREY or -GAREY group,
 , with the before-described preferences, and A is -CH₂-, -O-CH₂-, -CH₂-O-, -S-CH₂- or -CH₂-S-.

Another group of preferred compounds for use in
 a contemplated process has a structure that
 15 corresponds to formulas VIII and VIIIA, below, or a
 pharmaceutically acceptable salt thereof:



20

wherein

R^{3a}, R^{3b} and R²⁰ are as defined before, ,
 with the before-described preferences; and
 m is zero, 1 or 2;

-45-

n is zero, 1 or 2;


p is zero, 1 or 2;

the sum of $m + n + p = 1, 2, 3$ or 4;


(a) one of X, Y and Z is selected from the
 5 group consisting of $C(O)$, NR^6 , O, S, $S(O)$, $S(O)_2$ and
 $NS(O)_2R^7$, and the remaining two of X, Y and Z are
 CR^8R^9 , and $CR^{10}R^{11}$, or

(b) X and Z or Z and Y together constitute
 a moiety that is selected from the group consisting
 10 of $NR^6C(O)$, $NR^6S(O)$, $NR^6S(O)_2$, NR^6S , NR^6O , SS, NR^6NR^6
 and $OC(O)$, with the remaining one of X, Y and Z being
 CR^8R^9 , or

(c) n is zero and X, Y and Z together
 constitute a moiety selected from the group
 15 consisting of



-46-



wherein wavy lines are bonds to the atoms of the depicted ring;

R^6 and $R^{6'}$ are independently selected from the group consisting of hydrido, C_1 - C_6 -alkanoyl, C_6 -aryl- C_1 - C_6 -alkyl, aroyl, bis(C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl)- C_1 - C_6 -alkyl, C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl, C_1 - C_6 -perfluoroalkyl, C_1 - C_6 -trifluoromethylalkyl, C_1 - C_6 -perfluoroalkoxy- C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl, C_3 - C_6 -cycloalkyl, C_3 - C_8 -heterocycloalkyl, C_3 - C_8 -heterocycloalkylcarbonyl, C_6 -aryl, C_5 - C_6 -heterocyclo, C_5 - C_6 -heteroaryl, C_3 - C_8 -cycloalkyl- C_1 - C_6 -alkyl, C_6 -aryloxy- C_1 - C_6 -alkyl, heteroaryloxy- C_1 - C_6 -alkyl, heteroaryl- C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl, heteroarylthio- C_1 - C_6 -alkyl, C_6 -arylsulfonyl, C_1 - C_6 -alkylsulfonyl, C_5 - C_6 -heteroarylsulfonyl, carboxy- C_1 - C_6 -alkyl, C_1 - C_4 -alkoxycarbonyl- C_1 - C_6 -alkyl, aminocarbonyl, C_1 - C_6 -alkyliminocarbonyl, C_6 -aryliminocarbonyl, C_5 - C_6 -heterocycloiminocarbonyl,

C₆-arylthio-C₁-C₆-alkyl, C₁-C₆-alkylthio-C₁-C₆-alkyl, C₆-arylthio-C₃-C₆-alkenyl, C₁-C₄-alkylthio-C₃-C₆-alkenyl, C₅-C₆-heteroaryl-C₁-C₆-alkyl, halo-C₁-C₆-alkanoyl, hydroxy-C₁-C₆-alkanoyl, thiol-C₁-C₆-alkanoyl, C₃-C₆-alkenyl, C₃-C₆-alkynyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₅-alkoxycarbonyl, aryloxycarbonyl, NR⁸R⁹-C₁-C₅-alkylcarbonyl, hydroxy-C₁-C₅-alkyl, an aminocarbonyl wherein the aminocarbonyl nitrogen is (i) unsubstituted or (ii) substituted with one or two radicals independently selected from the group consisting of C₁-C₆-alkyl, ar-C₁-C₆-alkyl, C₃-C₈-cycloalkyl and a C₁-C₆-alkanoyl group, hydroxyaminocarbonyl, an aminosulfonyl group wherein the aminosulfonyl nitrogen is (i) unsubstituted or (ii) substituted with one or two radicals independently selected from the group consisting of C₁-C₆-alkyl, ar-C₁-C₆-alkyl, C₃-C₈-cycloalkyl and a C₁-C₆-alkanoyl group, an amino-C₁-C₆-alkylsulfonyl group wherein the amino-C₁-C₆-alkylsulfonyl nitrogen is (i) unsubstituted or (ii) substituted with one or two radicals independently selected from the group consisting of C₁-C₆-alkyl, ar-C₁-C₆-alkyl, C₃-C₈-cycloalkyl and a C₁-C₆-alkanoyl group and an amino-C₁-C₆-alkyl group wherein the aminoalkyl nitrogen is (i) unsubstituted or (ii) substituted with one or two radicals independently selected from the group consisting of C₁-C₆-alkyl, ar-C₁-C₆-alkyl, C₃-C₈-cycloalkyl and a C₁-C₆-alkanoyl group;

R^7 is selected from the group consisting of a arylalkyl, aryl, heteroaryl, heterocyclo, C_1 - C_6 -alkyl, C_3 - C_6 -alkynyl, C_3 - C_6 -alkenyl, C_1 - C_6 -carboxyalkyl and a C_1 - C_6 -hydroxyalkyl group;

- 5 R^8 and R^9 and R^{10} and R^{11} are independently selected from the group consisting of a hydrido, hydroxy, C_1 - C_6 -alkyl, aryl, ar- C_1 - C_6 -alkyl, heteroaryl, heteroar- C_1 - C_6 -alkyl, C_2 - C_6 -alkynyl, C_2 - C_6 -alkenyl, thiol- C_1 - C_6 -alkyl, C_1 - C_6 -alkylthio- C_1 - C_6 -
10 alkyl cycloalkyl, cycloalkyl- C_1 - C_6 -alkyl, heterocycloalkyl- C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl, aralkoxy- C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy- C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl, hydroxy- C_1 - C_6 -alkyl, hydroxycarbonyl- C_1 - C_6 -alkyl, hydroxycarbonylar- C_1 - C_6 -
15 alkyl, aminocarbonyl- C_1 - C_6 -alkyl, aryloxy- C_1 - C_6 -alkyl, heteroaryloxy- C_1 - C_6 -alkyl, arylthio- C_1 - C_6 -alkyl, heteroarylthio- C_1 - C_6 -alkyl, the sulfoxide or sulfone of any said thio substituents, perfluoro- C_1 - C_6 -alkyl, trifluoromethyl- C_1 - C_6 -alkyl, halo- C_1 - C_6 -
20 alkyl, alkoxycarbonylamino- C_1 - C_6 -alkyl and an amino- C_1 - C_6 -alkyl group wherein the aminoalkyl nitrogen is (i) unsubstituted or (ii) substituted with one or two radicals independently selected from the group consisting of C_1 - C_6 -alkyl, ar- C_1 - C_6 -alkyl, cycloalkyl
25 and C_1 - C_6 -alkanoyl, or wherein R^8 and R^9 or R^{10} and R^{11} and the carbon to which they are bonded form a carbonyl group, or wherein R^8 and R^9 or R^{10} and R^{11} , or R^8 and R^{10} together with the atoms to which they


are bonded form a 5- to 8-membered carbocyclic ring, or a 5- to 8-membered heterocyclic ring containing one or two heteroatoms that are nitrogen, oxygen, or sulfur, with the proviso that only one of R⁸ and R⁹
5 or R¹⁰ and R¹¹ is hydroxy;

R¹² and R^{12'} are independently selected from the group consisting of a hydrido, C₁-C₆-alkyl, aryl, ar-C₁-C₆-alkyl, heteroaryl, heteroaralkyl, C₂-C₆-alkynyl, C₂-C₆-alkenyl, thiol-C₁-C₆-alkyl,
10 cycloalkyl, cycloalkyl-C₁-C₆-alkyl, heterocycloalkyl-C₁-C₆-alkyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, aryloxy-C₁-C₆-alkyl, amino-C₁-C₆-alkyl, C₁-C₆-alkoxy-C₁-C₆-alkoxy-C₁-C₆-alkyl, hydroxy-C₁-C₆-alkyl, hydroxycarbonyl-C₁-C₆-alkyl, hydroxycarbonylar-C₁-C₆-alkyl,
15 aminocarbonyl-C₁-C₆-alkyl, aryloxy-C₁-C₆-alkyl, heteroaryloxy-C₁-C₆-alkyl, C₁-C₆-alkylthio-C₁-C₆-alkyl, arylthio-C₁-C₆-alkyl, heteroarylthio-C₁-C₆-alkyl, the sulfoxide or sulfone of any said thio substituents, perfluoro-C₁-C₆-alkyl, trifluoromethyl-C₁-C₆-alkyl, halo-C₁-C₆-alkyl, alkoxycarbonylamino-C₁-C₆-alkyl and an amino-C₁-C₆-alkyl group wherein the aminoalkyl nitrogen is (i) unsubstituted or (ii) substituted with one or two radicals independently selected from the group consisting of C₁-C₆-alkyl,
20 ar-C₁-C₆-alkyl, cycloalkyl and C₁-C₆-alkanoyl;

R¹³ is selected from the group consisting of a hydrido, benzyl, phenyl, C₁-C₆-alkyl, C₂-C₆-alkynyl, C₂-C₆-alkenyl and a C₁-C₆-hydroxyalkyl group.


A compound of formulas VIII and VIIIA thus includes a compound illustrated by formulas V, VA, VII and VIIA discussed above, as well as other compounds of formulas I and II.

5 A group of particularly preferred compounds of formula VIII and VIIIA correspond in structure to formula IX or IX, below, or a pharmaceutically acceptable salt thereof:



wherein R^6 , R^{20} and -AREY are as described before, with the before-described preferences.


15 A still more preferred group of contemplated compounds correspond in structure to formula X, formula XA or a pharmaceutically acceptable salt thereof:




wherein substituents R^6 , R^{20} and -EY are as discussed before as part of an -AREY or -GAREY group, with the before-described preferences, and wherein A is -CH₂-, -O-CH₂-, -CH₂-O-, -S-CH₂- or -CH₂-S-.

-51-

A group of contemplated compounds that is even still more preferred contain a $-NR^{3a}R^{3b}$ group in which R^{3a} and R^{3b} taken together with the nitrogen atom to which they are bonded form a group $-GAREY$ where G is a disubstituted piperazinyl group correspond in structure to formula XI, formula XIA or a pharmaceutically acceptable salt thereof:



10




wherein the definitions for X, Y, Z, m, n, p A, E, Y and R^{20} are as before discussed, with the before-described preferences and A being absent.

Without wishing to be bound by theory, it is believed that when substituent A is absent so that R is bonded directly to G, the bond flexibility provided by the non-sulfamido nitrogen of the piperazinyl group, $-G-$, to the remainder of the $-AREY$ substituent present provides enhanced fit of an inhibitor into the binding pocket of gelatinase and MMP-13 enzymes, while not appreciably altering the binding to MMP-1. It is also believed that similar

flexibility and enhanced binding to these enzymes is achieved where $-\text{SO}_2\text{G}-$ is a substituted N-sulfonamidopiperadinyll group and substituent A is a single atom such as $-\text{O}-$, $-\text{S}-$, or $-\text{CH}_2-$ or $-\text{NH}-$.

- 5 Of the compounds of formulas XI and XIA, a compound corresponding in formula to formulas XII or XIII, formulas XIIA or XIII A or a pharmaceutically acceptable salt thereof is yet more preferred,




- 15 wherein R^6 , R^{20} and $-\text{EY}$ are as described before, with the before-described preferences.

It is particularly preferred that both substituents A and E be absent in a compound of formulas XII, XIII, XIIA and XIII A so that the substituted phenyl ring, substituent or moiety R, is bonded on one side directly to one nitrogen atom of the piperazinyl ring, and on the other side, that phenyl ring is bonded directly to the Y group.


25 The structures of several particularly preferred compounds of the above formulas are shown

- 53 -


below along with the Example in which the particular compound is synthesized.




EXAMPLE 1




EXAMPLE 3




EXAMPLE 2




EXAMPLE 2H




EXAMPLE 4




EXAMPLE 5




EXAMPLE 6




EXAMPLE 6H



EXAMPLE 7



EXAMPLE 8



EXAMPLE 9

Further particularly preferred compounds include:

- 5 4-[(hydroxyamino)-carbonyl]-4-[[4-[4-(trifluoromethyl)-phenoxy]-1-piperidinyl]sulfonyl]-1-piperidinecarboxylate;
N-hydroxy-2-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-sulfonyl]acetamide;
- 10 N-[(tetrahydro-2H-pyran-2-yl)oxy]-2-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]acetamide;
N-hydroxy-1-(2-methoxyethyl)-4-[[4-[4-(trifluoromethyl)-phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide, monohydrochloride;
- 15 1-(2-Methoxyethyl)-N-[(tetrahydro-2H-pyran-2-yl)oxy]-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide;
tetrahydro-N-hydroxy-4-[[4-(4-nitrophenoxy)-1-piperidinyl]sulfonyl]-2H-pyran-4-carboxamide;
- 20 tetrahydro-N-hydroxy-4-[[4-(4-nitrophenoxy)-1-piperidinyl]sulfonyl]-2H-pyran-4-carboxamide;
N-hydroxy-1-(phenylmethyl)-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide, monohydrochloride;
- 25 1-(phenylmethyl)-N-[(tetrahydro-2H-pyran-2-yl)oxy]-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]-sulfonyl]-4-piperidinecarboxylate;

N-hydroxy-1-(2-methoxyethyl)-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide, monohydrochloride;

1-(2-methoxyethyl)-N-[(tetrahydro-2H-pyran-2-yl)oxy]-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxylate;

N-hydroxy-2-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]sulfonyl]acetamide;

N-[(tetrahydro-2H-pyran-2-yl)oxy]-2-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]sulfonyl]-acetamide;

tetrahydro-N-hydroxy-4-[[4-[4-(trifluoromethoxy)-phenoxy]-1-piperidinyl]sulfonyl]-2H-pyran-4-carboxamide;

tetrahydro-N-[(tetrahydro-2H-pyran-2-yl)oxy]-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]sulfonyl]-2H-pyran-4-carboxamide;

N-hydroxy-1-(phenylmethyl)-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-sulfonyl]-4-piperidinecarboxamide, monohydrochloride;

1-(phenylmethyl)-N-[(tetrahydro-2H-pyran-2-yl)oxy]-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide;

N-hydroxy-1-(2-pyridinylmethyl)-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide, monohydrochloride;

1-(2-pyridinylmethyl)-N-[(tetrahydro-2H-pyran-2-yl)oxy]-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]-sulfonyl]-4-piperidinecarboxamide;

N-hydroxy-1-(2-pyrimidinyl)-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide, monohydrochloride;

1-(2-pyrimidinyl)-N-[(tetrahydro-2H-pyran-2-yl)oxy]-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-sulfonyl]-4-piperidinecarboxamide;

N-hydroxy-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-1-[4-(trifluoromethyl)-2-pyrimidinyl]-4-piperidinecarboxamide,
5 monohydrochloride;

N-[(tetrahydro-2H-pyran-2-yl)oxy]-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-1-[4-(trifluoromethyl)-2-pyrimidinyl]-4-piperidinecarboxamide;
10

1-(5-ethyl-2-pyrimidinyl)-N-hydroxy-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide, monohydrochloride;

15 1-(5-ethyl-2-pyrimidinyl)-N-[(tetrahydro-2H-pyran-2-yl)oxy]-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide;

tetrahydro-N-hydroxy-4-[[4-[4-(trifluoromethoxy)-phenoxy]-1-piperidinyl]sulfonyl]-2H-thiopyran-4-carboxamide;
20

tetrahydro-N-hydroxy-4-[[4-[4-(trifluoromethoxy)-phenoxy]-1-piperidinyl]sulfonyl]-2H-thiopyran-4-carboxamide;

tetrahydro-N-hydroxy-4-[[4-[4-(trifluoromethoxy)-phenoxy]-1-piperidinyl]sulfonyl]-2H-thiopyran-4-carboxamide, 1,1-dioxide;
25

tetrahydro-N-hydroxy-4-[[4-[4-(trifluoromethoxy)-phenoxy]-1-piperidinyl]sulfonyl]-2H-thiopyran-4-carboxamide, 1,1-dioxide;

30 tetrahydro-N-hydroxy-4-[[4-[4-(trifluoromethyl)-phenoxy]-1-piperidinyl]sulfonyl]-2H-thiopyran-4-carboxamide;

tetrahydro-*N*-hydroxy-4-[[4-[4-(trifluoromethyl)-phenoxy]-1-piperidinyl]sulfonyl]-2*H*-thiopyran-4-carboxamide;

N-hydroxy-4[[1'-(*n*-pentyl)[4,4'-bipiperidin]-1-yl]sulfonyl]-tetrahydro-2*H*-pyran-4-carboxamide;

N-hydroxy-4[[1'-(4-methoxybenzoyl)[4,4'-bipiperidin]-1-yl]sulfonyl]-tetrahydro-2*H*-pyran-4-carboxamide;

N-hydroxy-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide, monohydrochloride;

1-(2-furanylmethyl)-*N*-hydroxy-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide;

4-[[4-[4-[4-(trifluoromethyl)phenoxy]phenoxy]-1-piperidinyl]sulfonyl]tetrahydro-*N*-hydroxy-2*H*-pyran-4-carboxamide;

tetrahydro-*N*-hydroxy-4-[[4-(4-pentylphenyl)-1-piperazinyl]sulfonyl]-2*H*-pyran-4-carboxamide, monohydrochloride;

tetrahydro-*N*-hydroxy-4-[(4-phenyl-1-piperazinyl)-sulfonyl]-2*H*-pyran-4-carboxamide;

N-hydroxy-1-(2-methoxyethyl)-4-[[4-[[4-(trifluoromethyl)benzoyl]amino]-1-piperidinyl]-sulfonyl]-4-piperidinecarboxamide, monohydrochloride;

N-hydroxy-1-phenyl-4-[[4-[4-(trifluoromethoxy)-phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide, monohydrochloride;

N-hydroxy-1-phenyl-4-[[4-[4-(trifluoromethyl)-phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide, monohydrochloride;

- 4-[[4-[4-(1,1-dimethylethyl)phenyl]-1-piperazinyl]-sulfonyl]-N-hydroxy-1-(2-methoxyethyl)-4-piperidinecarboxamide, monohydrochloride;
- 4-[[4-(4-butoxyphenyl)-1-piperazinyl]sulfonyl]-N-hydroxy-1-(2-methoxyethyl)-4-piperidinecarboxamide, dihydrochloride;
- tetrahydro-N-hydroxy-4-[[4-[[4-[(trifluoromethyl)-thio]phenyl]-thio]-1-piperidinyl]sulfonyl]-2H-pyran-4-carboxamide;
- 4-[[4-(4-bromophenyl)-4-fluoro-1-piperidinyl]-sulfonyl]tetrahydro-N-hydroxy-2H-pyran-4-carboxamide;
- 4-[[4-[4-(3,5-dimethylphenoxy)phenoxy]-1-piperidinyl]sulfonyl]tetrahydro-N-hydroxy-2H-pyran-4-carboxamide;
- 1-cyclopropyl-N-hydroxy-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide, monohydrochloride;
- N-hydroxy-1-(iminophenylmethyl)-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide, monohydrochloride;
- N-hydroxy-1-[(4-hydroxyphenyl)iminomethyl]-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide, monohydrochloride;
- 1-(2-furanylcarbonyl)-N-hydroxy-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide;
- N-hydroxy-1-[2-(methylthio)-4-pyrimidinyl]-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide, monohydrochloride;

- 1-cyclopropyl-N-hydroxy-4-[[4-[4-(tri-fluoromethoxy)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide, monohydrochloride;
- N-hydroxy-4-[[1'-(2-methoxyphenyl)[4,4'-bipiperidin]-1-yl]sulfonyl]-1-(phenylmethyl)-4-piperidinecarboxamide, dihydrochloride;
- 4-(1,4-dioxo-8-azaspiro-[4.5]dec-8-ylsulfonyl)-tetrahydro-N-hydroxy-2H-pyran-4-carboxamide;
- 4-[[4-[(3R,5R)-rel-3,5-dimethyl-1-piperidinyl]-carbonyl]-1-piperidinyl]sulfonyl]tetrahydro-N-hydroxy-2H-pyran-4-carboxamide;
- 4-[[4-[(3R,5S)-rel-3,5-dimethyl-1-piperidinyl]-carbonyl]-1-piperidinyl]sulfonyl]tetrahydro-N-hydroxy-2H-pyran-4-carboxamide;
- N-hydroxy-1-(4-methylphenyl)-4-[[4-[4-(trifluoro-methyl)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide, monohydrochloride;
- N-hydroxy-1-(4-methylphenyl)-4-[[4-[4-(trifluoro-methoxy)phenoxy]-1-piperidinyl]-sulfonyl]-4-piperidinecarboxamide, monohydrochloride;
- tetrahydro-N-hydroxy-4-[[4-(phenylmethyl)-1-piperazinyl]-sulfonyl]-2H-pyran-4-carboxamide, monohydrochloride;
- N-hydroxy-1-(phenylmethyl)-4-[(4-phenyl-1-piperazinyl)sulfonyl]-4-piperidinecarboxamide, bis(trifluoroacetate);
- N-hydroxy-1-(phenylmethyl)-4-[(4-phenyl-1-piperazinyl)sulfonyl]-4-piperidinecarboxamide, dihydrochloride; and
- 4-[[4-(4-butoxy-3-methylphenyl)-1-piperazinyl]-sulfonyl]-tetrahydro-N-hydroxy-2H-pyran-4-carboxamide.

-60-

Most preferred are compounds or their salts are those of Examples 19, 51, 47, 17, 42, 15, 12, 14, 35, 32, 23, 3, 2, 36, 43, 44, 13, 6, 46, 28, 30, 10, 50, 9, 18, 31, 16, 8, 55, 11, 38, 53, 33, 41, 40, 4, 5 54, 34 and 5.

Table 1 through Table 221, below, illustrate several compounds useful in a process of this invention. Each group of compounds is illustrated by a generic formula, or formulae, 10 followed by a series of preferred moieties or groups that constitute various substituents that can be attached at the position clearly shown in the generic structure. The generic symbols, e.g., R^1 , R^2 and the like, are as shown in the tables and are not 15 necessarily as defined before. This system is well known in the chemical communication arts and is widely used in scientific papers and presentations. For example in Table 1, R^1 and R^2 groups of the generic structure shown and of formula I are 20 illustrated as being taken together with the carbon to which they are bonded illustrate structural variables that can substitute for the R^1 and R^2 groups shown in the balance of the table. There are 12 R^1 and R^2 groups shown that are used to represent, 25 in a non-limiting manner, 12 distinct compounds that can be prepared for use in the invention.

- 61 -

Table 1



















Table 2



Tabl 3



Tabl 4




 R^3 

Table 5

 $\sim R^3$

1 	9 	16
2 	10 	17
3 	11 	18
4 	12 	19
5 	13 	20
6 	14 	21
7 	15 	22
8 		

Table 6





Table 7



 R^3 

Table 8






Table 9

 R^3

1 	8 	15
2 	9 	16
3 	10 	17
4 	11 	18
5 	12 	19
6 	13 	20
7 	14 	21

Table 10



 R^3 

Table 11






Table 12



1 	9 	16
2 	10 	17
3 	11 	18
4 	12 	19
5 	13 	20
6 	14 	21
7 	15 	22
8 		

Table 13





Table 14





Table 15





Table 16

























Table 17












































Table 18



Tabl 19


























Table 20





Table 21





Table 22















Table 23





Table 24



 R^3 

Table 25








Table 26



1 	9 	16
2 	10 	17
3 	11 	18
4 	12 	19
5 	13 	20
6 	14 	21
7 	15 	22
8 		

Table 27



Tabl 28





Table 29















Table 30



 R^3 

Table 31




 R^3 

Table 32

 R^3

1 	9 	16
2 	10 	17
3 	11 	18
4 	12 	19
5 	13 	20
6 	14 	21
7 	15 	22
8 		

Table 33





Table 34





Table 35





Table 36

























Table 37





Table 38






Table 39



1 	9 	16
2 	10 	17
3 	11 	18
4 	12 	19
5 	13 	20
6 	14 	21
7 	15 	22
8 		

Table 40





Table 41





Table 42















Table 43





Table 44






















Table 45





Table 46





Table 47





Table 48

















Table 49



-110-

Table 50

























Table 51





Table 52





Table 53





Table 54





Table 55

















Table 56



Tabl 57





Table 58





Table 59

























Table 60



1 	9 	16
2 	10 	17
3 	11 	18
4 	12 	19
5 	13 	20
6 	14 	21
7 	15 	22
8 		

Table 61





Table 62





Table 63















Table 64





Table 65






















Table 66




 R^3 

Table 67



1 	9 	16
2 	10 	17
3 	11 	18
4 	12 	19
5 	13 	20
6 	14 	21
7 	15 	22
8 		

Table 68



 R^3 

Table 69





Table 70
















Table 71



1 	8 	15
2 	9 	16
3 	10 	17
4 	11 	18
5 	12 	19
6 	13 	20
7 	14 	21

Tabl 72






















Table 73





Table 74


























Table 75





Table 76





Table 77















Table 78





Table 79



 R^3 

Table 80





Table 81



 R^3 

Table 82





Table 83


















Table 84



Tabl 85



1	8	15
2	9	16
3	10	17
4	11	18
5	12	19
6	13	20
7	14	21

Table 86





Table 87

























Table 88





Table 89





Table 90





Table 91
















Table 92



8	15
9	16
10	17
11	18
12	19
13	20
14	21

Table 93





Table 94





Table 95



 R^3 

Table 96





Table 97





Table 98
















Table 99



1 	8 	15
2 	9 	16
3 	10 	17
4 	11 	18
5 	12 	19
6 	13 	20
7 	14 	21

Table 100





Table 101


























Table 102



1 	9 	16
2 	10 	17
3 	11 	18
4 	12 	19
5 	13 	20
6 	14 	21
7 	15 	22
8 		

Table 103





Table 104





Table 105















Table 106







Table 107



Tabl 108





Table 109





Table 110





Table 111





Table 112





Table 113





Table 114





Table 115






Table 116



1 	9 	16
2 	10 	17
3 	11 	18
4 	12 	19
5 	13 	20
6 	14 	21
7 	15 	22
8 		

Table 117





Table 118















Table 119





Table 120





Table 121


























Table 122



1 	9 	16
2 	10 	17
3 	11 	18
4 	12 	19
5 	13 	20
6 	14 	21
7 	15 	22
8 		

Table 123





Table 124





Table 125















Table 126





Table 127





Table 128

























Table 129




Table 130




Table 131





Table 132



- | | | |
|---|----|----|
| 1 | 10 | 17 |
| 2 | 11 | 18 |
| 3 | 12 | 19 |
| 4 | 13 | 20 |
| 5 | 14 | 21 |
| 6 | 15 | 22 |
| 7 | 16 | 23 |
| 8 | | 24 |
| 9 | | 25 |
| | | 26 |

Table 133





Table 134




Table 135





Table 136





Table 137






Table 138



1		9	
2		10	
3		11	
4		12	
5		13	
6		14	
7		15	
8		16	

Table 139





Table 140





Table 141





Table 142





Table 143







Table 144



-205-

Table 145







Table 146



-207-

Table 147






























Table 148



1		7	
2		8	
3		9	
4		10	
5		11	
6		12	

Table 149



-210-

Table 150
















Table 151





Table 152




 R^3 

Table 153

 R^3

1 	9 	16
2 	10 	17
3 	11 	18
4 	12 	19
5 	13 	20
6 	14 	21
7 	15 	22
8 		

Table 154





Table 155







 $\sim R^3$ 

Table 156



-217-

Table 157

 R^3 

Tabl 158





Table 159



 R^3 

Table 160



 R^3 

Table 161





Table 162








Table 163




- 224 -

Table 164




1 	8 	15
2 	9 	16
3 	10 	17
4 	11 	18
5 	12 	19
6 	13 	20
7 	14 	21

Table 165

R¹


1	-CH ₃	16		30	
2	-CH ₂ CH ₃	17		31	
3	-CH(CH ₃) ₂	18		32	
4	(-C ₆ H ₅)	19		33	
5		20		34	
6		21		35	
7		22		36	
8	-CH ₂ -N(CH ₃) ₂	23		37	
9	-CH ₂ -N(CH ₃)CH ₂ C ₆ H ₅	24		38	
10	(-C ₆ H ₁₁)	25		39	
11		26		40	
12		27		41	
13		28		42	
14	-CH ₂ CH=CHCH ₃	29		43	
15		30			

Table 166




1	$-\text{CH}_3$	16		30	
2	$-\text{CH}_2\text{CH}_3$	17		31	
3	$-\text{CH}(\text{CH}_3)_2$	18	$-\text{C}\equiv\text{CH}$	32	
4	($-\text{C}_6\text{H}_5$)	19	$-\text{C}\equiv\text{CCH}_3$	33	
5		20		34	
6		21	$-\text{CH}_2\text{OCH}_3$	35	
7		22	$-\text{CH}_2\text{OC}_6\text{H}_5$	36	
8	$-\text{CH}_2-\text{N}(\text{CH}_3)_2$	23		37	
9	$-\text{CH}_2-\text{N}(\text{CH}_3)\text{CH}_2\text{C}_6\text{H}_5$	24		38	
10	($-\text{C}_6\text{H}_{11}$)	25		39	
11		26		40	
12		27		41	
13		28	$-\text{CH}_2\text{OH}$	42	
14	$-\text{CH}_2\text{CH}=\text{CHCH}_3$	29		43	$-\text{H}$
15		30			

Table 167

R¹


1	-CH ₃	16		30	
2	-CH ₂ CH ₃	17		31	
3	-CH(CH ₃) ₂	18		32	
4	(-C ₆ H ₅)	19		33	
5		20		34	
6		21		35	
7		22		36	
8	-CH ₂ -N(CH ₃) ₂	23		37	
9	-CH ₂ -N(CH ₃)CH ₂ C ₆ H ₅	24		38	
10	(-C ₆ H ₁₁)	25		39	
11		26		40	
12		27		41	
13		28		42	
14	-CH ₂ CH=CHCH ₃	29		43	
15		30			

Table 168

R¹


1	-CH ₃	16		30	
2	-CH ₂ CH ₃	17		31	
3	-CH(CH ₃) ₂	18		32	
4	(-C ₆ H ₅)	19		33	
5		20		34	
6		21		35	
7		22		36	
8	-CH ₂ -N(CH ₃) ₂	23		37	
9	-CH ₂ -N(CH ₃)CH ₂ C ₆ H ₅	24		38	
10	(-C ₆ H ₁₁)	25		39	
11		26		40	
12		27		41	
13		28		42	
14	-CH ₂ CH=CHCH ₃	29		43	
15		30			

Table 169




1	$-\text{CH}_3$	16		30	
2	$-\text{CH}_2\text{CH}_3$	17		31	
3	$-\text{CH}(\text{CH}_3)_2$	18	$-\text{C}\equiv\text{CH}$	32	
4		19	$-\text{C}\equiv\text{CCH}_3$	33	
5		20		34	
6		21	$-\text{OCH}_3$	35	
7		22	$-\text{OC}_6\text{H}_5$	36	
8	$-\text{CH}_2-\text{N}(\text{CH}_3)_2$	23		37	
9	$-\text{CH}_2-\text{N}(\text{CH}_3)\text{CH}_2\text{C}_6\text{H}_5$	24		38	
10		25		39	
11		26		40	
12		27		41	
13		28	$-\text{OH}$	42	
14	$-\text{CH}_2\text{CH}=\text{CHCH}_3$	29		43	
15		30			

Table 170




1	$-\text{CH}_3$	16		30	
2	$-\text{CH}_2\text{CH}_3$	17		31	
3	$-\text{CH}(\text{CH}_3)_2$	18	$-\text{C}\equiv\text{CH}$	32	
4	($-\text{C}_6\text{H}_5$)	19	$-\text{C}\equiv\text{CCH}_3$	33	
5		20		34	
6		21	$-\text{OCH}_3$	35	
7		22	$-\text{OC}_6\text{H}_5$	36	
8	$-\text{CH}_2\text{-N}(\text{CH}_3)_2$	23		37	
9	$-\text{CH}_2\text{-N}(\text{CH}_3)\text{CH}_2\text{C}_6\text{H}_5$	24		38	
10	($-\text{C}_6\text{H}_{11}$)	25		39	
11		26		40	
12		27		41	
13		28	$-\text{OH}$	42	
14	$-\text{CH}_2\text{CH}=\text{CHCH}_3$	29		43	
15		30			

Table 171




1	—CH ₃	16		30	
2	—CH ₂ CH ₃	17		31	
3	—CH(CH ₃) ₂	18		32	
4	(-C ₆ H ₅)	19		33	
5		20		34	
6		21		35	
7		22		36	
8	—CH ₂ -N(CH ₃) ₂	23		37	
9	—CH ₂ -N(CH ₃)CH ₂ C ₆ H ₅	24		38	
10	(-C ₆ H ₁₁)	25		39	
11		26		40	
12		27		41	
13		28		42	
14	—CH ₂ CH=CHCH ₃	29		43	
15		30			

Table 172




1	$-\text{CH}_3$	16		30	
2	$-\text{CH}_2\text{CH}_3$	17		31	
3	$-\text{CH}(\text{CH}_3)_2$	18	$-\text{C}\equiv\text{CH}$	32	
4	($-\text{C}_6\text{H}_5$)	19	$-\text{C}\equiv\text{CCH}_3$	33	
5		20		34	
6		21	$-\text{OCH}_3$	35	
7		22	$-\text{OC}_6\text{H}_5$	36	
8	$-\text{CH}_2\text{N}(\text{CH}_3)_2$	23		37	
9	$-\text{CH}_2\text{N}(\text{CH}_3)\text{CH}_2\text{C}_6\text{H}_5$	24		38	
10	($-\text{C}_6\text{H}_{11}$)	25		39	
11		26		40	
12		27		41	
13		28	$-\text{OH}$	42	
14	$-\text{CH}_2\text{CH}=\text{CHCH}_3$	29		43	$-\text{H}$
15		30			

Table 173




1	$-\text{CH}_3$	16		30	
2	$-\text{CH}_2\text{CH}_3$	17		31	
3	$-\text{CH}(\text{CH}_3)_2$	18	$-\text{C}\equiv\text{CH}$	32	
4	($-\text{C}_6\text{H}_5$)	19	$-\text{C}\equiv\text{CCH}_3$	33	
5		20		34	
6		21	$-\text{OCH}_3$	35	
7		22	$-\text{OC}_6\text{H}_5$	36	
8	$-\text{CH}_2\text{N}(\text{CH}_3)_2$	23		37	
9	$-\text{CH}_2\text{N}(\text{CH}_3)\text{CH}_2\text{C}_6\text{H}_5$	24		38	
10		25		39	
11		26		40	
12		27		41	
13		28	$-\text{OH}$	42	
14	$-\text{CH}_2\text{CH}=\text{CHCH}_3$	29		43	$-\text{H}$
15		30			

Table 174




1	$-\text{CH}_3$	16		30	
2	$-\text{CH}_2\text{CH}_3$	17		31	
3	$-\text{CH}(\text{CH}_3)_2$	18	$-\text{C}\equiv\text{CH}$	32	
4		19	$-\text{C}\equiv\text{CCH}_3$	33	
5		20		34	
6		21	$-\text{OCH}_3$	35	
7		22	$-\text{OC}_6\text{H}_5$	36	
8	$-\text{CH}_2\text{N}(\text{CH}_3)_2$	23		37	
9	$-\text{CH}_2\text{N}(\text{CH}_3)\text{CH}_2\text{C}_6\text{H}_5$	24		38	
10		25		39	
11		26		40	
12		27		41	
13		28	$-\text{OH}$	42	
14	$-\text{CH}_2\text{CH}=\text{CHCH}_3$	29		43	
15		30			

Table 175

R¹

1	—CH ₃	16		30	
2	—CH ₂ CH ₃	17		31	
3	—CH(CH ₃) ₂	18		32	
4	(—C ₆ H ₅)	19		33	
5		20		34	
6		21		35	
7		22		36	
8	—CH ₂ -N(CH ₃) ₂	23		37	
9	—CH ₂ -N(CH ₃)CH ₂ C ₆ H ₅	24		38	
10	(—C ₆ H ₁₁)	25		39	
11		26		40	
12		27		41	
13		28		42	
14	—CH ₂ CH=CHCH ₃	29		43	
15		30			


Table 176



1	$-\text{CH}_3$	16		30	
2	$-\text{CH}_2\text{CH}_3$	17		31	
3	$-\text{CH}(\text{CH}_3)_2$	18	$-\text{C}\equiv\text{CH}$	32	
4	($-\text{C}_6\text{H}_5$)	19	$-\text{C}\equiv\text{CCH}_3$	33	
5		20		34	
6		21	$-\text{OCH}_3$	35	
7		22	$-\text{OC}_6\text{H}_5$	36	
8	$-\text{CH}_2-\text{N}(\text{CH}_3)_2$	23		37	
9	$-\text{CH}_2-\text{N}(\text{CH}_3)\text{CH}_2\text{C}_6\text{H}_5$	24		38	
10	($-\text{C}_6\text{H}_{11}$)	25		39	
11		26		40	
12		27		41	
13		28	$-\text{OH}$	42	
14	$-\text{CH}_2\text{CH}=\text{CHCH}_3$	29		43	$-\text{H}$
15		30			


-237-

Table 177




1	$-\text{CH}_3$	16		30	
2	$-\text{CH}_2\text{CH}_3$	17		31	
3	$-\text{CH}(\text{CH}_3)_2$	18	$-\text{C}\equiv\text{CH}$	32	
4	($-\text{C}_6\text{H}_5$)	19	$-\text{C}\equiv\text{CCH}_3$	33	
5		20		34	
6		21	$-\text{OCH}_3$	35	
7		22	$-\text{OC}_6\text{H}_5$	36	
8	$-\text{CH}_2\text{N}(\text{CH}_3)_2$	23		37	
9	$-\text{CH}_2\text{N}(\text{CH}_3)\text{CH}_2\text{C}_6\text{H}_5$	24		38	
10	($-\text{C}_6\text{H}_{11}$)	25		39	
11		26		40	
12		27		41	
13		28	$-\text{OH}$	42	
14	$-\text{CH}_2\text{CH}=\text{CHCH}_3$	29		43	$-\text{H}$
15		30			

Table 178




1	$-\text{CH}_3$	16		30	
2	$-\text{CH}_2\text{CH}_3$	17		31	
3	$-\text{CH}(\text{CH}_3)_2$	18	$-\text{C}\equiv\text{CH}$	32	
4	($-\text{C}_6\text{H}_5$)	19	$-\text{C}\equiv\text{CCH}_3$	33	
5		20		34	
6		21	$-\text{OCH}_3$	35	
7		22	$-\text{OC}_6\text{H}_5$	36	
8	$-\text{CH}_2\text{N}(\text{CH}_3)_2$	23		37	
9	$-\text{CH}_2\text{N}(\text{CH}_3)\text{CH}_2\text{C}_6\text{H}_5$	24		38	
10	($-\text{C}_6\text{H}_{11}$)	25		39	
11		26		40	
12		27		41	
13		28	$-\text{OH}$	42	
14	$-\text{CH}_2\text{CH}=\text{CHCH}_3$	29		43	
15		30			

Table 179




1	$-\text{CH}_3$	16		30	
2	$-\text{CH}_2\text{CH}_3$	17		31	
3	$-\text{CH}(\text{CH}_3)_2$	18	$-\text{C}\equiv\text{CH}$	32	
4	($-\text{C}_6\text{H}_5$)	19	$-\text{C}\equiv\text{CCH}_3$	33	
5		20		34	
6		21	$-\text{OCH}_3$	35	
7		22	$-\text{OC}_6\text{H}_5$	36	
8	$-\text{CH}_2-\text{N}(\text{CH}_3)_2$	23		37	
9	$-\text{CH}_2-\text{N}(\text{CH}_3)\text{CH}_2\text{C}_6\text{H}_5$	24		38	
10	($-\text{C}_6\text{H}_{11}$)	25		39	
11		26		40	
12		27		41	
13		28	$-\text{OH}$	42	
14	$-\text{CH}_2\text{CH}=\text{CHCH}_3$	29		43	
15		30			

Table 180

R¹


1	-CH ₃	16		30	
2	-CH ₂ CH ₃	17		31	
3	-CH(CH ₃) ₂	18		32	
4	(-C ₆ H ₅)	19		33	
5		20		34	
6		21		35	
7		22		36	
8	-CH ₂ -N(CH ₃) ₂	23		37	
9	-CH ₂ -N(CH ₃)CH ₂ C ₆ H ₅	24		38	
10		25		39	
11		26		40	
12		27		41	
13		28		42	
14	-CH ₂ CH=CHCH ₃	29		43	
15		30			

Table 181



1	$-\text{CH}_3$	16		30	
2	$-\text{CH}_2\text{CH}_3$	17		31	
3	$-\text{CH}(\text{CH}_3)_2$	18	$-\text{C}\equiv\text{CH}$	32	
4		19	$-\text{C}\equiv\text{CCH}_3$	33	
5		20		34	
6		21	$-\text{OCH}_3$	35	
7		22	$-\text{OC}_6\text{H}_5$	36	
8	$-\text{CH}_2\text{N}(\text{CH}_3)_2$	23		37	
9	$-\text{CH}_2\text{N}(\text{CH}_3)\text{CH}_2\text{C}_6\text{H}_5$	24		38	
10		25		39	
11		26		40	
12		27		41	
13		28	$-\text{OH}$	42	
14	$-\text{CH}_2\text{CH}=\text{CHCH}_3$	29		43	$-\text{H}$
15		30			

Table 182














1	$-\text{CH}_3$	16		30	
2	$-\text{CH}_2\text{CH}_3$	17		31	
3	$-\text{CH}(\text{CH}_3)_2$	18	$-\text{C}\equiv\text{CH}$	32	
4	 ($-\text{C}_6\text{H}_5$)	19	$-\text{C}\equiv\text{CCH}_3$	33	
5		20		34	
6		21	$-\text{CH}_2\text{OCH}_3$	35	
7		22	$-\text{CH}_2\text{OC}_6\text{H}_5$	36	
8	$-\text{CH}_2\text{N}(\text{CH}_3)_2$	23		37	
9	$-\text{CH}_2\text{N}(\text{CH}_3)\text{CH}_2\text{C}_6\text{H}_5$	24		38	
10	 ($-\text{C}_6\text{H}_{11}$)	25		39	
11		26		40	
12		27		41	
13		28	$-\text{CH}_2\text{OH}$	42	
14	$-\text{CH}_2\text{CH}=\text{CHCH}_3$	29		43	
15		30			

Table 183




1	$-\text{CH}_3$	16		30	
2	$-\text{CH}_2\text{CH}_3$	17		31	
3	$-\text{CH}(\text{CH}_3)_2$	18	$-\text{C}\equiv\text{CH}$	32	
4	($-\text{C}_6\text{H}_5$)	19	$-\text{C}\equiv\text{CCH}_3$	33	
5		20		34	
6		21	$-\text{OCH}_3$	35	
7		22	$-\text{OC}_6\text{H}_5$	36	
8	$-\text{CH}_2\text{N}(\text{CH}_3)_2$	23		37	
9	$-\text{CH}_2\text{N}(\text{CH}_3)\text{CH}_2\text{C}_6\text{H}_5$	24		38	
10	($-\text{C}_6\text{H}_{11}$)	25		39	
11		26		40	
12		27		41	
13		28	$-\text{OH}$	42	
14	$-\text{CH}_2\text{CH}=\text{CHCH}_3$	29		43	$-\text{H}$
15		30			

Table 184




1	$-\text{CH}_3$	16		30	
2	$-\text{CH}_2\text{CH}_3$	17		31	
3	$-\text{CH}(\text{CH}_3)_2$	18	$-\text{C}\equiv\text{CH}$	32	
4		19	$-\text{C}\equiv\text{CCH}_3$	33	
5		20		34	
6		21	$-\text{OCH}_3$	35	
7		22	$-\text{OC}_6\text{H}_5$	36	
8	$-\text{CH}_2-\text{N}(\text{CH}_3)_2$	23		37	
9	$-\text{CH}_2-\text{N}(\text{CH}_3)\text{CH}_2\text{C}_6\text{H}_5$	24		38	
10		25		39	
11		26		40	
12		27		41	
13		28	$-\text{OH}$	42	
14	$-\text{CH}_2\text{CH}=\text{CHCH}_3$	29		43	$-\text{H}$
15		30			

Table 185




1	$-\text{CH}_3$	16		30	
2	$-\text{CH}_2\text{CH}_3$	17		31	
3	$-\text{CH}(\text{CH}_3)_2$	18	$-\text{C}\equiv\text{CH}$	32	
4		19	$-\text{C}\equiv\text{CCH}_3$	33	
5		20		34	
6		21	$-\text{OCH}_3$	35	
7		22	$-\text{OC}_6\text{H}_5$	36	
8	$-\text{CH}_2\text{-N}(\text{CH}_3)_2$	23		37	
9	$-\text{CH}_2\text{-N}(\text{CH}_3)\text{CH}_2\text{C}_6\text{H}_5$	24		38	
10		25		39	
11		26		40	
12		27		41	
13		28	$-\text{OH}$	42	
14	$-\text{CH}_2\text{CH}=\text{CHCH}_3$	29		43	$-\text{H}$
15		30			

Table 186




1	$-\text{CH}_3$	16		30	
2	$-\text{CH}_2\text{CH}_3$	17		31	
3	$-\text{CH}(\text{CH}_3)_2$	18	$-\text{C}\equiv\text{CH}$	32	
4	($-\text{C}_6\text{H}_5$)	19	$-\text{C}\equiv\text{CCH}_3$	33	
5		20		34	
6		21	$-\text{OCH}_3$	35	
7		22	$-\text{OC}_6\text{H}_5$	36	
8	$-\text{CH}_2-\text{N}(\text{CH}_3)_2$	23		37	
9	$-\text{CH}_2-\text{N}(\text{CH}_3)\text{CH}_2\text{C}_6\text{H}_5$	24		38	
10	($-\text{C}_6\text{H}_{11}$)	25		39	
11		26		40	
12		27		41	
13		28	$-\text{OH}$	42	
14	$-\text{CH}_2\text{CH}=\text{CHCH}_3$	29		43	$-\text{H}$
15		30			

Table 187




1	$-\text{CH}_3$	16		30	
2	$-\text{CH}_2\text{CH}_3$	17		31	
3	$-\text{CH}(\text{CH}_3)_2$	18	$-\text{C}\equiv\text{CH}$	32	
4	($-\text{C}_6\text{H}_5$)	19	$-\text{C}\equiv\text{CCH}_3$	33	
5		20		34	
6		21	$-\text{OCH}_3$	35	
7		22	$-\text{OC}_6\text{H}_5$	36	
8	$-\text{CH}_2-\text{N}(\text{CH}_3)_2$	23		37	
9	$-\text{CH}_2-\text{N}(\text{CH}_3)\text{CH}_2\text{C}_6\text{H}_5$	24		38	
10	($-\text{C}_6\text{H}_{11}$)	25		39	
11		26		40	
12		27		41	
13		28	$-\text{OH}$	42	
14	$-\text{CH}_2\text{CH}=\text{CHCH}_3$	29		43	$-\text{H}$
15		30			

Table 188




1	$-\text{CH}_3$	16		30	
2	$-\text{CH}_2\text{CH}_3$	17		31	
3	$-\text{CH}(\text{CH}_3)_2$	18	$-\text{C}\equiv\text{CH}$	32	
4	($-\text{C}_6\text{H}_5$)	19	$-\text{C}\equiv\text{CCH}_3$	33	
5		20		34	
6		21	$-\text{OCH}_3$	35	
7		22	$-\text{OC}_6\text{H}_5$	36	
8	$-\text{CH}_2-\text{N}(\text{CH}_3)_2$	23		37	
9	$-\text{CH}_2-\text{N}(\text{CH}_3)\text{CH}_2\text{C}_6\text{H}_5$	24		38	
10	($-\text{C}_6\text{H}_{11}$)	25	$-\text{N}-\text{CH}_3$	39	
11		26		40	
12		27		41	
13		28	$-\text{OH}$	42	
14	$-\text{CH}_2\text{CH}=\text{CHCH}_3$	29		43	
15		30			

Table 189




1	$-\text{CH}_3$	16		30	
2	$-\text{CH}_2\text{CH}_3$	17		31	
3	$-\text{CH}(\text{CH}_3)_2$	18	$-\text{C}\equiv\text{CH}$	32	
4		19	$-\text{C}\equiv\text{CCH}_3$	33	
5		20		34	
6		21	$-\text{OCH}_3$	35	
7		22	$-\text{OC}_6\text{H}_5$	36	
8	$-\text{CH}_2\text{N}(\text{CH}_3)_2$	23		37	
9	$-\text{CH}_2\text{N}(\text{CH}_3)\text{CH}_2\text{C}_6\text{H}_5$	24		38	
10		25		39	
11		26		40	
12		27		41	
13		28	$-\text{OH}$	42	
14	$-\text{CH}_2\text{CH}=\text{CHCH}_3$	29		43	
15		30			

Table 190


 R^1


1	$-\text{CH}_3$	16		30	
2	$-\text{CH}_2\text{CH}_3$	17		31	
3	$-\text{CH}(\text{CH}_3)_2$	18	$-\text{C}\equiv\text{CH}$	32	
4	($-\text{C}_6\text{H}_5$)	19	$-\text{C}\equiv\text{CCH}_3$	33	
5		20		34	
6		21	$-\text{OCH}_3$	35	
7		22	$-\text{OC}_6\text{H}_5$	36	
8	$-\text{CH}_2\text{-N}(\text{CH}_3)_2$	23		37	
9	$-\text{CH}_2\text{-N}(\text{CH}_3)\text{CH}_2\text{C}_6\text{H}_5$	24		38	
10	($-\text{C}_6\text{H}_{11}$)	25		39	
11		26		40	
12		27		41	
13		28	$-\text{OH}$	42	
14	$-\text{CH}_2\text{CH=CHCH}_3$	29		43	$-\text{H}$
15		30			

Table 191



1	$-\text{CH}_3$	16		30	
2	$-\text{CH}_2\text{CH}_3$	17		31	
3	$-\text{CH}(\text{CH}_3)_2$	18	$-\text{C}\equiv\text{CH}$	32	
4	($-\text{C}_6\text{H}_5$)	19	$-\text{C}\equiv\text{CCH}_3$	33	
5		20		34	
6		21	$-\text{OCH}_3$	35	
7		22	$-\text{OC}_6\text{H}_5$	36	
8	$-\text{CH}_2-\text{N}(\text{CH}_3)_2$	23		37	
9	$-\text{CH}_2-\text{N}(\text{CH}_3)\text{CH}_2\text{C}_6\text{H}_5$	24		38	
10	($-\text{C}_6\text{H}_{11}$)	25		39	
11		26		40	
12		27		41	
13		28	$-\text{OH}$	42	
14	$-\text{CH}_2\text{CH}=\text{CHCH}_3$	29		43	$-\text{H}$
15		30			

Table 192













1	$-\text{CH}_3$	16		30	
2	$-\text{CH}_2\text{CH}_3$	17		31	
3	$-\text{CH}(\text{CH}_3)_2$	18	$-\text{C}\equiv\text{CH}$	32	
4	 ($-\text{C}_6\text{H}_5$)	19	$-\text{C}\equiv\text{CCH}_3$	33	
5		20		34	
6		21	$-\text{OCH}_3$	35	
7		22	$-\text{OC}_6\text{H}_5$	36	
8	$-\text{CH}_2\text{N}(\text{CH}_3)_2$	23		37	
9	$-\text{CH}_2\text{N}(\text{CH}_3)\text{CH}_2\text{C}_6\text{H}_5$	24		38	
10	 ($-\text{C}_6\text{H}_{11}$)	25		39	
11		26		40	
12		27		41	
13		28	$-\text{OH}$	42	
14	$-\text{CH}_2\text{CH}=\text{CHCH}_3$	29		43	$-\text{H}$
15		30			

Table 193




1	$-\text{CH}_3$	16		30	
2	$-\text{CH}_2\text{CH}_3$	17		31	
3	$-\text{CH}(\text{CH}_3)_2$	18	$-\text{C}\equiv\text{CH}$	32	
4	($-\text{C}_6\text{H}_5$)	19	$-\text{C}\equiv\text{CCH}_3$	33	
5		20		34	
6		21	$-\text{OCH}_3$	35	
7		22	$-\text{OC}_6\text{H}_5$	36	
8	$-\text{CH}_2-\text{N}(\text{CH}_3)_2$	23		37	
9	$-\text{CH}_2-\text{N}(\text{CH}_3)\text{CH}_2\text{C}_6\text{H}_5$	24		38	
10	($-\text{C}_6\text{H}_{11}$)	25		39	
11		26		40	
12		27		41	
13		28	$-\text{OH}$	42	
14	$-\text{CH}_2\text{CH}=\text{CHCH}_3$	29		43	$-\text{H}$
15		30			

Table 194



1	$-\text{CH}_3$	16		30	
2	$-\text{CH}_2\text{CH}_3$	17		31	
3	$-\text{CH}(\text{CH}_3)_2$	18	$-\text{C}\equiv\text{CH}$	32	
4	($-\text{C}_6\text{H}_5$)	19	$-\text{C}\equiv\text{CCH}_3$	33	
5		20		34	
6		21	$-\text{OCH}_3$	35	
7		22	$-\text{OC}_6\text{H}_5$	36	
8	$-\text{CH}_2\text{N}(\text{CH}_3)_2$	23		37	
9	$-\text{CH}_2\text{N}(\text{CH}_3)\text{CH}_2\text{C}_6\text{H}_5$	24		38	
10	($-\text{C}_6\text{H}_{11}$)	25		39	
11		26		40	
12		27		41	
13		28	$-\text{OH}$	42	
14	$-\text{CH}_2\text{CH}=\text{CHCH}_3$	29		43	$-\text{H}$
15		30			


Table 195



R¹


1	-CH ₃	16		30	
2	-CH ₂ CH ₃	17		31	
3	-CH(CH ₃) ₂	18		32	
4	(-C ₆ H ₅)	19		33	
5		20		34	
6		21		35	
7		22		36	
8	-CH ₂ -N(CH ₃) ₂	23		37	
9	-CH ₂ -N(CH ₃)CH ₂ C ₆ H ₅	24		38	
10	(-C ₆ H ₁₁)	25		39	
11		26		40	
12		27		41	
13		28		42	
14	-CH ₂ CH=CHCH ₃	29		43	
15		30			

Table 196

R¹


1	-CH ₃	16		30	
2	-CH ₂ CH ₃	17		31	
3	-CH(CH ₃) ₂	18		32	
4	(-C ₆ H ₅)	19		33	
5		20		34	
6		21		35	
7		22		36	
8	-CH ₂ -N(CH ₃) ₂	23		37	
9	-CH ₂ -N(CH ₃)CH ₂ C ₆ H ₅	24		38	
10	(-C ₆ H ₁₁)	25		39	
11		26		40	
12		27		41	
13		28		42	
14	-CH ₂ CH=CHCH ₃	29		43	
15		30			

Table 197




1	$-\text{CH}_3$	16		30	
2	$-\text{CH}_2\text{CH}_3$	17		31	
3	$-\text{CH}(\text{CH}_3)_2$	18	$-\text{C}\equiv\text{CH}$	32	
4	($-\text{C}_6\text{H}_5$)	19	$-\text{C}\equiv\text{CCH}_3$	33	
5		20		34	
6		21	$-\text{OCH}_3$	35	
7		22	$-\text{OC}_6\text{H}_5$	36	
8	$-\text{CH}_2-\text{N}(\text{CH}_3)_2$	23		37	
9	$-\text{CH}_2-\text{N}(\text{CH}_3)\text{CH}_2\text{C}_6\text{H}_5$	24		38	
10	($-\text{C}_6\text{H}_{11}$)	25		39	
11		26		40	
12		27		41	
13		28	$-\text{OH}$	42	
14	$-\text{CH}_2\text{CH}=\text{CHCH}_3$	29		43	$-\text{H}$
15		30			

Table 198




1	$-\text{CH}_3$	16		30	
2	$-\text{CH}_2\text{CH}_3$	17		31	
3	$-\text{CH}(\text{CH}_3)_2$	18	$-\text{C}\equiv\text{CH}$	32	
4	($-\text{C}_6\text{H}_5$)	19	$-\text{C}\equiv\text{CCH}_3$	33	
5		20		34	
6		21	$-\text{OCH}_3$	35	
7		22	$-\text{OC}_6\text{H}_5$	36	
8	$-\text{CH}_2\text{N}(\text{CH}_3)_2$	23		37	
9	$-\text{CH}_2\text{N}(\text{CH}_3)\text{CH}_2\text{C}_6\text{H}_5$	24		38	
10		25		39	
11		26		40	
12		27		41	
13		28	$-\text{OH}$	42	
14	$-\text{CH}_2\text{CH}=\text{CHCH}_3$	29		43	$-\text{H}$
15		30			

Table 199

R¹

1	—CH ₃	16		30	
2	—CH ₂ CH ₃	17		31	
3	—CH(CH ₃) ₂	18		32	
4	(-C ₆ H ₅)	19		33	
5		20		34	
6		21		35	
7		22		36	
8	—CH ₂ -N(CH ₃) ₂	23		37	
9	—CH ₂ -N(CH ₃)CH ₂ C ₆ H ₅	24		38	
10		25		39	
11		26		40	
12		27		41	
13		28		42	
14	—CH ₂ CH=CHCH ₃	29		43	
15		30			

Table 200



1	$-\text{CH}_3$	16		30	
2	$-\text{CH}_2\text{CH}_3$	17		31	
3	$-\text{CH}(\text{CH}_3)_2$	18	$-\text{C}\equiv\text{CH}$	32	
4	($-\text{C}_6\text{H}_5$)	19	$-\text{C}\equiv\text{CCH}_3$	33	
5		20		34	
6		21	$-\text{OCH}_3$	35	
7		22	$-\text{OC}_6\text{H}_5$	36	
8	$-\text{CH}_2-\text{N}(\text{CH}_3)_2$	23		37	
9	$-\text{CH}_2-\text{N}(\text{CH}_3)\text{CH}_2\text{C}_6\text{H}_5$	24		38	
10	($-\text{C}_6\text{H}_{11}$)	25		39	
11		26		40	
12		27		41	
13		28	$-\text{OH}$	42	
14	$-\text{CH}_2\text{CH}=\text{CHCH}_3$	29		43	
15		30			

Table 201





Table 202







Table 203



Tabl 204







Table 205




1		11		20	
2		12		21	
3		13		22	
4		14		23	
5		15		24	
6		16		25	
7		17		26	
8		18		27	
9		19		28	
10				29	
				30	
				31	
				32	

Table 206




1		11		20	
2		12		21	
3		13		22	
4		14		23	
5		15		24	
6		16		25	
7		17		26	
8		18		27	
9		19		28	
10				29	
				30	
				31	
				32	

Table 207




1		11		20	
2		12		21	
3		13		22	
4		14		23	
5		15		24	
6		16		25	
7		17		26	
8		18		27	
9		19		28	
10				29	
				30	
				31	
				32	

Table 208




1		11		20	
2		12		21	
3		13		22	
4		14		23	
5		15		24	
6		16		25	
7		17		26	
8		18		27	
9		19		28	
10				29	
				30	
				31	
				32	

Table 209





1		11		20	
2		12		21	
3		13		22	
4		14		23	
5		15		24	
6		16		25	
7		17		26	
8		18		27	
9		19		28	
10				29	
				30	
				31	
				32	

Table 210



1		11		20	
2		12		21	
3		13		22	
4		14		23	
5		15		24	
6		16		25	
7		17		26	
8		18		27	
9		19		28	
10				29	
				30	
				31	
				32	

Table 211



Tabl 212





Table 213





Table 214








Table 215



1		11		20	
2		12		21	
3		13		22	
4		14		23	
5		15		24	
6		16		25	
7		17		26	
8		18		27	
9		19		28	
10				29	
				30	
				31	
				32	

Table 216



Tabl 217





Table 218





Table 219









Table 220



- 281 -


Table 221



Preparation Process

Also according to the present invention, there are provided processes for preparing the compounds of the present invention. Synthesis schemes, generic
5 Schemes 1-4 and specific Schemes A-D, illustrating such processes are illustrated below.

Scheme 1






P is a protecting group

P' is a protecting group


L is a leaving group

 $\text{R}^1, \text{R}^2, \text{R}^{3a}, \text{R}^{3b}$ are as defined herein


Schemes 2A and 2B




Scheme 3




Schemes 4A and 4B




Scheme 4A



Scheme 4B




Scheme A




- 288 -


Scheme B




Scheme C




Scheme D



Scheme E-1




Scheme E-2




- 292 -


Scheme 5




Scheme 6A



Scheme 6B



Scheme 7



As utilized herein, the term "alkyl", alone or in combination, means a straight-chain or branched-chain alkyl radical containing 1 to about 18 carbon atoms, preferably 1 to about 12 carbon atoms, and more preferably 1 to about 8 carbon atoms. Examples of such radicals include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, iso-amyl, hexyl, octyl and the like.

10 The term "alkenyl", alone or in combination, means a straight-chain or branched-chain hydrocarbon radical having one or more double bonds and containing 2 to about 18 carbon atoms preferably 2 to about 12 carbon atoms, and more preferably, 2 to about 8 carbon atoms. Examples of suitable alkenyl radicals include ethenyl (vinyl), 2-propenyl, 3-propenyl, 1,4-pentadienyl, 1,4-butadienyl, 1-butenyl, 2-butenyl, 3-butenyl, decenyl and the like.

 The term "alkynyl", alone or in
20 combination, means a straight-chain hydrocarbon radical having one or more triple bonds and containing 2 to about 18 carbon atoms, preferably 2 to about 12 carbon atoms, and more preferably, 2 to about 8 carbon atoms. Examples of alkynyl radicals
25 include ethynyl, 2-propynyl, 3-propynyl, decynyl, 1-butyne, 2-butyne, 3-butyne, and the like.

 The term "carbonyl" or "oxo", alone or in combination, means a $-C(=O)-$ group wherein the remaining two bonds (valences) can be independently substituted. The term carbonyl is also intended to encompass a hydrated carbonyl group $-C(OH)_2-$.
30

 The term "thiol" or "sulfhydryl", alone or in combination, means a $-SH$ group. The term "thio"

or "thia", alone or in combination, means a thiaether group; i.e., an ether group wherein the ether oxygen is replaced by a sulfur atom.

The term "amino", alone or in combination,
5 means an amine or -NH_2 group whereas the term mono-substituted amino, alone or in combination, means a substituted amine $\text{-N(H)(substituent)}$ group wherein one hydrogen atom is replaced with a substituent, and disubstituted amine means a -N(substituent)_2 wherein
10 two hydrogen atoms of the amino group are replaced with independently selected substituent groups.

Amines, amino groups and amides are compounds that can be designated as primary (I°), secondary (II°) or tertiary (III°) or unsubstituted,
15 mono-substituted or N,N-disubstituted depending on the degree of substitution of the amino nitrogen. Quaternary amine (ammonium) (IV°) means a nitrogen with four substituents $[\text{-N}^+(\text{substituent})_4]$ that is positively charged and accompanied by a counter ion,
20 whereas N-oxide means one substituent is oxygen and the group is represented as $[\text{-N}^+(\text{substituent})_3\text{-O}^-]$; i.e., the charges are internally compensated.

The term "cyano", alone or in combination, means a $\text{-C-triple bond-N (-C}\equiv\text{N, nitrile)}$ group. The
25 term "azido", alone or in combination, means a $\text{-N-triple bond-N (-N}\equiv\text{N)}$ group. The term "hydroxyl", alone or in combination, means a -OH group. The term "nitro", alone or in combination, means a -NO_2 group. The term "azo", alone or in combination, means a -
30 N=N- group wherein the bonds at the terminal positions can be independently substituted.

The term "hydrazino", alone or in combination, means a -NH-NH- group wherein the depicted remaining two bonds (valences) can be independently substituted. The hydrogen atoms of the hydrazino group can be replaced, independently, with substituents and the nitrogen atoms can form acid addition salts or be quaternized.

The term "sulfonyl", alone or in combination, means a -SO₂- group wherein the depicted remaining two bonds (valences) can be independently substituted. The term "sulfoxido", alone or in combination, means a -SO- group wherein the remaining two bonds (valences) can be independently substituted.

The term "sulfone", alone or in combination, means a -SO₂- group wherein the depicted remaining two bonds (valences) can be independently substituted. The term "sulfenamide", alone or in combination, means a -SON= group wherein the remaining three depicted bonds (valences) can be independently substituted. The term "sulfide", alone or in combination, means a -S- group wherein the remaining two bonds (valences) can be independently substituted.

The term "alkoxy", alone or in combination, means an alkyl ether radical or radicals with one, two or three oxygen atoms wherein the term alkyl is as defined above. Examples of suitable alkyl ether radicals include methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, iso-butoxy, sec-butoxy, tert-butoxy, methoxyethoxypropyl (CH₃OCH₂CH₂OCH₂CH₂CH₂-), 1,1-dimethoxyethane. 1,2-dimethoxyethane and the

like. The term "alkyloxy" is used to mean a substituted alkoxy group.

The term "cycloalkyl", alone or in combination, means a cyclic alkyl radical that contains 3 to about 8 carbon atoms. The term "cycloalkylalkyl" means an alkyl radical as defined above that is substituted by a cycloalkyl radical containing 3 to about 8, preferably 3 to about 6, carbon atoms. Examples of such cycloalkyl radicals include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and the like.

A heterocyclic (heterocyclo) group or the like alone or in combination is a saturated, unsaturated or partially unsaturated (non-aromatic) monocyclic, bicyclic or tricyclic heterocycle that contains one or more hetero atoms, typically one to three hetero atoms selected from nitrogen, oxygen and sulfur. A heterocyclic group can contain 4 to about 14 atoms in the one to three rings that also contain at least one nitrogen, oxygen or sulfur atom in addition to the carbon atoms. Preferably, a single ring is present and that ring contains 5 to 7 atoms and one hetero atom. Sulfur atoms, independently, may optionally be oxidized to, for example, -SO- or -SO₂-groups. Such a moiety can be optionally substituted on one or more ring carbon atoms by halogen, alkyl, alkoxy, oxo, and the like or as stated herein, and/or on a secondary nitrogen atom (i.e., -NH-) of the ring by alkyl, aralkoxycarbonyl, alkanoyl, aryl or arylalkyl or other groups listed herein or on a tertiary nitrogen atom (i.e., =N-) by oxido and that is attached via a carbon atom. The tertiary nitrogen atom with three substituents can

also attached to form a N-oxide [=N(O)-] group. A "heterocycloalkyl" group is an alkyl group substituted with a heterocyclo group.

The term "aryl", alone or in combination, means a 5- or 6-membered carbocyclic aromatic ring-containing moiety or a fused ring system containing two or three rings that have all carbon atoms in the ring; i.e., a carbocyclic aryl radical. Exemplary carbocyclic aryl radicals include phenyl, indenyl and naphthyl radicals.

The term "biaryl", alone or in combination means an aryl ring as define herein connected directly by a single bond to further aryl rings. Exemplary biaryl radicals include phenyl-phenyl (biphenyl), 2-phenylnaphtlenyl and phenylindenyl and 1-phenyl-anthracenyl radicals.

The term "heteroaryl" alone or in combination means a 5- or 6-membered aromatic ring-containing moiety or a fused ring system (radical) containing two or three rings that have carbon atoms and also one or more heteroatoms in the ring(s) such as sulfur, oxygen and nitrogen. Sulfur atoms, independently, may optionally be oxidized to, for example, -SO- or -SO₂-groups. Nitrogen atoms, independently, may be optionally oxidized to, for example, N-oxide groups or quaternized. Examples of such heterocyclic or heteroaryl groups are pyrrolidinyl, piperidyl, piperazinyl, morpholinyl, thiamorpholinyl, pyrrolyl, imidazolyl (e.g., imidazol-4-yl, 1-benzyloxycarbonylimidazol-4-yl, and the like), pyrazolyl, pyridyl, pyridyl-N-oxide, pyrazinyl, pyrimidinyl, furyl, tetrahydrofuryl, thienyl, thienyl-S-oxide, triazolyl, oxazolyl,

oxadiazoyl, thiazolyl, thiadiazoyl, indolyl (e.g.,
2-indolyl, and the like), quinolinyl, (e.g.,
2-quinolinyl, 3-quinolinyl, 1-oxido-2-quinolinyl, and
the like), isoquinolinyl (e.g., 1-isoquinolinyl,
5 3-isoquinolinyl, and the like), tetrahydroquinolinyl
(e.g., 1,2,3,4-tetrahydro-2-quinolyl, and the like),
1,2,3,4-tetrahydroisoquinolinyl (e.g., 1,2,3,4-
tetrahydro-1-oxo-isoquinolinyl, and the like),
quinoxalinyl, β -carbolinyl, 2-benzofurancarbonyl,
10 benzothiophenyl, 1-, 2-, 4- or 5-benzimidazolyl, and
the like radicals.

The term " heterocyclocarbonyl ", alone or
in combination, means a heterocyclogroup attached to
a $-(C=O)-$ group.

15 The term " heterocyclooxycarbony ", alone
or in combination, means a heterocyclogroup attached
to a $-O(C=O)-$ group.

The term " heterocycloalkoxycarbony ",
alone or in combination, means a heterocyclogroup
20 attached to a $-alkylO(C=O)-$ group.

The term " heterocycloalkyl", alone or in
combination, means a heterocyclogroup attached to an
alkyl group.

The term "aralkyl", alone or in
25 combination, means an alkyl radical as defined above
in which one hydrogen atom is replaced by an aryl
radical as defined above, such as benzyl, 2-
phenylethyl and the like.

The term "aralkoxycarbonyl", alone or in
30 combination, means a radical of the formula aralkyl-
 $O-C(O)-$ in which the term "aralkyl" has the
significance given above. An example of an
aralkoxycarbonyl radical is benzyloxycarbonyl.

The term "aryloxy" means a radical of the formula aryl-O- in which the term aryl has the significance given above. The phenoxy radical is an exemplary aryloxy radical.

5 The terms "heteroaralkyl" and "heteroaryloxy" mean radicals structurally similar to aralkyl and aryloxy that are formed from heteroaryl radicals. Exemplary radicals include 4-picolinyl and 2-pyrimidinoxy, respectively.

10 The terms "alkanoyl" or "alkylcarbonyl", alone or in combination, means an acyl radical derived from an alkanecarboxylic acid, examples of which include formyl, acetyl, propionyl, butyryl, valeryl, 4-methylvaleryl, and the like.

15 The term "cycloalkylcarbonyl" means an acyl group derived from a monocyclic or bridged cycloalkanecarboxylic acid such as cyclopropanecarbonyl, cyclohexanecarbonyl, adamantanecarbonyl, and the like, or from a benz-
20 fused monocyclic cycloalkanecarboxylic acid that is optionally substituted by, for example, alkanoylamino, such as 1,2,3,4-tetrahydro-2-naphthoyl, 2-acetamido-1,2,3,4-tetrahydro-2-naphthoyl.

25 The terms "aralkanoyl" or "aralkylcarbonyl" mean an acyl radical derived from an aryl-substituted alkanecarboxylic acid such as phenylacetyl, 3-phenylpropionyl (hydrocinnamoyl), 4-phenylbutyryl, (2-naphthyl)acetyl, 4-chlorohydrocinnamoyl,
30 4-aminohydrocinnamoyl, 4-methoxyhydrocinnamoyl and the like.

The terms "aroyl" or "arylcarbonyl" means an acyl radical derived from an aromatic carboxylic

acid. Examples of such radicals include aromatic carboxylic acids, an optionally substituted benzoic or naphthoic acid such as benzoyl, 4-chlorobenzoyl, 4-carboxybenzoyl, 4-(benzyloxycarbonyl)benzoyl, 1-naphthoyl, 2-naphthoyl, 6-carboxy-2 naphthoyl, 6-(benzyloxycarbonyl)-2-naphthoyl, 3-benzyloxy-2-naphthoyl, 3-hydroxy-2-naphthoyl, 3-(benzyloxyformamido)-2-naphthoyl, and the like.

The term "cycloalkylalkoxycarbonyl" means an acyl group of the formula cycloalkylalkyl-O-CO- wherein cycloalkylalkyl has the significance given above. The term "aryloxyalkanoyl" means an acyl radical of the formula aryl-O-alkanoyl wherein aryl and alkanoyl have the significance given above. The term "heterocyclooxycarbonyl" means an acyl group having the formula heterocyclo-O-CO- wherein heterocyclo is as defined above.

The term "heterocycloalkanoyl" is an acyl radical of the formula heterocyclo-substituted alkane carboxylic acid wherein heterocyclo has the significance given above. The term "heterocycloalkoxycarbonyl" means an acyl radical of the formula heterocyclo-substituted alkane-O-CO- wherein heterocyclo has the significance given above. The term "heteroaryloxy carbonyl" means an acyl radical represented by the formula heteroaryl-O-CO- wherein heteroaryl has the significance given above.

The term "aminocarbonyl" (carboxamide) alone or in combination, means an amino-substituted carbonyl (carbamoyl) group derived from an amine reacted with a carboxylic acid wherein the amino (amido nitrogen) group is unsubstituted (-NH₂) or a substituted primary or secondary amino group

containing one or two substituents selected from the group consisting of hydrogen, alkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl radicals and the like, as recited. A hydroxamate is a N-hydroxycarboxamide.

5 The term "aminoalkanoyl" means an acyl group derived from an amino-substituted alkanecarboxylic acid wherein the amino group can be a primary or secondary amino group containing substituents independently selected from hydrogen,
10 alkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl radicals and the like.

 The term "halogen" means fluoride, chloride, bromide or iodide. The term "haloalkyl" means an alkyl radical having the significance as
15 defined above wherein one or more hydrogens are replaced with a halogen. Examples of such haloalkyl radicals include chloromethyl, 1-bromoethyl, fluoromethyl, difluoromethyl, trifluoromethyl, 1,1,1-trifluoroethyl and the like.

20 The term "perfluoroalkyl" means an alkyl group wherein each hydrogen has been replaced by a fluorine atom. Examples of such perfluoroalkyl groups, in addition to trifluoromethyl above, are perfluorobutyl, perfluoroisopropyl, perfluorododecyl
25 and perfluorodecyl.

 The term "perfluoroalkoxy" alone or in combination, means a perfluoroalkyl ether radical wherein the term perfluoroalkyl is as defined above. Examples of such perfluoroalkoxy groups, in addition
30 to trifluoromethoxy ($\text{F}_3\text{C-O-}$), are perfluorobutoxy, perfluoroisopropoxy, perfluorododecoxy and perfluorodecoxy.

The term "perfluoroalkylthio" alone or in combination, means a perfluoroalkyl thioether radical wherein the term perfluoroalkyl is as defined above. Examples of such perfluoroalkylthio groups, in
5 addition to trifluoromethylthio ($\text{F}_3\text{C-S-}$), are perfluorobutylthio, perfluoroisopropylthio, perfluorododecylthio and perfluorodecylthio.

The term "aromatic ring" in combinations such as substituted-aromatic ring sulfone or
10 substituted-aromatic ring sulfoxide means aryl or heteroaryl as defined before.

Compounds contemplated herein can possess one or more asymmetric carbon atoms and are thus capable of existing in the form of optical isomers,
15 enantiomers, diastereoisomers, as well as in the form of racemic or nonracemic mixtures. A compound can also exist in other isomeric forms such as ortho, meta and para isomers, cis and trans isomers, syn and anti isomers, E and Z isomers, tautomeric isomers,
20 alpha and beta isomers, axial and equatorial isomers and isomers due to hindered rotation. An isomer can exist in equilibrium with another isomer in a mammal or a test system. Such isomeric equilibria can also occur during synthesis, storage, formulation, as
25 formulated pharmaceuticals, as liquids, solutions, solids, polymorphs and the like. Such a compound can also exist as an isomeric equilibrium system with a solvent or water, for example, as a hydrated ketone or aldehyde, hemiketal, hemiacetal, ketal, acetal or
30 other class or type of solvate as is well known in the art. All isomers are included as compounds of this invention.

The chemical reactions described herein are generally disclosed in terms of their broadest application to the preparation of the compounds of this invention. Occasionally, the reactions may not
5 be applicable as described to each compound included within the disclosed scope. The compounds for which this occurs will be readily recognized by those skilled in the art. In all such cases, either the reactions can be successfully performed by
10 conventional modifications known to those skilled in the art, e.g., by appropriate protection of interfering groups, by changing to alternative conventional reagents, by routine modification of reaction conditions, and the like, or other reactions
15 disclosed herein or otherwise conventional, are applicable to the preparation of the corresponding compounds that are contemplated.

"M" utilized in the reaction schemes that follow represents a leaving group such as halogen,
20 phosphate ester or sulfate ester.

It is understood that the definition of the compounds of the various formulas herein that contain asymmetric carbons, encompass all possible stereoisomers and mixtures thereof that possess the
25 activity discussed herein. In particular, it encompasses racemic modifications and any optical isomers which possesses the indicated activity. Optical isomers can be obtained in pure form by standard separation techniques.

30

Treatment Process

A process for treating a host mammal having a condition associated with pathological matrix

metalloprotease activity is also contemplated. That process comprises administering a compound described hereinbefore in an MMP enzyme-inhibiting effective amount to a mammalian host having such a condition.

5 The use of administration repeated a plurality of times is particularly contemplated.

A contemplated compound is used for treating a host mammal such as a mouse, rat, rabbit, dog, horse, primate such as a monkey, chimpanzee or
10 human that has a condition associated with pathological matrix metalloprotease activity.

Also contemplated is the similar use of a contemplated compound in the treatment of a disease state that can be affected by the activity of
15 metalloproteases such as TNF- α convertase or a member of the adamalysin family of enzymes such as ADAM 10. Exemplary of such disease states are the acute phase responses of shock and sepsis, coagulation responses, hemorrhage and cardiovascular effects, fever and
20 inflammation, anorexia and cachexia.

In treating a disease condition associated with pathological matrix metalloproteinase activity, a contemplated MMP inhibitor compound can be used, where appropriate, in the form of a pharmaceutically
25 acceptable amine salt derived from an inorganic or organic acid. Exemplary acid salts include but are not limited to the following: acetate, adipate, alginate, citrate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, camphorate,
30 camphorsulfonate, digluconate, cyclopentane-propionate, dodecylsulfate, ethanesulfonate, formate, glutamate, glucoheptanoate, gluconate, glucurantaes, glycerophosphate, hemisulfate, heptanoate, hexanoate,

fumarate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, isocitrate, lactate, malate, maleate, methanesulfonate, nicotinate, 2-naphthalenesulfonate, oxalate, oxalacetate, palmoate, 5 pectinate, persulfate, 3-phenylpropionate, phosphate, monohydrogen phosphate, dihydrogen phosphate, picrate, pivalate, propionate, pyruvate, succinate, tartrate, thiocyanate, tosylate, mesylate and undecanoate.

10 Also, a basic nitrogen-containing group can be quaternized with such agents as lower alkyl (C_1 - C_6) halides, such as methyl, ethyl, propyl, and butyl chloride, bromides, and iodides; dialkyl sulfates like dimethyl, diethyl, dibutyl, and diamyl sulfates, 15 long chain (C_8 - C_{20}) halides such as decyl, lauryl, myristyl and dodecyl chlorides, bromides and iodides, aralkyl halides like benzyl and phenethyl bromides, and others to provide enhanced water-solubility. Water or oil-soluble or dispersible products are 20 thereby obtained as desired. The salts are formed by combining the basic compounds with the desired acid.

Other compounds useful in this invention that are acids can also form pharmaceutically acceptable salts. Pharmaceutically acceptable cations include 25 metallic ions and organic ions. More preferred metallic ions include, but are not limited to appropriate alkali metal (Group Ia) salts, alkaline earth metal (Group IIa) salts and other physiological acceptable metal ions. Exemplary ions include 30 aluminum, calcium, lithium, magnesium, potassium, sodium and zinc in their usual valences. Preferred organic ions include protonated tertiary amines and

quaternary ammonium cations, including in part,
trimethylamine, diethylamine, N,N'-dibenzyl-
ethylenediamine, chlorprocaine, choline,
diethanolamine, ethylenediamine, meglumine (N-
5 methylglucamine) and procaine.

The term "pharmaceutically acceptable" is
used adjectivally herein to mean that the modified
noun is appropriate for use in a pharmaceutical
product.

10 In some cases, the salts can also be used
as an aid in the isolation, purification or
resolution of the compounds of this invention.

Total daily dose administered to a host
mammal in single or divided doses of an MMP enzyme-
15 inhibiting effective amount can be in amounts, for
example, of about 0.001 to about 100 mg/kg body
weight daily, preferably about 0.001 to about 30
mg/kg body weight daily and more usually about 0.01
to about 10 mg. Dosage unit compositions can contain
20 such amounts or submultiples thereof to make up the
daily dose.

A suitable dose can be administered, in
multiple sub-doses per day. Multiple doses per day
can also increase the total daily dose, should such
25 dosing be desired by the person prescribing the drug.
Such composition can be administered 1 to 6 times a
day, more usually 1 to 4 times a day.

The dosage regimen for treating a disease
condition with a compound and/or composition of this
30 invention is selected in accordance with a variety of
factors, including the type, age, weight, sex, diet
and medical condition of the patient, the severity of
the disease, the route of administration,

pharmacological considerations such as the activity, efficacy, pharmacokinetic and toxicology profiles of the particular compound employed, whether a drug delivery system is utilized and whether the compound
5 is administered as part of a drug combination. Thus, the dosage regimen actually employed can vary widely and therefore can deviate from the preferred dosage regimen set forth above.

A compound useful in the present invention
10 can be formulated as a pharmaceutical composition. Such a composition can then be administered orally, which is preferred, parenterally, by inhalation spray, rectally, or topically in dosage unit formulations containing conventional nontoxic
15 pharmaceutically acceptable carriers, adjuvants, and vehicles as desired. Topical administration can also involve the use of transdermal administration such as transdermal patches or iontophoresis devices. The term parenteral as used herein includes subcutaneous
20 injections, intravenous, intramuscular, intrasternal injection, or infusion techniques. Formulation of drugs is discussed in, for example, Hoover, John E., Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, Pennsylvania; 1975 and Liberman, H.A.
25 and Lachman, L., Eds., Pharmaceutical Dosage Forms, Marcel Decker, New York, N.Y., 1980.

Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions can be formulated according to the known art using
30 suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation can also be a sterile injectable solution or suspension in a nontoxic parenterally acceptable diluent or solvent,

for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that can be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile,
5 fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of
10 injectables. Dimethyl acetamide, surfactants including ionic and non-ionic detergents, polyethylene glycols can be used. Mixtures of solvents and wetting agents such as those discussed above are also useful.

15 Suppositories for rectal administration of the drug can be prepared by mixing the drug with a suitable nonirritating excipient such as cocoa butter, synthetic mono- di- or triglycerides, fatty acids and polyethylene glycols that are sold at
20 ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum and release the drug.

 Solid dosage forms for oral administration can include capsules, tablets, pills, powders, and
25 granules. In such solid dosage forms, the compounds of this invention are ordinarily combined with one or more adjuvants appropriate to the indicated route of administration. If administered *per os*, the compounds can be admixed with lactose, sucrose,
30 starch powder, cellulose esters of alcanoic acids, cellulose alkyl esters, talc, stearic acid, magnesium stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulfuric acids, gelatin, acacia

gum, sodium alginate, polyvinylpyrrolidone, and/or polyvinyl alcohol, and then tableted or encapsulated for convenient administration. Such capsules or tablets can contain a controlled-release formulation
5 as can be provided in a dispersion of active compound in hydroxypropylmethyl cellulose. In the case of capsules, tablets, and pills, the dosage forms can also comprise buffering agents such as sodium citrate, magnesium or calcium carbonate or
10 bicarbonate. Tablets and pills can additionally be prepared with enteric coatings.

For therapeutic purposes, formulations for parenteral administration can be in the form of aqueous or non-aqueous isotonic sterile injection
15 solutions or suspensions. These solutions and suspensions can be prepared from sterile powders or granules having one or more of the carriers or diluents mentioned for use in the formulations for oral administration. The compounds can be dissolved
20 in water, polyethylene glycol, propylene glycol, ethanol, corn oil, cottonseed oil, peanut oil, sesame oil, benzyl alcohol, sodium chloride, and/or various buffers. Other adjuvants and modes of administration are well and widely known in the pharmaceutical art.

25 Liquid dosage forms for oral administration can include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs containing inert diluents commonly used in the art, such as water. Such compositions can also comprise
30 adjuvants, such as wetting agents, emulsifying and suspending agents, and sweetening, flavoring, and perfuming agents.

The amount of active ingredient that can be combined with the carrier materials to produce a single dosage form varies depending upon the mammalian host treated and the particular mode of administration.

Best Mode for Carrying Out the Invention

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limiting of the remainder of the disclosure in any way whatsoever.

Abbreviations are often used for reagents and solvents in the specific examples that follow. Those abbreviations and their meanings are as follows:

BOC = t-butoxycarbonyl
DEAD = diethyl azodicarboxylate
DMF = dimethylformamide
DMPU = 1,3-dimethyl-3,4,5,6-tetrahydro-
2(1H)-pyrimidinone
EtOAc = ethyl acetate
EDC = 1-ethyl-3-[3-(dimethylamino)-
propyl]carbodiimide hydrochloride
Et₂O = diethyl ether
HOBT = 1-hydroxybenzotriazole
MeOH = methanol
MeCl₂ = methylene chloride
MsCl = methanesulfonyl chloride
NMM = N-methyl morpholine
THF = tetrahydrofuran


TsCl = toluenesulfonyl chloride

THP-O-hydroxylamine = O-tetrahydropyran-
hydroxylamine and O-tetrahydro-2H-
pyran-2-yl-hydroxylamine


5

Example 1: Preparation of

10



15



20


25

Part A: To a slurry of 4-hydroxypiperidine (46.3 g, 458 mmol) in tetrahydrofuran (400 mL) was added triethylamine (67 mL, 481 mmol), followed by slow addition of a solution of di-tert-butyl-dicarbonate (100 g, 458 mmol) in tetrahydrofuran (200 mL). The temperature was monitored and maintained below 32°C. The mixture was stirred for 4 hours before working up. Work up comprised stripping the tetrahydrofuran by rotary evaporation and taking the residue up in ethyl acetate (300 mL). The organic was then washed with 5% KHSO₄ (3x-150 mL), saturated NaHCO₃ (3x-150 mL), and brine (2x-150 mL). The organic portion was then dried over anhydrous MgSO₄,

-315-


filtered, and concentrated to afford a crude yellow oil. The oil was crystallized from hexanes providing the N-BOC-4-hydroxypiperidine product as a tan solid (86 g, 93% yield). ¹H NMR showed the desired

5 compound.




Part B: To a slurry of NaH (60% oil dispersion,
10 2.4 g, 60 mmol) in N,N-dimethylformamide (DMF; 70 mL)
cooled to zero degrees C under N₂ was slowly added a
solution of the N-BOC-4-hydroxypiperidine product
from Part A (10 g, 50 mmol) in DMF (20 mL). The
temperature was monitored and maintained at < 5°C.
15 The mixture was stirred 15 minutes before slowly
adding a solution of benzyl bromide (9 mL, 60 mmol)
in DMF (10 mL), keeping temperature at < 10°C. The
reaction was allowed to come to room temperature and
was stirred 12 hours. To quench, the reaction was
20 cooled to zero degrees C and H₂O (50 mL) was added.
Work up comprised stripping the solvents by rotary
evaporation and dissolving the residue in ethyl
acetate (150 mL) and H₂O (100 mL). The layers were
separated and the aqueous was extracted via ethyl
25 acetate (2x- 150 mL). The organic portions were
washed with saturated NaHCO₃ (2x-100 mL), H₂O (1x-150
mL), and brine (1x- 150 mL), then dried over Na₂SO₄,
filtered, and concentrated to afford a crude oil (18
g, 100% crude yield). ¹H NMR showed the desired

compound along with the benzyl bromide starting material.




5 **Part C:** To a solution of the crude product of Part B in 1,4-dioxane (10 mL) was added 4N HCl in dioxane (50 mL, 200 mmol). The mixture stirred at room temperature until starting material was gone by LC (~1 h). The solvents were then stripped and the residue was slurried in diethyl ether and filtered. The solid was washed with diethyl ether (2x-50 mL) and dried in vacuo to afford a white solid (11.5 g, 100% yield). ¹H NMR showed the desired compound as the HCl salt.




20 **Part D:** The HCl salt of Part C (10g, 44 mmol) and triethylamine (15.3 mL, 110 mmol) were slurried in CH₂Cl₂ (170 mL) and cooled to zero degrees C. A solution of methane sulfonyl chloride (5.1 mL, 66 mmol) in CH₂Cl₂ (50 mL) was slowly added, maintaining the temperature below 10°C with an ice bath. After the addition, the ice bath was removed and the reaction stirred for 1 hour as it came to room

temperature. After the disappearance of the starting material, the solvent was stripped and the residue was dissolved in ethyl acetate (100 mL) and H₂O (30 mL). Once separated, the organic layer was washed
5 with 5% KHSO₄ (3x-50 mL) and brine (1x- 50 mL). The organic layer was then dried over Na₂SO₄, filtered, and concentrated to afford an oily solid that was recrystallized from diethylether and hexanes, affording an off-white solid (12.3 g, 95% yield), SC
10 79767. ¹H NMR showed the desired compound. HPLC showed 100% at t_r= 12.1 minutes.




Part E: Oven-dried glassware was charged with
15 the product of Part D (5.0 g, 16.9 mmol) and tetrahydrofuran (34 mL) and the composition was cooled to -75°C. Lithium bis(trimethylsilyl)amide (1.0 M in tetrahydrofuran, 34 mL, 34 mmol) was slowly added, keeping temperature < -60°C. Reaction was
20 stirred for 30 minutes after the addition, and was then charged with a solution of methyl chloroformate (1.3 mmol, 16.9 mmol) in tetrahydrofuran (17 mL), again keeping the temperature < -60°C. After stirring for 1 hour at -75°C, the reaction was quenched with
25 saturated NH₄Cl, keeping temperature < -20°C. The aqueous portion froze into a solid chunk of ice. After warming to 5°C, the mixture was extracted with ethyl acetate (3x- 200 mL). The resulting organic

portions were washed with saturated NH_4Cl (2x-100 mL) and brine (1x-100mL), then dried over Na_2SO_4 and concentrated to afford the depicted product as a tan oil (5.0 g, 91% crude yield). ^1H NMR showed the
5 desired compound with some starting material present. HPLC showed 90% at $t_r = 13.9$ minutes, 10% at 12.1 minutes.




10 **Part F:** To a solution of the product of Part E (4.5 g, 13.7 mmol) and dibromodiethylether (1.9 mL, 15.1 mmol) in DMF (28 mL) was added 18-Crown-6 (500 mg, cat.) followed by potassium carbonate (3.8 g, 27.4 mmol). The mixture was heated at 60°C for 4
15 hours after which more potassium carbonate (1.9 g, 13.7 mmol) was added, and the reaction continued at 60°C for 14 hours. Liquid chromatography showed <10% starting material remained. The reaction was worked up by pouring into stirring 10% HCl_{aq} (200 mL). A
20 gummy solid resulted that was extracted with ethyl acetate (3x- 300 mL). The resulting organic portions were washed with brine (2x- 200 mL), dried over Na_2SO_4 , and concentrated to afford a dark brown oil. The product was crystallized from diethylether and
25 hexanes. The product was dried to afford the pyran methyl ester as an orange solid (3.7 g, 69% yield). ^1H NMR showed the desired compound. HPLC showed 96% at $t_r=25.2$ minutes.

-319-




Part G: To a solution of the product of Part F (3.5 g, 8.8 mmol) in tetrahydrofuran (20 mL) was added potassium trimethylsilonate (2.7 g, 21.1 mmol). The reaction stirred overnight (about eighteen hours) at room temperature. Liquid chromatography showed that < 3% starting material remained. Work up comprised stripping the tetrahydrofuran and taking the residue up in H₂O (100 mL). The solution was washed with diethylether (50 mL). The aqueous portion was then cooled to zero degrees C and 10% HCl_{aq} was slowly added until about pH 3. The acidic mixture was then extracted with ethyl acetate (3x-150 mL). The organic portions were washed with brine (1x- 100 mL), dried over Na₂SO₄, and concentrated to afford a wet solid. The solid was dried in vacuo with phosphorous pentoxide yielding an orange solid (2.4 g, 72% yield). ¹H NMR showed the desired carboxylic acid compound. HPLC showed 97% at t_r=21.3 minutes.



Part H: To a solution of the Part G acid product (2.4 g, 6.2 mmol) in dimethylacetamide (10 mL) was added N-methylmorpholine (2.0 mL, 18.6 mmol) followed by N-hydroxybenzotriazole hydrate (1.0 g,


7.4 mmol), O-(tetrahydro-2H-pyran-2-yl)hydroxylamine (1.1 g, 9.4 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.8 g, 9.4 mmol). The mixture stirred overnight (about eighteen hours) and was then stripped of solvent. The residue was dissolved in ethyl acetate (250 mL) and washed with 5% NaHSO₄ (1x- 150 mL), saturated potassium carbonate (1x-150 mL), and brine (1x-150 mL). The organic portions were then dried over Na₂SO₄ and concentrated to afford a viscous oil (3.2g, 100+% crude yield). ¹H NMR showed the desired compound. HPLC showed 95% at t_r=23.5 minutes.



Part I: The crude oil product of Part H (3.0 g, 6.2 mmol) was dissolved in acetonitrile (10 mL) and stirred with 10% HCl_{aq} (15 mL) for 2 hours at which time liquid chromatography showed no more starting material present. The acetonitrile was removed using an N₂ stream, affording a solid that was collected, washed with H₂O (1x-20 mL), and dried *in vacuo* to afford the product as a tan solid (1.6 g, 64% yield). ¹H NMR showed the desired compound. HPLC showed 99% at t_r=18.8 minutes. Mass spectroscopy showed M⁺_{found} = 399 (M⁺_{calc} = 399).

Example 2: Preparation of tetrahydro-N-hydroxy-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-2H-pyran-4-carboxamide

-321-



Part A: In dry equipment under nitrogen, 4-
5 hydroxypiperidine (20.2 g, 0.2 mol) was dissolved in
tetrahydrofuran (200 mL) and triethylamine (29 mL,
0.21 mol). A solution of di-*t*-butyldicarbonate
(43.65 g, 0.2 mol) was added at such a rate that the
temperature remained below thirty degrees Celsius.
10 After stirring at ambient temperature for four hours,
the reaction was concentrated *in vacuo*. The residue
was taken up in ethyl acetate, washed with water, 5%
KHSO₄, saturated NaHCO₃, brine, dried over Na₂SO₄,
filtered, and concentrated *in vacuo* to give the BOC
15 piperidine as a white solid (37.7 g, 94%).

Part B: Preparation of 1,1-Dimethylethyl
4-[4-(trifluoromethyl)phenoxy]-1-
piperidinecarboxylate

To a solution of the BOC piperidine from Part A (6.03
20 g, 30 mmol) in dimethylformamide (60 mL) was added
cesium carbonate (9.77 g, 30 mmol) and 4-
fluorobenzotrifluoride (3.8 mL, 30 mmol). The slurry
was stirred at ninety degrees Celsius. After
nineteen hours, cesium carbonate (3.26g, 10 mmol) and
25 4-fluorobenzotrifluoride (0.95ml mL, 10 mmol) were
added and the reaction continued at ninety degrees
Celsius. After a total of forty six hours, the
reaction was concentrated *in vacuo*. The residue was
taken up in ethyl acetate, washed with water three
30 times, brine, dried over Na₂SO₄, filtered, and

concentrated in vacuo. Chromatography (on silica, ethyl acetate/hexanes) provided the substituted BOC piperidine as a white solid (6.0 g, 58%).

Part C: Preparation of 4-[4-(trifluoromethyl)-
5 phenoxy]piperidine

To a slurry of the substituted BOC piperidine from Part B (5.95 g, 17.2 mmol) in 1,4-dioxane (10 mL) was added 4N HCl dioxane solution (17 mL). After one hour at ambient temperature the reaction was
10 concentrated in vacuo. The residue was slurried in diethyl ether and vacuum filtration of the resulting precipitate provided the hydrochloride salt as a white solid (4.6 g, 100%).

Part D: Preparation of 1-(Methylsulfonyl)-4-[4-
15 (trifluoromethyl)phenoxy]piperidine

To a solution of the hydrochloride salt from Part C (4.6 g, 16.9 mmol) and triethylamine (5.9 mL, 42.4 mmol) in methylene chloride (45 mL) at zero degrees Celsius was added a solution of
20 methanesulfonyl chloride (1.97 mL, 25.4 mmol) in methylene chloride (10 mL). After one hour at ambient temperature, the solvent was removed in vacuo. The residue was taken up in ethyl acetate, washed with water two times, brine, dried over Na₂SO₄,
25 filtered, and concentrated in vacuo. The residue was slurried in diethyl ether and vacuum filtration of the resulting precipitate provided the sulfonamide as an off-white solid (5.25 g, 96%).

Part E: Preparation of methyl [[4-[4-
30 (trifluoromethyl)phenoxy]-1-
piperidinyl]]sulfonyl]acetate

In dry equipment under nitrogen, the sulfonamide from Part D (4.2 g, 13 mmol) was dissolved in dry

tetrahydrofuran (26 mL), chilled to minus seventy-five degrees Celsius, and a 1M solution of lithium bis(trimethylsilyl)amide (26 mL) was added while maintaining the temperature below minus sixty-five
5 degrees. After thirty minutes at minus seventy-five degrees Celsius, a solution of methyl chloroformate (1.0 mL, 13 mmol) in dry tetrahydrofuran (13 mL) was added maintaining the temperature below minus sixty degrees. After thirty minutes at minus seventy-five
10 degrees Celsius, the reaction was quenched with saturated ammonium chloride solution (100 mL) and extracted with ethyl acetate. The combined extracts were washed with saturated ammonium chloride solution, brine, dried over Na₂SO₄, filtered, and
15 concentrated *in vacuo* to give the methylene sulfonamide as a yellow oil (4.95 g, 100%).

Part F: Preparation of Methyl tetrahydro-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-2H-pyran-4-
20 carboxylate

To a solution of the methylene sulfonamide from Part E (6.15 g, 16 mmol) in dimethylformamide (32 mL) was added potassium carbonate (7.8 g, 56.6 mmol), bis-(2-bromoethyl)ether (2.1 mL, 16 mmol) and 18-
25 Crown-6 (500 mg). The slurry was stirred at sixty degrees Celsius. After sixteen hours, potassium carbonate (2.0 g, 14 mmol) and bis-(2-bromoethyl)ether (0.2 mL, 1.6 mmol) were added and the reaction stirred at sixty degrees Celsius. After
30 a total of twenty two hours, the reaction was concentrated *in vacuo*. The residue was taken up in ethyl acetate, washed with water three times, brine, dried over Na₂SO₄, filtered, and concentrated *in*

-324-

WO 00/46221

vacuo. Chromatography (on silica, ethyl acetate/hexanes) provided the THP-substituted sulfonamide as a white solid (4.75 g, 65%).

Part G: Preparation of tetrahydro-4-[[4-[4-(trifluoromethyl)sulfonyl]-2H-pyran-4-piperidinyl]sulfonyl]-2H-pyran-4-carboxylic acid

5 In dry equipment under nitrogen, the THP-substituted sulfonamide from Part F (0.9 g, 2 mmol) was dissolved in dry tetrahydrofuran (4.0 mL) and potassium trimethylsilylate (0.38 g, 3.0 mmol) was added at ambient temperature. After twenty four hours water (100 mL) was added and the solution concentrated in vacuo. The residue was taken up in water and extracted with ethyl acetate to remove unreacted starting material. The aqueous solution was treated with 6 N HCl until pH=1. The slurry was extracted with ethyl acetate and the combined extracts washed with water, dried over Na₂SO₄, filtered, and concentrated in vacuo. The residue was heated in diethyl ether, the solid filtered and dried to give the carboxylic acid as a white solid (635 mg, 73%).

Part H: Preparation of tetrahydro-N-[[4-(trifluoromethyl)phenoxy]-1-[4-(trifluoromethyl)sulfonyl]-2H-pyran-4-carboxamide

25 In dry equipment under nitrogen, the carboxylic acid from Part G (3.0 g, 6.86 mmol) was dissolved in dry dimethylformamide (17 mL) and the remaining reagents were added to the solution in the following order: N-hydroxybenzotriazole hydrate (1.11

g, 8.24 mmol), N-methylmorpholine (2.26 mL, 20.6 mmol), O-(tetrahydro-2H-pyran-2-yl)hydroxylamine (2.49 g, 21.3 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.84 g, 9.6 mmol).


- 5 After two hours at ambient temperature, the reaction was concentrated *in vacuo*. The residue was taken up in ethyl acetate, washed with water, 5% KHSO₄, saturated NaHCO₃, brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Chromatography (on silica,
10 ethyl acetate/hexanes) provided the THP hydroxamate as a white foam (2.3 g, 63%). HRMS (ES+) M+ H⁺ calculated for C₂₃H₃₁N₂O₇ S₁F₃ : 537.1882, found 537.1856.

Part I: Preparation of tetrahydro-N-hydroxy-4-


- 15 [[4-[4-(trifluoromethyl)phenoxy]-1-
piperidinyllsulfonyl]-2H-pyran-4-carboxamide

- To a solution of the THP hydroxamate from Part H (1.55 g, 2.89 mmol) in 1,4-dioxane (7 mL) was added 4
20 N HCl dioxane solution (7 mL) and methanol (7 mL). After one hour at ambient temperature the reaction was diluted with ethyl acetate and washed with water, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The product was recrystallized
25 (acetone/hexanes) to give the depicted compound as a white solid (1.23 g, 95%). HRMS (ES+) M+ H⁺ calculated for C₁₈H₂₃N₂O₆ S₁F₃ : 453.1307, found 453.1319.

- 30 Example 3: Preparation of tetrahydro-N-hydroxy-4-
 [[4-[[4-[(trifluoromethyl)thio]phenyl]-
 thio]-1-piperidinyllsulfonyl]-2H-pyran-
 4-carboxamide




Part 1: Preparation of



5 To a slurry of Cs_2CO_3 (Aldrich, 20 g, 50 mmol) and 4-(methylsulfonyl)hydroxy-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester of Example 14, part B (5 g, 25 mmol) in acetone (70 mL) at 25°C under N_2 was slowly added 4-trifluoromethyl thiophenol (10 g, 50 mmol). The mixture was stirred 48 hours. After this time the acetone was removed by roto-evaporation and taking the residue up in ethyl acetate (150 mL) and H_2O (100 mL). The layers were separated and the aqueous was extracted via ethyl acetate (2x- 150 mL). The organics were washed with saturated K_2CO_3 (2x-100 mL), H_2O (1x-150 mL), and brine (1x- 150 mL), then dried over Na_2SO_4 , filtered, and concentrated to afford the crude N-Boc piperidine as an oil. The oil was purified on silica gel to give 6 g of a clear oil. ^1H NMR and mass spectrum were consistent with the desired compound.

Part 2: Preparation of




25 To a solution of the product (6 g) of Part 1 in 1,4-dioxane (10 mL) was added 4 N HCl in dioxane (50

-327-

mL, 200 mmol). The mixture stirred at room temperature until starting material was gone by LC (about one hour). The solvents were then removed and the residue was slurried in diethyl ether and
5 filtered. The solid was washed with diethyl ether (2x-50 mL) and dried in vacuo to afford the piperidine HCl salt as a white solid (6 g). ¹H NMR and mass spectrum showed the desired compound as the HCl salt.


10 Part 3: Preparation of



The HCl salt of Part 2 (6 g, 30 mmol) and triethylamine (Aldrich, 10 mL, 110 mmol) were slurried in CH₂Cl₂ (100 mL) and cooled to zero°C. A
15 solution of methane sulfonyl chloride (Aldrich, 4 g, 45 mmol) in CH₂Cl₂ (20 mL) was slowly added, maintaining the temperature below 10°C. After the addition, the ice bath was removed and the reaction stirred 1 hour as it warmed to ambient temperature.
20 After the disappearance of the starting material, the solvent was removed and the residue was taken up in ethyl acetate (100 mL) and H₂O (30 mL). Once separated, the organic layer was washed with 5% KHSO₄ (3x-50 mL) and brine (1x- 50 mL). The organic layer
25 was then dried over Na₂SO₄, filtered, and concentrated to afford the piperidine as an oily solid that was recrystallized from diethyl ether, affording an off-white solid (3.5 g). ¹H NMR and mass spectrum showed the desired compound.


30 Part 4: Preparation of

-328-



Oven-dried glassware was charged with the compound from Part 3 (3.0 g, 12 mmol), tetrahydrofuran (34 mL,) and cooled to -75°C. Lithium bis(trimethylsilyl)amide (Aldrich, 1.0 M in tetrahydrofuran, 35 mL, 33 mmol) was slowly added, keeping temperature less than -60°C. The reaction was stirred for 30 minutes after the addition and was then charged with a solution of methylchloroformate (Aldrich, 1.3 mmol, 16.9 mmol) in tetrahydrofuran (17 mL), again keeping the temperature at less than -60°C. After stirring for 1 hour at -75°C, the reaction was quenched with saturated NH₄Cl, keeping temperature at less than -20°C. The aqueous portion freezes into a solid chunk of ice. After warming to 5°C, the mixture was extracted via ethyl acetate (3x- 200 mL). Organics were washed with saturated NH₄Cl (2x-100 mL) and brine (1x-100mL), then dried over Na₂SO₄ and concentrated to afford the methylene piperidine as a tan oil (5.0 g, 91% crude yield). ¹H NMR and mass spectrum indicated desired compound.

Part 5: Preparation of




To a solution of compound from Part 4 (3 g, 11 mmol) and dibromo-diethylether (Lancaster, 1.8 mL, 15.1 mmol) in dimethylformamide (28 mL) was added 18-Crown-6 (Aldrich, 500 mg, cat.), followed by

-329-

potassium carbonate (Aldrich, 3.8 g, 27.4 mmol). The mixture was heated at 60°C for 16 hours. The product was isolated by pouring the reaction mixture into stirring 10% HCl_{aq} (200 mL) and extraction with ethylacetate (3x- 300 mL). Organics were washed with brine (2x- 200 mL), dried over Na₂SO₄, and concentrated to afford the ester as an oil. The oil was crystallized from diethylether (1.6 g). ¹H NMR and mass spectrum showed the desired compound.


10 Part 6: Preparation of



To a solution from Part 5 (2 g, 7 mmol) in tetrahydrofuran (20 mL) was added potassium trimethylsilonate (Aldrich, 2 g, 18 mmol). The reaction stirred overnight (about 18 hours) at room temperature. LC showed less than 3% starting material remained. Work up comprised removing the tetrahydrofuran and taking the residue up in H₂O (100 mL). The solution was washed with diethylether (50 mL). The aqueous was then cooled to zero°C and 10% HCl_{aq} was slowly added until pH = 3. The acidic mixture was then extracted with ethyl acetate (3x-150 mL). The organics were washed with brine (1x- 100 mL), dried over Na₂SO₄, and concentrated to afford a wet solid. The solid was dried in vacuo with phosphorous pentoxide yielding the acid as an orange solid (2.4 g, 72% yield). ¹H NMR and mass spectrum showed the desired compound.

25 Part 7: Preparation of

-330-





To a solution of the acid product in Part 6 (2.4 g, 6.2 mmol) in dimethylacetamide (10 mL) was added N-methylmorpholine (Aldrich, 2.0 mL, 18.6 mmol) followed by N-hydroxybenzotriazole hydrate (Aldrich, 1.0 g, 7.4 mmol), O- (tetrahydro-2H-pyran-2-yl) hydroxylamine (1.1 g, 9.4 mmol), and, lastly, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (Sigma, 1.8 g, 9.4 mmol). The mixture stirred overnight (about 18 hours) and was then stripped of solvent. The residue was taken up in ethyl acetate (250 mL) and washed with 5% NaHSO₄ (1x-150 mL), saturated potassium carbonate (1x-150 mL), and brine (1x-150 mL). The organic layer was dried over Na₂SO₄ and concentrated to afford a viscous oil. ¹H NMR and MS showed the desired compound.

The viscous crude oil (3.0 g, 6.2 mmol) was dissolved in acetonitrile (10 mL) and stirred with 10% HCl_{aq} (15 mL) for 2 hours, after which, liquid chromatography (LC) showed no more starting material. The acetonitrile was removed with N₂ stream over the surface of the solution affording a solid that was collected, washed with H₂O (1x-20 mL), and dried in vacuo to afford the product as a tan solid (1.6 g, 64% yield). ¹H NMR showed the desired compound. Mass spectroscopy showed: C₁₈H₂₃F₃N₂O₅S₃ M⁺_{found} = 500 (M⁺_{calc} = 500).


Example 4: Preparation of

-331-




5 **Part A:** To a slurry of 4-hydroxypiperidine (46.3 g, 458 mmol) in tetrahydrofuran (400 mL) was added triethylamine (67 mL, 481 mmol), followed by slow addition of a solution of di-tert-butyl-
 10 dicarbonate (100 g, 458 mmol) in tetrahydrofuran (200 mL). The temperature was monitored and maintained below 32°C. The mixture was stirred for 4 hours before working up. Work up consisted of removing the tetrahydrofuran *in vacuo* and taking the residue up in ethyl acetate (300 mL). The organic phase was then
 15 washed with 5% KHSO₄ (3x-150 mL), saturated NaHCO₃ (3x- 150 mL), and brine (2x-150 mL). The organic phase was then dried over anhydrous MgSO₄, filtered, and concentrated to afford a crude yellow oil. The oil was crystallized from hexanes providing the N-
 20 BOC-4-hydroxypiperidine product as a tan solid (86 g, 93% yield). ¹H NMR showed the desired compound.

-332-




Part B: To a slurry of NaH (60% oil dispersion,
5 2.4 g, 60 mmol) in N,N-dimethylformamide (70 mL)
cooled to zero degrees C under N₂ was slowly added a
solution of the N-BOC-4-hydroxypiperidine product
from Part A (10 g, 50 mmol) in N,N-dimethylformamide
(20 mL). The temperature was monitored and
10 maintained at < 5°C. The mixture was stirred 15
minutes before slowly adding a solution of benzyl
bromide (9 mL, 60 mmol) in N,N-dimethylformamide (10
mL), keeping temperature < 10°C. The reaction was
permitted to come to room temperature and was stirred
15 12 hours. To quench, the reaction was cooled to zero
degrees C and H₂O (50 mL) was added. Work up
consisted of removing the solvents in vacuo and
taking the residue up in ethyl acetate (150 mL) and
H₂O (100 mL). The layers were separated and the
20 aqueous layer was extracted with ethyl acetate (2x-
150 mL). The organic phase was washed with saturated
NaHCO₃ (2x-100 mL), H₂O (1x-150 mL), and brine (1x-
150 mL), then dried over Na₂SO₄, filtered, and
concentrated to afford a crude oil (18 g, 100% crude
25 yield). ¹H NMR showed the desired compound along with
the benzyl bromide starting material.

-333-



Part C: To a solution of the crude product of Part B in 1,4-dioxane (10 mL) was added 4 N HCl in dioxane (50 mL, 200 mmol). The mixture stirred at room temperature until starting material was gone by liquid chromatography (LC; about 1 hour). The solvents were then removed and the residue was slurried in diethyl ether and filtered. The solid was washed with diethyl ether (2x-50 mL) and dried in vacuo to afford a white solid (11.5 g, 100% yield). ¹H NMR showed the desired compound as the HCl salt.




15

Part D: The HCl salt of Part C (10g, 44 mmol) and triethylamine (15.3 mL, 110 mmol) were slurried in CH₂Cl₂ (170 mL) and cooled to zero degrees C with an ice bath. A solution of methanesulfonyl chloride (5.1 mL, 66 mmol) in CH₂Cl₂ (50 mL) was slowly added, maintaining the temperature below 10°C. After the addition, the ice bath was removed and the reaction stirred 1 h as it came to room temperature. After the disappearance of the starting material, the

25

solvent was removed and the residue was taken up in ethyl acetate (100 mL) and H₂O (30 mL). Once separated, the organic layer was washed with 5% KHSO₄ (3x-50 mL) and brine (1x- 50 mL). The organic layer was then dried over Na₂SO₄, filtered, and concentrated to afford an oily solid that was recrystallized from diethyl ether and hexanes, affording an off-white solid (12.3 g, 95% yield), the sulfonamide. ¹H NMR showed the desired compound. HPLC showed 100% at t_r= 12.1 minutes.




Part E: Oven-dried glassware was charged with the sulfonamide of Part D (5.0 g, 16.9 mmol) and tetrahydrofuran (34 mL) and cooled to -75°C. Lithium bis(trimethylsilyl)amide (1.0 M in tetrahydrofuran, 34 mL, 34 mmol) was slowly added, keeping temperature < -60°C. Reaction mixture was stirred for 30 minutes after the addition and was then charged with a solution of methyl chloroformate (1.3 mmol, 16.9 mmol) in tetrahydrofuran (17 mL) again keeping the temperature < -60°C. After stirring for 1 hour at -75°C, the reaction was quenched with saturated NH₄Cl, keeping temperature < -20°C. The aqueous portion froze into a solid chunk of ice. After warming to 5°C, the mixture was extracted via ethyl acetate (3x-200 mL). Organic layers were washed with saturated NH₄Cl (2x-100 mL) and brine (1x-100 mL), then dried over Na₂SO₄ and concentrated to afford the methyl

-335-


formate as a tan oil (5.0 g, 91% crude yield). ^1H NMR showed the desired compound with some starting material present. HPLC showed 90% at $t_r = 13.9$ minutes, 10% at 12.1 minutes.

5




Part F: To a solution of the methyl formate of Part E (4.5 g, 13.7 mmol) and dibromo-diethylether (Lancaster, 1.9 mL, 15.1 mmol) in dimethylformamide (28 mL) was added 18-Crown-6 (500 mg, cat.), followed by potassium carbonate (3.8 g, 27.4 mmol). The mixture was heated at 60°C for 4 hours, after which more potassium carbonate (1.9g, 13.7 mmol) was added, and the reaction continued at 60°C for 14 hours. LC showed <10% starting material remained. The reaction was worked up by pouring into stirring 10% HCl_{aq} (200 mL). A gummy solid resulted that was extracted with ethyl acetate (3x- 300 mL). Organic layers were washed with brine (2x- 200 mL), dried over Na_2SO_4 , and concentrated to afford a dark brown oil. Oil was crystallized from diethyl ether and hexanes. The solid was dried to afford the pyran methyl ester as an orange solid (3.7 g, 69% yield). ^1H NMR showed the desired compound. HPLC showed 96% at $t_r = 25.2$ minutes.

- 336 -




Part G: To a solution of the pyran methyl ester of Part F (3.5 g, 8.8 mmol) in tetrahydrofuran (20 mL) was added potassium trimethylsilonate (2.7 g, 21.1 mmol). The reaction stirred overnight (about eighteen hours) at room temperature. LC showed < 3% starting material remained. Work up consisted of removing the tetrahydrofuran and taking the residue up in H₂O (100 mL). The solution was washed with diethyl ether (50 mL). The aqueous phase was then cooled to zero degrees C and 10% HCl_{aq} was slowly added until pH value of about 3. The acidic mixture was then extracted with ethyl acetate (3x-150 mL). The organic layers were washed with brine (1x- 100 mL), dried over Na₂SO₄, and concentrated to afford a wet solid. The solid was dried in vacuo with phosphorous pentoxide yielding the carboxylic acid as an orange solid (2.4 g, 72% yield). ¹H NMR showed the desired compound. HPLC showed 97% at t_r=21.3 minutes.



Part H: To a solution of the Part G carboxylic acid product (2.4 g, 6.2 mmol) in dimethylacetamide

-337-


(10 mL) was added N-methylmorpholine (2.0 mL, 18.6 mmol), followed by N-hydroxybenzotriazole hydrate (1.0 g, 7.4 mmol), O-(tetrahydro-2H-pyran-2-yl) hydroxylamine (1.1 g, 9.4 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.8 g, 9.4 mmol). The mixture was stirred overnight (about eighteen hours) and then solvent was removed. The residue was taken up in ethyl acetate (250 mL) and washed with 5% NaHSO₄ (1x-150 mL), saturated potassium carbonate (1x-150 mL), and brine (1x-150 mL). The organic phase was then dried over Na₂SO₄ and concentrated to afford a viscous oil (3.2g, 100+% crude yield). ¹H NMR showed the desired compound. HPLC showed 95% at t_r=23.5 minutes.




Part I: The viscous oil product of Part H (3.0 g, 6.2 mmol) was dissolved in acetonitrile (10 mL) and stirred with 10% HCl_{aq} (15 mL) for 2 hours. After which time, LC showed no more starting material. The acetonitrile was removed via N₂ stream affording a solid that was collected, washed with H₂O (1x-20 mL), and dried in vacuo to afford the depicted hydroxamate as a tan solid (1.6 g, 64% yield). ¹H NMR showed the desired compound. HPLC showed 99% at t_r=18.8 minutes. Mass spectroscopy showed M⁺_{found} = 399 (M⁺_{calc} = 399).

Example 5: Preparation of 4-[[4-(4-bromophenyl)-4-hydroxy-1-piperidinyl]sulfonyl]-tetrahydro-N-hydroxy-2H-pyran-4-carboxamide

5



Part 1: Preparation of




4-(4-Bromophenyl)-4-hydroxypiperidine (Aldrich, 3 g, 1.3 mmol) and N-methylmorpholine (Aldrich, 1.5 g, 2.6 mmol) were slurried in CH_2Cl_2 (50 mL) and cooled to zero°C. A solution of methane sulfonyl chloride (Aldrich, 2 g, 2.1 mmol) in CH_2Cl_2 (20 mL) was slowly added, maintaining the temperature below 10°C. After the addition, the ice bath was removed and the reaction stirred 1 hour as it warmed to ambient temperature. After the disappearance of the starting material, the solvent was removed under reduced pressure. Water (100 mL) was added and 10% aqueous hydrochloride acid and the product filtered to result in the methylsulfonyl amide as an off-white solid (3.5 g). ^1H NMR and mass spectroscopy showed the desired compound.

25


-339-

Part 2: Preparation of




Oven-dried glassware was charged with the compound from Part 1 (5.0 g, 15 mmol) and tetrahydrofuran (30 mL,) and cooled to -75°C. Lithium bis(trimethylsilyl)amide (Aldrich, 1.0 M in tetrahydrofuran, 50 mL, 33 mmol) was slowly added, keeping temperature less than -60°C. Reaction stirred for 30 minutes after the addition and was then charged with a solution of methyl chloroformate (Aldrich, 1.3 mmol, 16.9 mmol) in tetrahydrofuran (17 mL) again keeping the temperature at less -60°C. After stirring for 1 hour at -75°C, the reaction was quenched with saturated NH₄Cl, keeping temperature less than -20°C. The aqueous does freeze into a solid chunk of ice. After warming to 5°C, the mixture was extracted via ethyl acetate (3x- 200 mL). Organics were washed with saturated NH₄Cl (2x-100 mL) and brine (1x-100mL), then dried over Na₂SO₄ and concentrated to afford the methylene sulfonamide as an amber oil (6.0 g, 91% crude yield). ¹H NMR and mass spectroscopy indicated desired compound.

Part 3: Preparation of



To a solution of compound from Part 2 (5 g, 13 mmol) and dibromo-diethylether (Lancaster, 1.8 mL, 15.1 mmol) in dimethylformamide (28 mL) was added 18-Crown-6 (Aldrich, 500 mg, cat.) followed by potassium carbonate (Aldrich, 3.8 g, 27.4 mmol). The mixture was heated at 60°C for 16 hours. The product was isolated by pouring into stirring 10% HCl_{aq} (200 mL) and extracted with ethyl acetate (3x- 300 mL). Organics were washed with brine (2x- 200 mL), dried over Na₂SO₄, and concentrated to afford the ester as an oil. The crystallized to result in 3 grams of a tan solid. ¹H NMR and mass spectroscopy showed the desired compound.

Part 4: Preparation of




15

To a solution from Part 3 (3 g, 7 mmol) in tetrahydrofuran (20 mL) was added potassium trimethylsilanolate (Aldrich, 3 g, 23 mmol). The reaction stirred overnight (about 18 hours) at room temperature. Liquid chromatography (LC) showed less than 3% starting material remained. Work up comprised removing the tetrahydrofuran and taking the residue up in H₂O (100 mL). The solution was washed with diethylether (50 mL). The aqueous was then cooled to 0°C and 10% HCl_{aq} was slowly added until pH = 3. The acidic mixture was then extracted via ethyl acetate (3x-150 mL). The organics were washed with brine (1x-100 mL), dried over Na₂SO₄, and concentrated to afford a wet solid. The solid was dried in vacuo

25

with phosphorous pentoxide yielding the acid as an orange solid (3 g, 72% yield). ^1H NMR and mass spectroscopy showed the desired compound.

Part 5: Preparation of



5

To a solution of the acid product in Part 4 (3.5 g, 8 mmol) in dimethylacetamide (10 mL) was added N-methylmorpholine (Aldrich, 2.0 mL, 18.6 mmol) followed by N-hydroxybenzotriazole hydrate (Aldrich, 1.0 g, 12 mmol), O-(tetrahydro-2H-pyran-2-yl) hydroxylamine (1.1 g, 12 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (Sigma, 2.3 g, 12 mmol). The mixture was stirred overnight (about 18 hours) and was then stripped of solvent. The residue was taken up in ethyl acetate (250 mL) and washed with 5% NaHSO₄ (1x-150 mL), saturated potassium carbonate (1x-150 mL), and brine (1x-150 mL). The organic layer was then dried over Na₂SO₄ and concentrated to afford a viscous oil. ^1H NMR and mass spectroscopy showed the desired compound.

The viscous crude oil (4.0 g, 6.2 mmol) was dissolved in acetonitrile (10 mL) and stirred with 10% HCl_{aq} (15 mL) for 2 hours, after which, LC showed no more starting material. The acetonitrile was removed with N₂ stream over the surface of the solution affording a solid that was collected, washed with H₂O (1x-20 mL), and dried in vacuo to afford the product as a tan solid (2 g). ^1H NMR showed the


25

-342-

desired compound. Mass spectroscopy showed:

$C_{17}H_{23}F_3N_2O_6SBr$ $M^{+H}_{found} = 463$ ($M^{+H}_{calc} = 463$).

Example 6: Preparation of ethyl 4-[(hydroxyamino)-
carbonyl]-4-[[4-[4-(trifluoromethyl)-
phenoxy]-1-piperidinyl]sulfonyl]-1-
piperidinecarboxylate



Part A: Preparation of 1,1-Dimethylethyl 4-hydroxy-
1-piperidinecarboxylate

In dry equipment under nitrogen, 4-hydroxypiperidine (20.2 g, 0.2 mol) was dissolved in tetrahydrofuran (200 mL) and triethylamine (29 mL, 0.21 mol). A solution of di-t-butyldicarbonate (43.65 g, 0.2 mol) was added at such a rate that the temperature remained below thirty degrees Celsius. After stirring at ambient temperature for four hours, the reaction was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water, 5% KHSO₄, saturated NaHCO₃, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo to give the BOC piperidine as a white solid (37.7 g, 94%).

Part B: Preparation of 1,1-Dimethylethyl 4-[4-(trifluoromethyl)phenoxy]-1-
piperidinecarboxylate

To a solution of the BOC piperidine from part A (6.03 g, 30 mmol) in dimethylformamide (60 mL) were

-343-

added cesium carbonate (9.77 g, 30 mmol) and 4-fluorobenzotrifluoride (3.8 mL, 30 mmol). The slurry was stirred at ninety degrees Celsius. After nineteen hours cesium carbonate (3.26g, 10 mmol) and 4-fluorobenzotrifluoride (0.95ml mL, 10 mmol) were added and the reaction continued at ninety degrees Celsius. After a total of forty six hours, the reaction was concentrated *in vacuo*. The residue was taken up in ethyl acetate, washed with water three times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Chromatography (on silica, ethyl acetate/hexanes) provided the substituted BOC piperidine as a white solid (6.0 g, 58%).

Part C: Preparation of 4-[4-(trifluoromethyl)-phenoxy]piperidine

To a slurry of the substituted BOC piperidine from part B (5.95 g, 17.2 mmol) in 1,4-dioxane (10 mL) was added 4N HCl dioxane solution (17 mL). After one hour at ambient temperature, the reaction was concentrated *in vacuo*. The residue was slurried in diethyl ether and vacuum filtration of the resulting precipitate provided the hydrochloride salt as a white solid (4.6 g, 100%).

Part D: Preparation of 1-(methylsulfonyl)-4-[4-(trifluoromethyl)phenoxy]piperidine

To a solution of the hydrochloride salt from part C (4.6 g, 16.9 mmol) and triethylamine (5.9 mL, 42.4 mmol) in dichloromethane (45 mL) at zero degrees Celsius was added a solution of methane sulfonyl chloride (1.97 mL, 25.4 mmol) in dichloromethane (10 mL). After one hour at ambient temperature, the solvent was stripped *in vacuo*. The residue was taken

-344-

up in ethyl acetate, washed with water two times,
saturated sodium chloride solution, dried over Na₂SO₄,
filtered, and concentrated in vacuo. The residue was
slurried in diethyl ether and vacuum filtration of
5 the resulting precipitate provided the sulfonamide as
an off-white solid (5.25 g, 96%).

Part E: Preparation of methyl [[4-[4-
(trifluoromethyl)phenoxy]-1-piperidinyl]-
sulfonyl]acetate

10 In dry equipment under nitrogen, the sulfonamide
from part D (4.2 g, 13 mmol) was dissolved in dry
tetrahydrofuran (26 mL), chilled to minus seventy-
five degrees Celsius, and a 1 M solution of lithium
bis(trimethylsilyl)amide (26 mL) was added while
15 maintaining the temperature below minus sixty five
degrees.

After thirty minutes at minus seventy-five
degrees Celsius, a solution of methyl chloroformate
(1.0 mL, 13 mmol) in dry tetrahydrofuran (13 mL) was
20 added maintaining the temperature below minus sixty
degrees. After thirty minutes at minus seventy-five
degrees Celsius, the reaction was quenched with
saturated ammonium chloride solution (100 mL) and
extracted with ethyl acetate. The combined extracts
25 were washed with saturated ammonium chloride
solution, saturated sodium chloride solution, dried
over Na₂SO₄, filtered, and concentrated in vacuo to
give the methylene sulfonamide as a yellow oil
(4.95 g, 100%).

30 Part F: Preparation of 1-Ethyl 4-methyl 4-[[4-[4-
(trifluoromethyl)phenoxy]-1-piperidinyl]-
sulfonyl]-1,4-piperidinecarboxylate

-345-

WO 00/46221

To a solution of the methylene sulfonamide from part E (6.15 g, 16 mmol) in dimethylformamide (32 mL) were added potassium carbonate (7.8 g, 56.6 mmol), bis-(2-bromoethyl)amine ethyl carbamate [3.0 g, 10.75 mmol; prepared by method found in Synth. Commun.; 11;1;1981;p.17-24] and 18-Crown-6 (500 mg). The slurry was stirred at sixty degrees Celsius. After sixteen hours, potassium carbonate (2.0 g, 14 mmol) was added and the reaction stirred at sixty degrees Celsius. After a total of twenty four hours, the reaction was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water three times, saturated sodium chloride solution, dried over Na_2SO_4 , filtered, and concentrated in vacuo. Chromatography (on silica, ethyl acetate/hexanes) provided the piperidine sulfonamide as a clear colorless oil (1.2 g, 31%).

Part G: Preparation of 1-Ethyl hydrogen 4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-sulfonyl]-1,4-piperidinecarboxylate

In dry equipment under nitrogen, the piperidine sulfonamide from part F (1.0 g, 1.9 mmol) was dissolved in dry tetrahydrofuran (7.0 mL) and potassium trimethylsilylate (0.38 g, 3.0 mmol) was added at ambient temperature. After eighteen hours, water (100 mL) was added and the solution concentrated in vacuo. The residue was taken up in water and extracted with ethyl acetate. The combined ethyl acetate extracts were concentrated in vacuo. The residue was wetted with water and 1 N HCl solution (1.5 mL) was added, the slurry was extracted with ethyl acetate and the combined extracts washed with saturated sodium chloride solution, dried over

Na₂SO₄, filtered, and concentrated in vacuo to give the carboxylic acid as a white foam (860 mg, 89%).

Part H: Preparation of Ethyl 4-[[[(tetrahydro-2H-pyran-2-yl)oxy]amino]carbonyl]-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-sulfonyl]-1-piperidinecarboxylate

In dry equipment under nitrogen, the carboxylic acid from part G (0.82 g, 1.6 mmol) was dissolved in dry dimethylformamide (4 mL) and the remaining reagents were added to the solution in the following order: N-hydroxybenzotriazole hydrate (0.26 g, 1.9 mmol), N-methylmorpholine (0.53 mL, 4.84 mmol), O-(tetrahydro-2H-pyran-2-yl)hydroxylamine (0.585 g, 5.0 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.43 g, 2.26 mmol). After two hours at ambient temperature, the reaction was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water, 5% KHSO₄, saturated NaHCO₃, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo. Chromatography (on silica, ethyl acetate/hexanes) provided the THP hydroxamate as a white foam (0.72 g, 73%).

Part I: Preparation of ethyl 4-[(hydroxyamino)-carbonyl]-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-1-piperidinecarboxylate


To a solution of the THP hydroxamate from part H (0.69 g, 1.1 mmol) in 1,4-dioxane (3 mL) were added 4 N HCl dioxane solution (3 mL) and methanol (3 mL). After two hours at ambient temperature, the reaction was diluted with ethyl acetate and washed with water, dried over Na₂SO₄, filtered, and concentrated in

-347-

vacuo. The product was recrystallized (acetone/hexanes) to give the title compound as a white solid (420 mg, 73%). HRMS (ES+) $M + NH_4^+$ calculated for $C_{21}H_{28}N_3O_7S_1F_3$: 541.1944, found 5 541.1904.

Example 7: Preparation of 4-[(3,5-dimethyl-1-piperidinyl)sulfonyl]tetrahydro-N-hydroxy-2H-pyran-4-carboxamide


10



Part A: 3,5-Dimethylpiperidine (70% cis/30% trans, 7.0 mL, 53 mmol) and triethylamine (11.2 mL, 80 mmol) were slurried in CH₂Cl₂ (75 mL) and cooled to zero degrees C. A solution of methanesulfonyl chloride (6.2 mL, 80 mmol) in CH₂Cl₂ (25 mL) was slowly added, maintaining the temperature < 10°C with an ice bath. After the addition, the ice bath was removed and the reaction stirred 1 hour as it came to room temperature. After the disappearance of the starting material, the solvent was removed and the residue was taken up in ethyl acetate (200 mL) and H₂O (50 mL). Once separated, the organic layer was washed with 5% KHSO₄ (3x-50 mL) and brine (1x- 50 mL). The organic layer was then dried over Na₂SO₄, filtered, and concentrated to afford the methyl sulfonamide an off white solid (10.0 g, 99% yield. ¹H NMR showed the desired compound.


30

-348-



Part B: Oven-dried glassware was charged with the methyl sulfonamide product of Part A (10.0 g, 52.3 mmol) and tetrahydrofuran (160 mL), and cooled to -75°C. Lithium bis(trimethylsilyl)amide (1.0 M in tetrahydrofuran, 157 mL, 157 mmol) was slowly added, keeping temperature < -60°C. Reaction was stirred for 30 minutes after the addition and was then charged with a solution of methyl chloroformate (4.0 mL, 52.3 mmol) in tetrahydrofuran (80 mL), again keeping the temperature at < -60°C. After stirring for 1 hour at -75°C, the reaction was quenched with saturated NH₄Cl, keeping temperature < -20°C. The aqueous portion froze into a solid chunk of ice.


After warming to 5°C, the mixture was extracted with ethyl acetate (3x- 200 mL). Organic phases were washed with saturated NH₄Cl (2x-100 mL) and brine (1x-100mL), then dried over Na₂SO₄ and concentrated to afford the methyl ester as a brown oil (10.7 g, 82 % crude yield). ¹H-NMR showed the desired compound.



Part C: To a solution of the methyl ester product from Part B (5.0 g, 20 mmol) and dibromo-diethylether (3.0 mL, 24.1 mmol) in dimethylformamide


-349-

(40 mL) was added 18-Crown-6 (500 mg, cat.) followed by potassium carbonate (8.3 g, 60.0 mmol). The mixture was heated at 60°C for 4 hours, after which time more potassium carbonate (1.9 g, 13.7 mmol) was added, and the reaction continued at 60°C for 14 hours. LC showed <5% starting material remained. The reaction was worked up by pouring into stirring 10% HCl_{aq} (200 mL). A gummy solid resulted that was extracted with ethyl acetate (3x- 300 mL). Organic layers were washed with brine (2x- 200 mL), dried over Na₂SO₄, and concentrated to afford the as an orange solid (3.6 g, 56% yield). ¹H-NMR showed the desired compound.



Part D: Potassium trimethylsilanolate (4.3 g, 34 mmol) was added to a solution of the pyran methyl ester product from Part C (3.6 g, 11.3 mmol) in tetrahydrofuran (30 mL). The reaction stirred overnight (about eighteen hours) at room temperature. Work up consisted of removing the tetrahydrofuran and taking the residue up in H₂O (100 mL). The solution was washed with diethyl ether (50 mL). The aqueous portion was then cooled to zero degrees C and 10% HCl was slowly added until about a pH value of 3. The acidic mixture was then extracted with ethyl acetate (3x-150 mL). The organic layers were washed with brine (1x- 100 mL), dried over Na₂SO₄, and


concentrated to afford the carboxylic acid as a yellow solid (2.5 g, 72% yield). ¹H-NMR showed the desired compound.



5


Part E: To a solution of the carboxylic acid product of Part D (2.2 g, 7.2 mmol) in dimethylacetamide (15 mL) was added N-methylmorpholine (2.4 mL, 22.0 mmol) followed by N-hydroxybenzotriazole hydrate (1.2 g, 8.6 mmol), O-(tetrahydro-2H-pyran-2-yl)hydroxylamine (1.3 g, 10.8 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (2.1 g, 10.8 mmol). The mixture was stirred overnight (about eighteen hours) and then the solvent was removed. The residue was taken up in ethyl acetate (250 mL) and washed with 5% NaHSO₄ (1x- 150 mL), saturated potassium carbonate (1x-150 mL), and brine (1x-150 mL). The organic layer was then dried over Na₂SO₄ and concentrated to afford the THP-protected hydroxamate as a brown oil (2.7 g, 93% yield). ¹H-NMR showed the desired compound.

-351-



Part F: The crude the THP-protected hydroxamate oil product of Part E (2.7 g, 6.7 mmol) was dissolved in acetonitrile (15 mL) and stirred with 10% HCl (15 mL) for 3 hours, after which time LC showed no more starting material. The solution was reduced to one-half volume, and then acetonitrile (10 mL) and trifluoroacetic acid (1 mL) were added. The solution was filtered and purified by preparatory reverse phase LC (acetonitrile/water) affording title product as a tan solid (1.7 g, 79% yield). ¹H-NMR showed the desired compound. HPLC showed product as a mixture of 70% cis and 30% trans. Mass spectroscopy showed $M^{+H}_{\text{found}} = 321$ ($M^{+H}_{\text{calc}} = 321$).

Example 8 : Preparation of tetrahydro-N-hydroxy-4-[[4-(phenylmethyl)-1-piperidinyl]-sulfonyl]-2H-pyran-4-carboxamide



Part A: To a solution of 4-benzylpiperidine (10.55 mL, 60 mmol) and triethylamine (12.5 mL, 90 mmol) in methylene chloride (250 mL) at zero degrees Celsius was added a solution of methanesulfonyl

-352-

chloride (7 mL, 60 mmol) in methylene chloride (50 mL). After one hour at ambient temperature, the solvent was removed *in vacuo*. The residue was taken up in ethyl acetate, washed with water two times, brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The product was recrystallized (ethyl acetate/hexanes) to give the sulfonamide as a beige solid (14.2 g, 94%). HRMS (ES+) M+ H⁺ calculated for C₁₃H₁₉N₁O₂S₁ : 254.1215, found 254.1211.

10 Part B: Preparation of methyl [[4-(phenylmethyl)-1-piperidinyl]sulfonyl]acetate

In dry equipment under nitrogen, the sulfonamide from Part A (2.53 g, 10 mmol) was dissolved in dry tetrahydrofuran (20 mL), chilled to minus seventy-five degrees Celsius, and a 1M solution of lithium bis(trimethylsilyl)amide (20 mL) was added while maintaining the temperature below minus sixty-five degrees. After thirty minutes at minus seventy-five degrees Celsius, a solution of methyl chloroformate (0.77mL, 10 mmol) in dry tetrahydrofuran (10 mL) was added maintaining the temperature below minus sixty degrees. After thirty minutes at minus seventy-five degrees Celsius, the reaction was quenched with saturated ammonium chloride solution (100 mL) and extracted with ethyl acetate. The combined extracts were washed with saturated ammonium chloride solution, brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo* to give the methylene sulfonamide as a yellow oil (3.05 g, 100%).

30 Part C: Preparation of methyl tetrahydro-4-[[4-(phenylmethyl)-1-piperidinyl]sulfonyl]-2H-pyran-4-carboxylate

-353-

To a solution of the methylene sulfonamide from Part B (4.1 g, 13.2 mmol) in dimethylformamide (26 mL) was added potassium carbonate (7.8 g, 56.6 mmol), bis-(2-bromoethyl)ether (1.73 mL, 13.2 mmol) and 18-
5 Crown-6 (500mg). The slurry was stirred at sixty degrees Celsius. After sixteen hours, potassium carbonate (2.0 g, 14 mmol) and bis-(2-bromoethyl)ether (0.2 mL, 1.6 mmol) were added and the reaction stirred at sixty degrees Celsius .
10 After a total of twenty-eight hours, the reaction was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water three times, brine, dried over Na₂SO₄, filtered, and concentrated in vacuo. Chromatography (on silica, ethyl
15 acetate/hexanes) provided the THP-substituted sulfonamide as a white solid (2.85 g, 57%).

Part D: Preparation of tetrahydro-4-[[4-(phenylmethyl)-1-piperidinyl]sulfonyl]-2H-
pyran-4-carboxylic acid

20 In dry equipment under nitrogen, the THP-substituted sulfonamide from Part C (2.8 g, 7.3 mmol) was dissolved in dry tetrahydrofuran (25 mL) and potassium trimethylsilylate (1.4 g, 11.0 mmol) was added at ambient temperature. After sixteen hours,
25 water (100 mL) was added and the solution concentrated in vacuo. The residue was taken up in water and extracted with ethyl acetate to remove unreacted starting material. The aqueous solution was treated with 6 N HCl until pH=1. The slurry was
30 extracted with ethyl acetate and the combined extracts washed with water, dried over Na₂SO₄, filtered, and concentrated in vacuo. The residue was heated in diethyl ether, the solid filtered and dried

-354-

to give the carboxylic acid as a white solid (1.96 g, 73%).

Part E: Preparation of tetrahydro-4-[[4-(phenylmethyl)-1-piperidinyl]sulfonyl]-N-
5 [(tetrahydro-2H-pyran-4-carboxamide


In dry equipment under nitrogen, the carboxylic acid from Part D (1.9 g, 5.18 mmol) was dissolved in dry dimethylformamide (13 mL) and the remaining reagents were added to the solution in the following
10 order: N-hydroxybenzotriazole hydrate (0.84 g, 6.2 mmol), N-methylmorpholine (1.71 mL, 15.5 mmol), O-(tetrahydro-2H-pyran-2-yl)hydroxylamine (1.88 g, 16.0 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.39 g, 7.25 mmol).
15 After two hours at ambient temperature, the reaction was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water, 5% KHSO₄, saturated NaHCO₃, brine, dried over Na₂SO₄, filtered, and concentrated in vacuo. Chromatography (on silica,
20 ethyl acetate/hexanes) provided the TPH hydroxamate as a white foam (2.22 g, 93%).

Part F: Preparation of tetrahydro-N-hydroxy-4-[[4-(phenylmethyl)-1-piperidinyl]sulfonyl]-2H-
pyran-4-carboxamide

25 To a solution of the THP hydroxamate from Part E (2.15 g, 4.6 mmol) in 1,4-dioxane (12 mL) was added 4 N HCl dioxane solution (12 mL) and methanol (12 mL). After thirty minutes at ambient temperature, the reaction was diluted with ethyl acetate and washed
30 with water, dried over Na₂SO₄, filtered, and concentrated in vacuo. The product was recrystallized (acetone/hexanes) to give the depicted compound as a white solid (1.45 g, 82%). HRMS (ES+)

M+ H⁺ calculated for C₁₈H₂₆N₂O₅ S₁: 383.1641, found 383.1640.

Example 9: Preparation of N-hydroxy-2-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-sulfonyl]butanamide



10 Part A: Preparation of 1,1-dimethylethyl 4-hydroxy-1-
piperidinecarboxylate

In dry equipment under nitrogen, 4-hydroxypiperidine (20.2 g, 0.2 mol) was dissolved in tetrahydrofuran (200 mL) and triethylamine (29 mL, 0.21 mol). A solution of di-*t*-butyldicarbonate (43.65 g, 0.2 mol) was added at such a rate that the temperature remained below thirty degrees Celsius. After stirring at ambient temperature for four hours, the reaction was concentrated *in vacuo*. The residue was taken up in ethyl acetate, washed with water, 5% KHSO_4 , saturated NaHCO_3 , brine, dried over Na_2SO_4 , filtered, and concentrated *in vacuo* to give the BOC piperidine as a white solid (37.7 g, 94%).

Part B: Preparation of 1,1-Dimethylethyl 4-[4-(trifluoromethyl)phenoxy]-1-piperidinecarboxylate

To a solution of the BOC piperidine from Part A (6.03 g, 30 mmol) in dimethylformamide (60 mL) were added cesium carbonate (9.77 g, 30 mmol) and 4-fluorobenzotrifluoride (3.8 mL, 30 mmol). The resulting slurry was stirred at ninety degrees

Celsius. After nineteen hours, cesium carbonate (3.26 g, 10 mmol) and 4-fluorobenzotrifluoride (0.95 mL, 10 mmol) were added and the reaction continued at ninety degrees Celsius. After a total of forty-six hours, the reaction was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water three times, brine, dried over Na_2SO_4 , filtered, and concentrated in vacuo. Chromatography (on silica, ethyl acetate/hexanes) provided the substituted BOC piperidine as a white solid (6.0 g, 58%).

Part C: Preparation of 4-[4-(trifluoromethyl)-phenoxy]piperidine
To a slurry of the substituted BOC piperidine from Part B (5.95 g, 17.2 mmol) in 1,4-dioxane (10 mL) was added 4 N HCl dioxane solution (17 mL). After one hour at ambient temperature, the reaction was concentrated in vacuo. The residue was slurried in diethyl ether and vacuum filtration of the resulting precipitate provided the hydrochloride salt as a white solid (4.6 g, 100%).

Part D: Preparation of 1-(methylsulfonyl)-4-[4-(trifluoromethyl)phenoxy]piperidine
To a solution of the hydrochloride salt from Part C (4.6 g, 16.9 mmol) and triethylamine (5.9 mL, 42.4 mmol) in methylene chloride (45 mL) at zero degrees Celsius was added a solution of methanesulfonyl chloride (1.97 mL, 25.4 mmol) in methylene chloride (10 mL). After one hour at ambient temperature, the solvent was removed in vacuo. The residue was taken up in ethyl acetate, washed with water two times, brine, dried over Na_2SO_4 , filtered, and concentrated in vacuo. The residue was slurried in diethyl ether and vacuum filtration of

-357-

the resulting precipitate provided the sulfonamide as an off-white solid (5.25 g, 96%).

Part E: Preparation of Methyl [[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-
5 sulfonyl]acetate

In dry equipment under nitrogen, the sulfonamide from Part D (4.2 g, 13 mmol) was dissolved in dry tetrahydrofuran (26 mL), chilled to minus seventy-five degrees Celsius, and a 1 M solution of lithium
10 bis(trimethylsilyl)amide (26 mL) was added while maintaining the temperature below minus sixty-five degrees. After thirty minutes at minus seventy-five degrees Celsius, a solution of methyl chloroformate (1.0 mL, 13 mmol) in dry tetrahydrofuran (13 mL) was
15 added while maintaining the temperature below minus sixty degrees. After thirty minutes at minus seventy-five degrees Celsius, the reaction was quenched with saturated ammonium chloride solution (100 mL) and extracted with ethyl acetate. The
20 combined extracts were washed with saturated ammonium chloride solution, brine, dried over Na₂SO₄, filtered, and concentrated in vacuo to give the methylene sulfonamide as a yellow oil (4.95 g, 100%).

Part F: Preparation of methyl 2-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-
25 sulfonyl]butanoate

To a solution of the methylene sulfonamide from Part E (6.15 g, 16 mmol) in dimethylformamide (32 mL) were added potassium carbonate (7.8 g, 56.6 mmol),
30 bis-(2-bromoethyl)amine ethyl carbamate (3.0 g, 10.75 mmol; partially purified) and 18-Crown-6 (500 mg). The resulting slurry was stirred at sixty degrees Celsius. After sixteen hours, potassium

carbonate (2.0 g, 14 mmol) was added and the reaction stirred at sixty degrees Celsius. After a total of twenty four hours, the reaction was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water three times, brine, dried over Na_2SO_4 , filtered, and concentrated *in vacuo*. Chromatography (on silica, ethyl acetate/hexanes) provided the ethyl sulfonamide as a light yellow oil (0.3 g, 10%).

10 Part G: Preparation of 2-[[4-[4-(trifluoromethyl)-
phenoxy]-1-piperidinyl]sulfonyl]butanoic acid

In dry equipment under nitrogen, the ethyl sulfonamide from Part F (0.82 g, 12.0 mmol) was dissolved in dry tetrahydrofuran (5.0 mL) and potassium trimethylsilonate (0.51 g, 4.0 mmol) was added at ambient temperature. After eighteen hours water (100 mL) was added and the solution concentrated *in vacuo*. The residue was taken up in water and extracted with ethyl acetate. The combined extracts were concentrated *in vacuo*. The residue was wetted with water and 1N HCl solution (3.0 mL) was added, the resulting slurry was extracted with ethyl acetate and the combined extracts washed with brine, dried over Na_2SO_4 , filtered, and concentrated *in vacuo* to give the carboxylic acid as an off-white solid (669 mg, 85%).

20 Part H: Preparation of N-[(tetrahydro-2H-pyran-2-yl)oxy]-2-[[4-[4-(trifluoromethyl)phenoxy]-1-
30 piperidinyl]sulfonyl]butanamide

In dry equipment under nitrogen, the carboxylic acid from Part G (0.627 g, 1.59 mmol) was dissolved in dry dimethylformamide (4 mL) and the remaining


reagents were added to the solution in the following order: N-hydroxybenzotriazole hydrate (0.26 g, 1.9 mmol), N-methylmorpholine (0.53 mL, 4.84 mmol), O-(tetrahydro-2H-pyran-2-yl)hydroxylamine (0.585 g, 5.0 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.43 g, 2.26 mmol). After fourteen hours at ambient temperature, the reaction was concentrated *in vacuo*. The residue was taken up in ethyl acetate, washed with water, 5% KHSO₄, saturated NaHCO₃, brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Chromatography (on silica, ethyl acetate/hexanes) provided the THP hydroxamate as a white solid (0.666 g, 85%).

Part I: Preparation of N-Hydroxy-2-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-sulfonyl]butanamide

To a solution of the THP hydroxamate from Part H (0.61 g, 1.23 mmol) in 1,4-dioxane (3 mL) were added 4 N HCl dioxane solution (3 mL) and methanol (3 mL). After two hours at ambient temperature, the reaction was diluted with ethyl acetate and washed with water, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The product was recrystallized (acetone/hexanes) to give the depicted compound as a white solid (490 mg, 97%). HRMS (ES+) M+ NH₄⁺ calculated for C₁₆H₂₁N₂O₅ S₁F₃ : 428.1467, found 428.1451.

Example 10: Preparation of tetrahydro-N-hydroxy-4-[[4-[4-(trifluoromethoxy)phenyl]-1-piperazinyl]sulfonyl]-2H-pyran-4-carboxamide, monohydrochloride

-360-



Part A: To a solution of tert-butyl-piperazine (25.0 g, 134 mmol) in dichloromethane (200 mL),
5 cooled to zero degrees Celsius, was added triethylamine (28.3 mL, 201 mmol) followed by methanesulfonyl chloride (15.5 mL, 201 mmol). Once the addition was complete the cooling bath was removed and the reaction mixture was stirred at
10 ambient temperature. After 1 hour the reaction mixture was concentrated *in vacuo*. The residue was partitioned between H₂O and ethyl acetate and the aqueous was further extracted with ethyl acetate. The combined organic layers were washed with
15 saturated NaCl and dried over Na₂SO₄. Concentration *in vacuo* provided the sulfonamide as an off-white solid (36.4 g, >100%).

Part B: To a solution of the sulfonamide of part A (35.0 g, 132 mmol) in tetrahydrofuran (200 mL)
20 was slowly added a solution of lithium bis(trimethylsilyl)amide (66.5 g, 397 mmol) in tetrahydrofuran (300 mL) at such a rate that the temperature never exceeded minus sixty degrees Celsius. After stirring at minus seventy-eight
25 degrees Celsius for 1.5 hours a solution of dimethyl carbonate (10.2 mL, 132 mmol) in tetrahydrofuran (100 mL) was slowly added at such a rate that the temperature of the reaction never exceeded minus sixty degrees Celsius. After stirring for 1 hour at

-361-

minus seventy-eight degrees Celsius the reaction was quenched by the addition of saturated NH_4Cl . The reaction mixture was diluted with H_2O and extracted with ethyl acetate. The combined organic layers were washed with 5% HCl , saturated NH_4Cl , saturated NaCl and dried over Na_2SO_4 . Concentration *in vacuo* provided the sulfonamide ester as an orange oil (34.1 g, 80%).


Part C: To a solution of the sulfonamide ester of part B (15.00 g, 46.53 mmol) in N,N -dimethylformamide (200 mL) was added K_2CO_3 (19.29 g, 139.59 mmol), 18-crown-6 (4.65 g) and bis(2-bromoethyl)ether (10.79 g, 46.53 mmol). The resulting mixture was heated to sixty degrees Celsius for 22 hours and then additional K_2CO_3 (19.3 g, 139.6 mmol) and bis(2-bromoethyl)ether (5.8 mL, 23.2 mmol) was added and stirring was continued at sixty degrees Celsius for 22 hours. The reaction mixture was then concentrated *in vacuo*. The residue was dissolved in acetonitrile and filtered through a pad of Celite®. The filtrate was concentrated *in vacuo* and the residue was partitioned between H_2O and ethyl acetate. The organic layers were washed with saturated NaHCO_3 , saturated NaCl and dried over Na_2SO_4 . Chromatography (on silica, ethyl acetate/hexanes) provided the cyclized ester as a pale yellow solid (7.23 g, 40%).

Part D: The cyclized ester of part C (7.23 g, 18.42 mmol) was treated with a solution of 4N HCl in dioxane (46 mL). After stirring at ambient temperature for 1 hour the reaction mixture was concentrated *in vacuo* to provide the piperazine sulfonamide as a tan solid (5.83 g, 96%).

- Part E: To a suspension of the piperazine sulfonamide of part D (2.15 g, 6.53 mmol) in toluene (15 mL), precooled to zero degrees Celsius, was added sodium tert-butoxide (1.43 g, 14.93 mmol). Once the addition was complete the cooling bath was removed and the reaction mixture was allowed to warm to ambient temperature at which time 4-(trifluoromethoxy)bromobenzene (1.50 g, 6.22 mmol), BINAP (0.116 g, 0.187 mmol) and tris(dibenzylideneacetone)dipalladium (0) (0.057 g, 0.062 mmol) were added. The resulting mixture was stirred at eighty degrees Celsius for 12 hours and then concentrated in vacuo. The residue was heated in boiling methanol, decanted from the salts and concentrated in vacuo. The residue was dissolved in H₂O and acidified (pH-2) with 1N HCl. The aqueous layer was then extracted with ethyl acetate. The combined organic layers were dried over Na₂SO₄ to give the acid as a yellow solid (1.06 g, 39%).
- Part F: To a solution of the acid of part E (1.06 g, 2.42 mmol) in N,N-dimethylformamide (5.0 mL) was added 1-hydroxybenzotriazole (0.392 g, 2.90 mmol), N-methylmorpholine (0.790 mL, 7.26 mmol), O-(tetrahydropuranyl) hydroxylamine (0.425 g, 3.63 mmol) and 1-3-[(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (0.650 g, 3.39 mmol). After stirring at ambient temperature for 7 hours the reaction was diluted with H₂O and extracted with ethyl acetate. The combined organic layers were washed with saturated NaHCO₃, saturated NaCl and dried over Na₂SO₄. Chromatography (on silica, ethyl acetate with 5% methanol/hexanes) provided the protected hydroxamate as a white solid (1.02 g, 78%).

Part G: The protected hydroxamate of part F (1.02 g, 1.90 mmol) was treated with a solution of 2N HCl in diethyl ether (9.5 mL) and methanol (0.77 mL, 18.97 mmol). The reaction mixture became gelatinous and a solution of 4 N HCl in dioxane (4.0 mL) was added. After stirring at ambient temperature for 2 hours the solids were collected by filtration, washing with diethyl ether, to give the title compound as a white solid (0.869 g, 93%). MS MH⁺ calculated for C₁₇H₂₃O₆N₃S₁F₃: 454, found 454.

Example 11: Preparation of N-hydroxy-2-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-sulfonyl]acetamide



Part A: Preparation of 1,1-Dimethylethyl 4-hydroxy-1-piperidinecarboxylate

In dry equipment under nitrogen, 4-hydroxypiperidine (20.2 g, 0.2 mol) was dissolved in tetrahydrofuran (200 mL) and triethylamine (29 mL, 0.21 mol). A solution of di-t-butyl dicarbonate (43.65 g, 0.2 mol) was added at such a rate that the temperature remained below thirty degrees Celsius. After stirring at ambient temperature for four hours, the reaction was concentrated *in vacuo*. The residue was taken up in ethyl acetate, washed with water, 5% KHSO₄, saturated NaHCO₃, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and

-364-

concentrated *in vacuo* to give the BOC piperidine as a white solid (37.7 g, 94%).

Part B: Preparation of 1,1-dimethylethyl 4-[4-(trifluoromethyl)phenoxy]-1-
5 piperidinecarboxylate

To a solution of the BOC piperidine from part A (6.03 g, 30 mmol) in dimethylformamide (60 mL) were added cesium carbonate (9.77 g, 30 mmol) and 4-fluorobenzotrifluoride (3.8 mL, 30 mmol). The slurry
10 was stirred at ninety degrees Celsius. After nineteen hours cesium carbonate (3.26g, 10 mmol) and 4-fluorobenzotrifluoride (0.95ml mL, 10 mmol) were added and the reaction continued at ninety degrees Celsius. After a total of forty six hours, the
15 reaction was concentrated *in vacuo*. The residue was taken up in ethyl acetate, washed with water three times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Chromatography (on silica, ethyl acetate/hexanes)
20 provided the substituted BOC piperidine as a white solid (6.0 g, 58%).

Part C: Preparation of 4-[4-(trifluoromethyl)-
phenoxy]piperidine

A solution of 4 N HCl dioxane solution (17 mL)
25 was added to a slurry of the substituted BOC piperidine from part B (5.95 g, 17.2 mmol) in 1,4-dioxane (10 mL). After one hour at ambient temperature the reaction was concentrated *in vacuo*. The residue was slurried in diethyl ether and vacuum
30 filtration of the resulting precipitate provided the hydrochloride salt as a white solid (4.6 g, 100%).

Part D: Preparation of 1-(Methylsulfonyl)-4-[4-(trifluoromethyl)phenoxy]piperidine

-365-

A solution of methane sulfonyl chloride (1.97 mL, 25.4 mmol) in dichloromethane (10 mL) was added to a solution of the hydrochloride salt from part C (4.6 g, 16.9 mmol) and triethylamine (5.9 mL, 42.4 mmol) in dichloromethane (45 mL) at zero degrees Celsius. After one hour at ambient temperature, the solvent was stripped *in vacuo*. The residue was taken up in ethyl acetate, washed with water two times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The residue was slurried in diethyl ether and vacuum filtration of the resulting precipitate provided the sulfonamide as an off-white solid (5.25 g, 96%).

Part E: Preparation of methyl [[4-[4-(trifluoromethyl)-phenoxy]-1-piperidinyl]-sulfonylacetate

In dry equipment under nitrogen, the sulfonamide from part D (4.2 g, 13 mmol) was dissolved in dry tetrahydrofuran (26 mL), chilled to minus seventy-five degrees Celsius, and a 1 M solution of lithium bis(trimethylsilyl)amide (26 mL) was added maintaining the temperature below minus sixty five degrees. After thirty minutes at minus seventy-five degrees Celsius, a solution of methyl chloroformate (1.0 mL, 13 mmol) in dry tetrahydrofuran (13 mL) was added while maintaining the temperature below minus sixty degrees. After thirty minutes at minus seventy-five degrees Celsius, the reaction was quenched with saturated ammonium chloride solution (100 mL) and extracted with ethyl acetate. The combined extracts were washed with saturated ammonium chloride solution, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and

-366-

concentrated *in vacuo* to give the methylene sulfonamide as a yellow oil (4.95 g, 100%).

Part F: Preparation of [[4-[4-(trifluoromethyl)-phenoxy]-1-piperidinyl]sulfonyl]acetic acid

5 In dry equipment under nitrogen, the methylene sulfonamide from part E (1.52 g, 4.0 mmol) was dissolved in dry tetrahydrofuran (10 mL) and potassium trimethylsilylate (1.03 g, 8.0 mmol) was added at ambient temperature. After fifteen hours, 10 water (50 mL) was added, and at 5 degrees Celsius 6 N HCl was added until pH=1. The slurry was extracted with ethyl acetate and the combined extracts washed with water, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The residue was heated in 15 diethyl ether, the solid filtered and dried to give the carboxylic acid as a white solid (1.25 g, 85%).

Part G: Preparation of N-[(tetrahydro-2H-pyran-2-yl)oxy]-2-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl] acetamide

20 In dry equipment under nitrogen, the carboxylic acid from part F (1.2 g, 3.3 mmol) was dissolved in dry dimethylformamide (8 mL) and the remaining reagents were added to the solution in the following order: N-hydroxybenzotriazole hydrate (0.53 g, 3.9 25 mmol), N-methylmorpholine (1.08 mL, 9.8 mmol), O-(tetrahydro-2H-pyran-2-yl)hydroxylamine (1.19 g, 10.1 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.88 g, 4.6 mmol). After two hours at ambient temperature, the reaction 30 was concentrated *in vacuo*. The residue was taken up in ethyl acetate, washed with water, 5% KHSO₄, saturated NaHCO₃, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in*

vacuo. Chromatography (on silica, ethyl acetate/hexanes) provided the THP hydroxamate as a white solid (1.16 g, 76%).

Part H: Preparation of N-hydroxy-2-[[4-[4-


5 (trifluoromethyl)phenoxy]-1-piperidinyl]-
sulfonyllacetamide

To a solution of the THP hydroxamate from part G (1.11 g, 2.4 mmol) in 1,4-dioxane (6 mL) were added 4 N HCl dioxane solution (6 mL) and methanol (6 mL).

10 After ninety minutes at ambient temperature, the reaction was diluted with ethyl acetate and washed with water, dried over Na₂SO₄, filtered, and concentrated in vacuo. The product was recrystallized (acetone/hexanes) to give the title
15 compound as a white solid (0.60 g, 67%). HRMS (ES+) M+ H⁺ calculated for C₁₄H₁₇F₃N₂O₅S₁: 383.0889, found 383.0885.

Example 12: Preparation of N-hydroxy-1-(2-

20 methoxyethyl)-4-[[4-[4-(trifluoromethyl)-
phenoxy]-1-piperidinyl)sulfonyl]-4-
piperidinecarboxamide, monohydrochloride



25

Part A: Preparation of 1,1-Dimethylethyl 4-hydroxy-1-
piperidinecarboxylate

-368-

In dry equipment under nitrogen, 4-hydroxypiperidine (20.2 g, 0.2 mol) was dissolved in tetrahydrofuran (200 mL) and triethylamine (29 mL, 0.21 mol). A solution of di-t-butylidicarbonate (43.65 g, 0.2 mol) was added at such a rate that the temperature remained below thirty degrees Celsius. After stirring at ambient temperature for four hours, the reaction was concentrated *in vacuo*. The residue was taken up in ethyl acetate, washed with water, 5% KHSO₄, saturated NaHCO₃, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo* to give the BOC piperidine as a white solid (37.7 g, 94%).

Part B: Preparation of 1,1-dimethylethyl 4-[4-(trifluoromethyl)phenoxy]-1-piperidinecarboxylate

To a solution of the BOC piperidine from part A (6.03 g, 30 mmol) in dimethylformamide (60 mL) were added cesium carbonate (9.77 g, 30 mmol) and 4-fluorobenzotrifluoride (3.8 mL, 30 mmol). The slurry was stirred at ninety degrees Celsius. After nineteen hours cesium carbonate (3.26g, 10 mmol) and 4-fluorobenzotrifluoride (0.95ml mL, 10 mmol) were added and the reaction continued at ninety degrees Celsius. After a total of forty six hours, the reaction was concentrated *in vacuo*. The residue was taken up in ethyl acetate, washed with water three times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Chromatography (on silica, ethyl acetate/hexanes) provided the substituted BOC piperidine as a white solid (6.0 g, 58%).

Part C: Preparation of 4-[4-(trifluoromethyl)-

phenoxy)piperidine

To a slurry of the substituted BOC piperidine from part B (5.95 g, 17.2 mmol) in 1,4-dioxane (10 mL) was added 4 N HCl dioxane solution (17 mL).

- 5 After one hour at ambient temperature the reaction was concentrated *in vacuo*. The residue was slurried in diethyl ether and vacuum filtration of the resulting precipitate provided the hydrochloride salt as a white solid (4.6 g, 100%).

10 Part D: Preparation of 1-(methylsulfonyl)-4-[4-(trifluoromethyl)phenoxy]piperidine

- To a solution of the hydrochloride salt from part C (4.6 g, 16.9 mmol) and triethylamine (5.9 mL, 42.4 mmol) in dichloromethane (45 mL) at zero degrees Celsius was added a solution of methane sulfonyl chloride (1.97 mL, 25.4 mmol) in dichloromethane (10 mL). After one hour at ambient temperature, the solvent was stripped *in vacuo*. The residue was taken up in ethyl acetate, washed with water two times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The residue was slurried in diethyl ether and vacuum filtration of the resulting precipitate provided the sulfonamide as an off-white solid (5.25 g, 96%).
- 20
- 25

Part E: Preparation of methyl [[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-sulfonylacetate

- In dry equipment under nitrogen, the sulfonamide from part D (4.2 g, 13 mmol) was dissolved in dry tetrahydrofuran (26 mL), chilled to minus seventy-five degrees Celsius, and a 1 M solution of lithium bis(trimethylsilyl)amide (26 mL) was added
- 30

maintaining the temperature below minus sixty five degrees. After thirty minutes at minus seventy-five degrees Celsius, a solution of methyl chloroformate (1.0 mL, 13 mmol) in dry tetrahydrofuran (13 mL) was added maintaining the temperature below minus sixty degrees. After thirty minutes at minus seventy-five degrees Celsius, the reaction was quenched with saturated ammonium chloride solution (100 mL) and extracted with ethyl acetate. The combined extracts were washed with saturated ammonium chloride solution, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo* to give the methylene sulfonamide as a yellow oil (4.95 g, 100%).

Part F: Preparation of methyl 1-(phenylmethyl)-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-sulfonyl]-4-piperidinecarboxylate

To a solution of the methylene sulfonamide from part E (1.14 g, 3 mmol) in dimethylformamide (6 mL) were added potassium carbonate (1.24 g, 9 mmol), bis-(2-chloroethyl)benzyl amine (0.7 g, 3 mmol; and 18-Crown-6 (500mg). The slurry was stirred at sixty degrees Celsius. After sixteen hours the reaction was concentrated *in vacuo*. The residue was taken up in ethyl acetate, washed with water three times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Chromatography (on silica, ethyl acetate/hexanes) provided the piperidine sulfonamide as a white solid (950 mg, 59%).

Part G: Preparation of Methyl 1-(2-methoxyethyl)-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxylate

To a slurry of the piperidine sulfonamide from part F (420 mg, 0.8 mmol) in methanol (15 mL) was added ammonium formate (147 mg, 2.3 mmol). The system was purged with nitrogen for 10 minutes. The nitrogen stream was removed and palladium on carbon (80 mg of 10 weight % on activated carbon, 50%water) was added. The reaction was refluxed for forty five minutes, cooled, filtered through Celite under nitrogen, and concentrated in vacuo. The residue was dissolved in dry dimethylformamide (5 mL) and potassium carbonate (150 mg, 1.07 mmol) and 2-bromoethyl methyl ether (100 uL, 1.07 mmol) were added. The reaction was stirred at thirty five degrees Celsius for thirty hours and then concentrated in vacuo. The residue was taken up in ethyl acetate and filtered through Celite. The filtrate washed with water two times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo. The product was recrystallized (methanol) to give the N-methoxyethyl piperidine sulfonamide as an off-white solid (190 mg, 53%).

Part H: Preparation of 1-(2-methoxyethyl)-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-sulfonyl]-4-piperidinecarboxylic acid

In dry equipment under nitrogen, the N-methoxyethyl piperidine sulfonamide from part G (1.51 g, 2.97 mmol) was dissolved in dry tetrahydrofuran (30 mL) and potassium trimethylsilonate (1.27 g, 8.91 mmol) was added at ambient temperature. After five hours, water (10 mL) was added and at 5 degrees Celsius, 6 N HCl was added until pH=1. The slurry was filtered, washed with water and dried in vacuo to

give the carboxylic acid as a white solid (1.47 g, 100%).

Part I: Preparation of 1-(2-methoxyethyl)-N-

5 [(tetrahydro-2H-pyran-2-yl)oxy]-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-
sulfonyl]-4-piperidinecarboxamide

In dry equipment under nitrogen, the carboxylic acid from part H (1.4 g, 2.83 mmol) was dissolved in dry dimethylformamide (12 mL) and the remaining
10 reagents were added to the solution in the following order: N-hydroxybenzotriazole hydrate (0.46 g, 3.4 mmol), N-methylmorpholine (0.93 mL, 8.5 mmol), O-(tetrahydro-2H-pyran-2-yl)hydroxylamine (1.03 g, 8.8 mmol), and 1-(3-dimethylaminopropyl)-3-
15 ethylcarbodiimide hydrochloride (0.76 g, 4.0 mmol). The reaction was stirred at forty five degrees Celsius. After forty eight hours, the reaction was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water, 5% KHSO₄, saturated
20 NaHCO₃, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo. Chromatography (on silica, ethyl acetate/hexanes) provided the THP hydroxamate as a white solid (800 mg, 48%).


25 Part J: Preparation of N-hydroxy-1-(2-methoxyethyl)-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-4-
piperidinecarboxamide, monohydrochloride

To a solution of the THP hydroxamate from part I
30 (760 mg, 1.28 mmol) in 1,4-dioxane (3 mL) was added 4N HCl dioxane solution (3.2 mL) and methanol (3.5 mL). After thirty minutes at ambient temperature the reaction was poured into 100 mL acetonitrile. The

-373-

slurry was filtered under nitrogen, washed with acetonitrile and dried *in vacuo* to give the title compound as an off white solid (0.62 g, 89%). HRMS (ES+) $M + H^+$ calculated for $C_{21}H_{30}F_3N_3O_6S_1$: 510.1886, found 510.1862.

Example 13: Preparation of tetrahydro-N-hydroxy-4-[[4-(4-nitrophenoxy)-1-piperidinyl]sulfonyl]-2H-pyran-4-carboxamide



Part A: Preparation of 1,1-dimethylethyl 4-hydroxy-1-piperidinecarboxylate

In dry equipment under nitrogen, 4-hydroxypiperidine (20.2 g, 0.2 mol) was dissolved in tetrahydrofuran (200 mL) and triethylamine (29 mL, 0.21 mol). A solution of di-*t*-butyldicarbonate (43.65 g, 0.2 mol) was added at such a rate that the temperature remained below thirty degrees Celsius. After stirring at ambient temperature for four hours, the reaction was concentrated *in vacuo*. The residue was taken up in ethyl acetate, washed with water, 5% $KHSO_4$, saturated $NaHCO_3$, saturated sodium chloride solution, dried over Na_2SO_4 , filtered, and concentrated *in vacuo* to give the BOC piperidine as a white solid (37.7 g, 94%).

Part B: Preparation of 1,1-dimethylethyl 4-(4-nitrophenoxy)-1-piperidinyl carboxylate

-374-

To a solution of the BOC piperidine from part A (10.05 g, 0.05 mol) in dimethylformamide (100 mL) were added cesium carbonate (16.3 g, 0.05 mol) and 4-fluoronitrobenzene (5.3 mL, 0.05 mol). The slurry
5 was stirred at eighty five degrees Celsius. After eight hours cesium carbonate (1.6g, 5 mmol) was added and the reaction continued at ninety degrees Celsius. After a total of twenty four hours, the reaction was concentrated in vacuo. The residue was taken up in
10 ethyl acetate, washed with water three times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo. Chromatography (on silica, ethyl acetate/hexanes) provided the substituted BOC piperidine as an yellow solid (13.6
15 g, 84%).

Part C: Preparation of 4-(4-nitrophenoxy)piperidine,
monohydrochloride

To a slurry of the substituted BOC piperidine from part B (6.44 g, 20 mmol) in 1,4-dioxane (5 mL)
20 was added 4 N HCl dioxane solution (20 mL). After two hour at ambient temperature the reaction was concentrated in vacuo. The residue was slurried in diethyl ether and vacuum filtration of the resulting precipitate provided the hydrochloride salt as a
25 light yellow solid (5.2 g, 100%).

Part D: Preparation of 1-(methylsulfonyl)-4-(4-
nitrophenoxy)piperidine

To a solution of the hydrochloride salt from part C (5.17 g, 20 mmol) and triethylamine (7.0 mL,
30 50 mmol) in dichloromethane (80 mL) at zero degrees Celsius was added a solution of methane sulfonyl chloride (2.32 mL, 30 mmol) in dichloromethane (20 mL). After one hour at ambient temperature, the

-375-

WO 00/46221

solvent was stripped in vacuo. The residue was taken up in ethyl acetate, washed with water two times, saturated sodium chloride solution, dried over Na_2SO_4 , filtered, and concentrated in vacuo. The product was recrystallized (ethyl acetate/hexanes) to give the sulfonamide as a yellow solid (4.57 g, 76%).

5 Part E: Preparation of methyl [(4-(4-nitrophenoxy)-1-piperidinyl)sulfonyl]acetate

In dry equipment under nitrogen, the sulfonamide from part D (4.48 g, 14.9 mmol) was dissolved in dry tetrahydrofuran (30 mL), chilled to minus seventy-five degrees Celsius, and a 1 M solution of lithium bis(trimethylsilyl)amide (30 mL) was added maintaining the temperature below minus sixty-five degrees. After thirty minutes at minus sixty-five degrees Celsius, a solution of methyl chloroformate (1.15 mL, 14.9 mmol) in dry tetrahydrofuran (15 mL) was added while maintaining the temperature below minus sixty degrees. After thirty minutes at minus seventy-five degrees Celsius, the reaction was quenched with saturated ammonium chloride solution (100 mL) and extracted with ethyl acetate. The combined extracts were washed with saturated sodium chloride solution, dried over Na_2SO_4 , filtered, and concentrated in vacuo. Chromatography (on silica, ethyl acetate/hexanes) provided the methylene sulfonamide as a white solid (3.9 g, 73%).

25 Part F: Preparation of methyl tetrahydro-4-[(4-(4-nitrophenoxy)-1-piperidinyl)sulfonyl]-2H-pyran-4-carboxylate

To a solution of the methylene sulfonamide from part E (3.73 g, 10.4 mmol) in dimethylformamide (20

30

-376-

mL) was added potassium carbonate (5.75 g, 41.6 mmol), bis-(2-bromoethyl)ether (1.36 mL, 10.4 mmol) and 18-Crown-6 (500 mg). The slurry was stirred at sixty degrees Celsius. After sixteen hours the
5 reaction was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water three times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo. The product was recrystallized (ethyl acetate) to give
10 the tetrahydropyran sulfonamide as a white solid (2.1 g, 47%).

Part G: Preparation of tetrahydro-4-[[4-(4-nitrophenoxy)-1-piperidinyl)sulfonyl]-2H-pyran-4-carboxylic acid

15 In dry equipment under nitrogen, the tetrahydropyran sulfonamide from part F (2.04 g, 4.77 mmol) was dissolved in dry tetrahydrofuran (12 mL) and potassium trimethylsilonate (2.04 g, 14.3 mmol) was added at ambient temperature. After five hours
20 water (50 mL) was added and the tetrahydrofuran was stripped off in vacuo. The residue was washed with ethyl acetate, the layers were separated, the aqueous was chilled to 5 degrees Celsius and 6 N HCl was added until pH=1. The slurry was filtered, washed
25 with water and dried in vacuo to give the carboxylic acid as a white solid (1.66 g, 84%).

Part H: Preparation of tetrahydro-4-[[4-(4-nitrophenoxy)-1-piperidinyl)sulfonyl]-N-((tetrahydro-2H-pyran-2-yl)oxy)-2H-pyran-4-carboxamide

30

In dry equipment under nitrogen, the carboxylic acid from part G (1.63 g, 3.9 mmol) was dissolved in dry dimethylformamide (10 mL) and the remaining


-377-

reagents were added to the solution in the following order: N-hydroxybenzotriazole hydrate (0.640 g, 4.7 mmol), N-methylmorpholine (1.3 mL, 11.8 mmol), O-(tetrahydro-2H-pyran-2-yl)hydroxylamine (1.43 g, 12.2 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.1 g, 5.5 mmol). The reaction was stirred at forty five degrees Celsius. After forty eight hours, the reaction was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water, 5% KHSO₄, saturated NaHCO₃, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo. Chromatography (on silica, ethyl acetate/methanol/hexanes) provided the THP hydroxamate as a white solid (1.89 g, 95%).

Part I: Preparation of tetrahydro-N-hydroxy-4-[[4-(4-nitrophenoxy)-1-piperidinyl]sulfonyl]-2H-pyran-4-carboxamide

To a solution of the THP hydroxamate from part H (1.85 g, 3.6 mmol) in 1,4-dioxane (9 mL) were added 4 N HCl dioxane solution (9 mL) and methanol (1 mL). After fifteen minutes at ambient temperature the product precipitated from the reaction. The reaction was diluted with diethyl ether, the solids filtered under nitrogen and dried in vacuo to give the title compound as a white solid (1.4 g, 93%). HRMS (ES+) M+ H⁺ calculated for C₁₇H₂₃N₃O₈S₁: 430.1284, found 430.1314.

Example 14: Preparation of N-hydroxy-1-(phenylmethyl)-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide, monohydrochloride



Part A: Preparation of 1,1-dimethylethyl 4-
hydroxy-1-piperidinecarboxylate

In dry equipment under nitrogen, 4-

5 hydroxypiperidine (20.2 g, 0.2 mol) was dissolved in tetrahydrofuran (200 mL) and triethylamine (29 mL, 0.21 mol). A solution of di-*t*-butyldicarbonate (43.65 g, 0.2 mol) was added at such a rate that the temperature remained below thirty degrees Celsius.

10 After stirring at ambient temperature for four hours, the reaction was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water, 5% KHSO_4 , saturated NaHCO_3 , saturated sodium chloride solution, dried over Na_2SO_4 , filtered, and

15 concentrated in vacuo to give the BOC piperidine as a white solid (37.7 g, 94%).

Part B: Preparation of 4-(methylsulfonyl)hydroxy-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester

20 To a solution of the BOC piperidine of part A
(5.00 g, 24.84 mmol) in dichloromethane (50 mL) at
zero degrees Celsius, was added triethylamine (3.81
mL, 27.32 mmol) followed by methane sulfonyl chloride
(2.02 mL, 26.08 mmol). Once the addition was
25 complete the cooling bath was removed. After
stirring for two hours the reaction mixture was
concentrated in vacuo. The residue was taken up in
ethyl acetate, washed with water two times, saturated

sodium chloride solution, dried over Na_2SO_4 , filtered, and concentrated *in vacuo* to provide the mesylate as an off-white solid (7.34 g, >100%).

Part C: Preparation of 4-[4-(trifluoromethoxy)-
5 phenoxy]-1-piperidinecarboxylic acid, 1,1-
 dimethylethyl ester

In dry equipment under nitrogen, 4-trifluoromethoxyphenol (10.15 g, 57 mmol) was dissolved in dry dimethylformamide (125 mL) and at
10 minus five degrees Celsius sodium hydride (2.74g, 68.4 mmol of the 60% oil dispersion) was added and the ice bath was removed. After one hour at ambient temperature, the mesylate from part B (15.9 g, 57 mmol) was added and the reaction stirred at eighty
15 degrees Celsius. After stirring at ambient temperature for four hours, the reaction was concentrated *in vacuo*. The residue was taken up in diethyl ether, washed with water, saturated sodium chloride solution, dried over Na_2SO_4 , filtered, and
20 concentrated *in vacuo* to provide the substituted BOC-piperidine as a beige solid (20.6 g, 100%).

Part D: Preparation of 4-[4-(trifluoromethoxy)-
 phenoxy]piperidine

At fifteen degrees Celsius, 4 N HCl in dioxanes
25 (125 mL) was slowly added to the substituted BOC-piperidine from part C (20.6 g, 57 mmol) and stirred for ninety minutes. The reaction was concentrated *in vacuo*. The residue was dissolved in water (150 mL) and washed two times with ethyl acetate. The aqueous
30 solution was cooled to five degrees Celsius and the pH adjusted to eleven with 5 N sodium hydroxide solution and extracted with ethyl acetate. The ethyl acetate was dried over Na_2SO_4 , filtered, and

concentrated *in vacuo* to provide the substituted piperidine as a beige solid (11.9 g, 80%).

Part E: Preparation of 1-(methylsulfonyl)-4-[4-(trifluoromethoxy)phenoxy]piperidine

5

The substituted piperidine from part D (11.5 g, 44.1 mmol) was dissolved in dichloromethane (125 mL) with triethylamine (12.3 mL, 88.1 mmol), and at zero degrees Celsius a solution of methane sulfonyl chloride (5.1 mL, 66.1 mmol) in dichloromethane (20 mL) was added. After one hour at ambient temperature, the solvent was stripped *in vacuo*. The residue was taken up in ethyl acetate, washed with water two times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The residue was recrystallized (ethyl acetate/hexanes) to give the sulfonamide as an off-white solid (10.77 g, 72%).

Part F: Preparation of methyl [[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]-sulfonyl]-4-piperidinecarboxylate

In dry equipment under nitrogen, the sulfonamide from part E (10.77 g, 31.8 mmol) was dissolved in dry tetrahydrofuran (64 mL), chilled to minus seventy-five degrees Celsius, and a 1 M solution of lithium bis(trimethylsilyl)amide (80 mL) was added maintaining the temperature below minus sixty five degrees. After thirty minutes at minus seventy-five degrees Celsius, a solution of methyl chloroformate (2.45 mL, 31.8 mmol) in dry tetrahydrofuran (32 mL) was added maintaining the temperature below minus sixty degrees. After thirty minutes at minus seventy-five degrees Celsius, the reaction was

-381-

quenched with saturated ammonium chloride solution (125 mL) and extracted with ethyl acetate. The combined extracts were washed with saturated ammonium chloride solution, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo* to give the methylene sulfonamide as an yellow oil (12.69 g, 100%).

Part G: Preparation of Methyl 1-(phenylmethyl)-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxylate

To a solution of the methylene sulfonamide from part F (20.46 g, 51.5 mmol) in dimethylformamide (90 mL) was added potassium carbonate (21.3 g, 154.7 mmol), bis-(2-chloroethyl)benzyl amine (12.0 g, 51.5 mmol; and 18-Crown-6 (700 mg). The slurry was stirred at sixty degrees Celsius. After twenty four hours the reaction was concentrated *in vacuo*. The residue was taken up in ethyl acetate, washed with water three times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The residue was recrystallized (methanol) to give the N-benzyl piperidine sulfonamide as a white solid (15 g, 52%).

Part H: Preparation of 1-(Phenylmethyl)-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]-sulfonyl]-4-piperidinecarboxylic acid

In dry equipment under nitrogen, the N-benzyl piperidine sulfonamide from part G (1.5 g, 2.7 mmol) was dissolved in dry tetrahydrofuran (30 mL) and potassium trimethylsilonate (1.15 g, 8.1 mmol) was added at ambient temperature. After twenty four hours, water (10 mL) was added and at 5 degrees

Celsius, 6 N HCl was added until pH=7. The slurry was filtered, washed with water and dried in vacuo to give the carboxylic acid as a white solid (1.24 g, 85%).

5 Part I: Preparation of 1-(phenylmethyl)-N-
[(tetrahydro-2H-pyran-2-yl)oxy-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]-
sulfonyl]-4-piperidinecarboxylate

In dry equipment under nitrogen, the carboxylic
10 acid from part H (1.2 g, 2.2 mmol) was dissolved in dry dimethylformamide (6 mL) and the remaining reagents were added to the solution in the following order: N-hydroxybenzotriazole hydrate (0.36 g, 2.66 mmol), N-methylmorpholine (0.73 mL, 6.64 mmol), O-
15 (tetrahydro-2H-pyran-2-yl)hydroxylamine (0.83 g, 6.86 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.59 g, 3.1 mmol). The reaction was stirred at forty five degrees Celsius. After two hours, the reaction was
20 concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water, 5% KHSO₄, saturated NaHCO₃, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo. Chromatography (on silica, ethyl acetate/hexanes)
25 provided the THP hydroxamate as a white solid (1.24 mg, 88%).


Part J: Preparation of N-Hydroxy-1-(phenylmethyl)-4-
[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]sulfonyl]-4-
30 piperidinecarboxamide, monohydrochloride

To a solution of the THP hydroxamate from part I (1.18 g, 1.84 mmol) in 1,4-dioxanes (5 mL) was added 4 N HCl dioxane solution (5 mL) and methanol (1 mL).

-383-

After thirty minutes, the reaction was diluted with diethyl ether, the solids filtered under nitrogen and dried in vacuo to give the title compound as a white solid (0.96 g, 88%). HRMS (ES+) $M + H^+$ calculated for $C_{25}H_{30}F_3N_3O_6S_1$: 558.1886, found 558.1961.

Example 15: Preparation of N-hydroxy-1-(2-methoxyethyl)-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide, monohydrochloride



Part A: Preparation of 1,1-dimethylethyl 4-hydroxy-1-piperidinecarboxylate

In dry equipment under nitrogen, 4-hydroxypiperidine (20.2 g, 0.2 mol) was dissolved in tetrahydrofuran (200 mL) and triethylamine (29 mL, 0.21 mol). A solution of di-*t*-butyldicarbonate (43.65 g, 0.2 mol) was added at such a rate that the temperature remained below thirty degrees Celsius. After stirring at ambient temperature for four hours, the reaction was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water, 5% $KHSO_4$, saturated $NaHCO_3$, saturated sodium chloride solution, dried over Na_2SO_4 , filtered, and

-384-

concentrated in vacuo to give the BOC piperidine as a white solid (37.7 g, 94%).

Part B: Preparation of 4-(methylsulfonyl)hydroxy-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester

5 To a solution of the BOC piperidine of part A (5.00 g, 24.84 mmol) in dichloromethane (50 mL) at zero degrees Celsius, was added triethylamine (3.81 mL, 27.32 mmol) followed by methane sulfonyl chloride
10 (2.02 mL, 26.08 mmol). Once the addition was complete the cooling bath was removed. After stirring for two hours the reaction mixture was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water two times, saturated
15 sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo to provide the mesylate as an off-white solid (7.34 g, >100%).

Part C: Preparation of 4-[4-(trifluoromethoxy)-phenoxy]-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester

20 In dry equipment under nitrogen, 4-trifluoromethoxyphenol (10.15 g, 57 mmol) was dissolved in dry dimethylformamide (125 mL), and at minus five degrees Celsius sodium hydride (2.74g, 25 68.4 mmol of the 60% oil dispersion) was added and the ice bath was removed. After one hour at ambient temperature, the mesylate from part B (15.9 g, 57 mmol) was added and the reaction stirred at eighty degrees Celsius. After stirring at ambient
30 temperature for four hours, the reaction was concentrated in vacuo. The residue was taken up in diethyl ether, washed with water, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and

concentrated *in vacuo* to provide the substituted BOC-piperidine as a beige solid (20.6 g, 100%).

Part D: Preparation of 4-[4-(trifluoromethoxy)-
phenoxy]piperidine

5 At fifteen degrees Celsius, 4 N HCl in dioxane (125 mL) was slowly added to the substituted BOC-piperidine from part C (20.6 g, 57 mmol) and stirred for ninety minutes. The reaction was concentrated *in vacuo*. The residue was dissolved in water (150 mL)
10 and washed two times with ethyl acetate. The aqueous solution was cooled to five degrees Celsius and the pH adjusted to eleven with 5 N sodium hydroxide solution and extracted with ethyl acetate. The ethyl acetate was dried over Na₂SO₄, filtered, and
15 concentrated *in vacuo* to provide the substituted piperidine as a beige solid (11.9 g, 80%).

Part E: Preparation of 1-(methanesulfonyl)-4-[4-
(trifluoromethoxy)phenoxy]piperidine

20 The substituted piperidine from part D (11.5 g, 44.1 mmol) was dissolved in dichloromethane (125 mL) with triethylamine (12.3 mL, 88.1 mmol), and at zero degrees Celsius, a solution of methane sulfonyl chloride (5.1 mL, 66.1 mmol) in dichloromethane (20
25 mL) was added. After one hour at ambient temperature, the solvent was stripped *in vacuo*. The residue was taken up in ethyl acetate, washed with water two times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in*
30 *vacuo*. The residue was recrystallized (ethyl acetate/hexanes) to give the sulfonamide as an off-white solid (10.77 g, 72%).

Part F: Preparation of methyl [[4-[4-

(trifluoromethoxy)phenoxy]-1-
piperidinyl)sulfonyl]-4-piperidinecarboxylate

In dry equipment under nitrogen, the sulfonamide
5 from part E (10.77 g, 31.8 mmol) was dissolved in dry
tetrahydrofuran (64 mL), chilled to minus seventy-
five degrees Celsius, and a 1 M solution of lithium
bis(trimethylsilyl)amide (80 mL) was added
maintaining the temperature below minus sixty five
10 degrees. After thirty minutes at minus seventy-five
degrees Celsius, a solution of methyl chloroformate
(2.45 mL, 31.8 mmol) in dry tetrahydrofuran (32 mL)
was added maintaining the temperature below minus
sixty degrees. After thirty minutes at minus
15 seventy-five degrees Celsius, the reaction was
quenched with saturated ammonium chloride solution
(125 mL) and extracted with ethyl acetate. The
combined extracts were washed with saturated ammonium
chloride solution, saturated sodium chloride
20 solution, dried over Na₂SO₄, filtered, and
concentrated in vacuo to give the methylene
sulfonamide as an yellow oil (12.69 g, 100%).

Part G: Preparation of methyl 1-(phenylmethyl)-4-[[4-
[4- (trifluoromethoxy)phenoxy]-1-
25 piperidinyl)sulfonyl]-4-piperidinecarboxylate

To a solution of the methylene sulfonamide from
part F (20.46 g, 51.5 mmol) in dimethylformamide (90
mL) was added potassium carbonate (21.3 g, 154.7
30 mmol), bis-(2-chloroethyl)benzyl amine (12.0 g, 51.5
mmol; and 18-Crown-6 (700 mg). The slurry was
stirred at sixty degrees Celsius. After twenty four
hours the reaction was concentrated in vacuo. The

-387-

WO 00/46221

residue was taken up in ethyl acetate, washed with water three times, saturated sodium chloride solution, dried over Na_2SO_4 , filtered, and concentrated in vacuo. The residue was recrystallized (methanol) to give the N-benzyl piperidine sulfonamide as a white solid (15 g, 52%).

5 Part H: Preparation of 4-[[4-[4-(trifluoromethoxy)-phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxylic acid, methyl ester

10 To a slurry of the N-benzyl piperidine sulfonamide from part G (4.17 g, 7.5 mmol) in methanol (50 mL) was added ammonium formate (1.46 g, 22.5 mmol). The system was purged with nitrogen for 10 minutes. The nitrogen stream was removed and

15 palladium on carbon (1.5 g of 10 weight % on activated carbon, 50% water) was added. The reaction was refluxed for thirty minutes, cooled, filtered through Celite under nitrogen, and concentrated in vacuo to give the unsubstituted piperidine

20 sulfonamide as a beige solid (3.15 g, 90%).
Part I: Preparation of methyl 1-(2-methoxyethyl)-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxylate

25 The unsubstituted piperidine sulfonamide from part H (3.0 g, 6.45 mmol) was dissolved in dry dimethylformamide (15 mL) and potassium carbonate (1.3 g, 9.7 mmol) and 2-bromoethyl methyl ether (908 μL , 9.7 mmol) were added. The reaction was stirred at thirty five degrees Celsius for sixteen hours and

30 then concentrated in vacuo. The residue was taken up in ethyl acetate and filtered through Celite. The filtrate washed with water two times, saturated

sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Chromatography (on silica, ethyl acetate/hexanes) provided the N-methoxyethyl piperidine sulfonamide as a white solid
5 (1.65 g, 50%).

Part J: Preparation of 1-(2-methoxyethyl)-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]-sulfonyl]-4-piperidinecarboxylic acid

In dry equipment under nitrogen, the N-methoxyethyl piperidine sulfonamide from part I (1.6 g, 3.0 mmol) was dissolved in dry tetrahydrofuran (25 mL) and potassium trimethylsilonate (1.3 g, 9.13 mmol) was added at ambient temperature. After twenty four hours, water (10 mL) was added and at 5 degrees
10 Celsius, 6 N HCl was added until pH=7. The slurry was filtered, washed with water and dried *in vacuo* to give the carboxylic acid as an off-white solid (1.39 g, 90%).

Part K: Preparation of 1-(2-methoxyethyl)-N-
20 [[(tetrahydro-2H-pyran-2-yl)oxy]-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]-sulfonyl]-4-piperidinecarboxylate

In dry equipment under nitrogen, the carboxylic acid from part J (1.36 g, 2.67 mmol) was dissolved in
25 dry dimethylformamide (9 mL) and the remaining reagents were added to the solution in the following order: N-hydroxybenzotriazole hydrate (0.43 g, 3.2 mmol), N-methylmorpholine (0.88 mL, 8.0 mmol), O-(tetrahydro-2H-pyran-2-yl)hydroxylamine (0.97 g, 8.0
30 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.72 g, 3.7 mmol). The reaction was stirred at forty degrees Celsius. After twenty hours, the reaction was concentrated in


-389-

vacuo. The residue was taken up in ethyl acetate, washed with water, 5% KHSO₄, saturated NaHCO₃, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Chromatography
5 (on silica, ethyl acetate/hexanes) provided the THP hydroxamate as a white solid (1.42 g, 90%).

Part L: Preparation of N-hydroxy-1-(2-methoxyethyl)-
4-[[4-[4-(trifluoromethoxy)phenoxy]-1-
piperidinyl]sulfonyl]-4-
10 piperidinecarboxamide, monohydrochloride

To a solution of the THP hydroxamate from part K (1.3 g, 2.13 mmol) in 1,4-dioxanes (2 mL) was added 4 N HCl dioxane solution (5.3 mL) and methanol (0.5
15 mL). After ten minutes, the reaction was diluted with diethyl ether, the solids filtered under nitrogen and dried *in vacuo* to give the title compound as a white solid (1.15 g, 96%). HRMS (ES+) M+ H⁺ calculated for C₂₁H₃₀F₃N₃O₇S₁: 526.1835, found
20 526.1805.

Example 16: Preparation of N-hydroxy-2-[[4-[4-(trifluoromethoxy)phenoxy]-1-
piperidinyl]sulfonyl]acetamide



25

Part A: Preparation of 1,1-dimethylethyl 4-hydroxy-1-
piperidinecarboxylate

In dry equipment under nitrogen, 4-
30 hydroxypiperidine (20.2 g, 0.2 mol) was dissolved in tetrahydrofuran (200 mL) and triethylamine (29 mL,

-390-

WO 00/46221

0.21 mol). A solution of di-t-butylidicarbonate (43.65 g, 0.2 mol) was added at such a rate that the temperature remained below thirty degrees Celsius. After stirring at ambient temperature for four hours,

5 the reaction was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water, 5% KHSO₄, saturated NaHCO₃, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo to give the BOC piperidine as a

10 white solid (37.7 g, 94%).
Part B: Preparation of 4-(methylsulfonyl)hydroxy-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester

15 To a solution of the BOC piperidine of part A (5.00 g, 24.84 mmol) in dichloromethane (50 mL) at zero degrees Celsius, was added triethylamine (3.81 mL, 27.32 mmol) followed by methane sulfonyl chloride (2.02 mL, 26.08 mmol). Once the addition was

20 complete the cooling bath was removed. After stirring for two hours the reaction mixture was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water two times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo to provide the mesylate as

25 an off-white solid (7.34 g, >100%).
Part C: Preparation of 4-[4-(trifluoromethoxy)-1,1-dimethylethyl ester]

30 In dry equipment under nitrogen, 4-trifluoromethoxyphenol (10.15 g, 57 mmol) was dissolved in dry dimethylformamide (125 mL) and at minus five degrees Celsius sodium hydride (2.74g, 68.4 mmol of the 60% oil dispersion) was added and

the ice bath was removed. After one hour at ambient temperature, the mesylate from part B (15.9 g, 57 mmol) was added and the reaction stirred at eighty degrees Celsius. After stirring at ambient
5 temperature for four hours, the reaction was concentrated in vacuo. The residue was taken up in diethyl ether, washed with water, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo to provide the substituted BOC-
10 piperidine as a beige solid (20.6 g, 100%).

Part D: Preparation of 4-[4-(trifluoromethoxy)-
phenoxy]piperidine

At fifteen degrees Celsius, 4 N HCl in dioxanes (125 mL) was slowly added to the substituted BOC-
15 piperidine from part C (20.6 g, 57 mmol) and stirred for ninety minutes. The reaction was concentrated in vacuo. The residue was dissolved in water (150 mL) and washed two times with ethyl acetate. The aqueous solution was cooled to five degrees Celsius and the
20 pH adjusted to eleven with 5 N sodium hydroxide solution and extracted with ethyl acetate. The ethyl acetate was dried over Na₂SO₄, filtered, and concentrated in vacuo to provide the substituted piperidine as a beige solid (11.9 g, 80%).

25 Part E: Preparation of 1-(methylsulfonyl)-4-[4-
(trifluoromethoxy)phenoxy]piperidine

The substituted piperidine from part D (11.5 g, 44.1 mmol) was dissolved in dichloromethane (125 mL) with triethylamine (12.3 mL, 88.1 mmol) and at zero
30 degrees Celsius a solution of methane sulfonyl chloride (5.1 mL, 66.1 mmol) in dichloromethane (20 mL) was added. After one hour at ambient temperature, the solvent was stripped in vacuo. The

residue was taken up in ethyl acetate, washed with water two times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo. The residue was recrystallized (ethyl acetate/hexanes) to give the sulfonamide as an off-white solid (10.77 g, 72%).

Part F: Preparation of methyl [[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxylate

10 In dry equipment under nitrogen, the sulfonamide from part E (10.77 g, 31.8 mmol) was dissolved in dry tetrahydrofuran (64 mL), chilled to minus seventy-five degrees Celsius, and a 1 M solution of lithium bis(trimethylsilyl)amide (80 mL) was added
15 maintaining the temperature below minus sixty five degrees. After thirty minutes at minus seventy-five degrees Celsius, a solution of methyl chloroformate (2.45 mL, 31.8 mmol) in dry tetrahydrofuran (32 mL) was added maintaining the temperature below minus
20 sixty degrees. After thirty minutes at minus seventy-five degrees Celsius, the reaction was quenched with saturated ammonium chloride solution (125 mL) and extracted with ethyl acetate. The combined extracts were washed with saturated ammonium
25 chloride solution, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo to give the methylene sulfonamide as an yellow oil (12.69 g, 100%).

Part G: Preparation of [[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]sulfonyl]acetic acid

30 In dry equipment under nitrogen, the methylene sulfonamide from part F (1.59 g, 4.0 mmol) was dissolved in dry tetrahydrofuran (10 mL) and

-393-

potassium trimethylsilonate (1.2 g, 8.5 mmol) was added at ambient temperature. After fifteen hours water (50 mL) was added and at 5 degrees Celsius 6N HCl was added until pH=1. The slurry was extracted with ethyl acetate and the combined extracts washed with water, dried over Na₂SO₄, filtered, and concentrated in vacuo. The product was recrystallized (hexanes/ethyl acetate) to give the carboxylic acid as a white solid (1.15 g, 75%).

10 Part H: Preparation of N-[(tetrahydro-2H-pyran-2-yl)oxy]-2-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]sulfonyl]acetamide


In dry equipment under nitrogen, the carboxylic acid from part G (1.0 g, 2.6 mmol) was dissolved in dry dimethylformamide (7 mL) and the remaining reagents were added to the solution in the following order: N-hydroxybenzotriazole hydrate (0.42 g, 3.1 mmol), N-methylmorpholine (0.86 mL, 7.8 mmol), O-(tetrahydro-2H-pyran-2-yl)hydroxylamine (0.9 g, 7.8 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.7 g, 3.7 mmol). After three hours at ambient temperature, the reaction was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water, 5% KHSO₄, saturated NaHCO₃, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo. The product was recrystallized (hexanes/ethyl acetate) to give the THP hydroxamate as a white solid (0.9 g, 71%).

30 Part I: Preparation of N-hydroxy-2-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]-sulfonyl]acetamide

-394-

To a solution of the THP hydroxamate from part H (0.84 g, 2.4 mmol) in 1,4-dioxane (4.5 mL) was added 4N HCl dioxane solution (4.5 mL) and methanol (64.5 mL). After thirty minutes at ambient temperature the
5 reaction was diluted with ethyl acetate and washed with water, dried over Na₂SO₄, filtered, and concentrated in vacuo. The product was recrystallized (ethyl acetate/hexanes) to give the title compound as a white solid (0.35 g, 51%). HRMS
10 (ES+) M+ H⁺ calculated for C₁₄H₁₇F₃N₂O₆S₁: 416.1103, found 416.1117.

Example 17: Preparation of tetrahydro-N-hydroxy-4-
[[4-[4-(trifluoromethoxy)phenoxy]-1-
15 piperidiny]sulfonyl]-2H-pyran-4-
carboxamide



20 Part A: Preparation of 1,1-dimethylethyl 4-hydroxy-1-
piperidinecarboxylate

In dry equipment under nitrogen, 4-hydroxypiperidine (20.2 g, 0.2 mol) was dissolved in tetrahydrofuran (200 mL) and triethylamine (29 mL,
25 0.21 mol). A solution of di-t-butyl dicarbonate (43.65 g, 0.2 mol) was added at such a rate that the temperature remained below thirty degrees Celsius. After stirring at ambient temperature for four hours, the reaction was concentrated in vacuo. The residue
30 was taken up in ethyl acetate, washed with water, 5%

KHSO₄, saturated NaHCO₃, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo to give the BOC piperidine as a white solid (37.7 g, 94%).

5 Part B: Preparation of 4-(methylsulfonyl)hydroxy-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester

To a solution of the BOC piperidine of part A (5.00 g, 24.84 mmol) in dichloromethane (50 mL) at
10 zero degrees Celsius, was added triethylamine (3.81 mL, 27.32 mmol) followed by methane sulfonyl chloride (2.02 mL, 26.08 mmol). Once the addition was complete the cooling bath was removed. After stirring for two hours the reaction mixture was
15 concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water two times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo to provide the mesylate as an off-white solid (7.34 g, >100%).

20 Part C: Preparation of 4-[4-(trifluoromethoxy)-phenoxy]-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester

In dry equipment under nitrogen, 4-trifluoromethoxyphenol (10.15 g, 57 mmol) was
25 dissolved in dry dimethylformamide (125 mL) and at minus five degrees Celsius sodium hydride (2.74g, 68.4 mmol of the 60% oil dispersion) was added and the ice bath was removed. After one hour at ambient temperature, the mesylate from part B (15.9 g, 57
30 mmol) was added and the reaction stirred at eighty degrees Celsius. After stirring at ambient temperature for four hours, the reaction was concentrated in vacuo. The residue was taken up in

diethyl ether, washed with water, saturated sodium chloride solution, dried over Na_2SO_4 , filtered, and concentrated in vacuo to provide the substituted BOC-piperidine as a beige solid (20.6 g, 100%).

5 Part D: Preparation of 4-[4-(trifluoromethoxy)-
phenoxy]piperidine

At fifteen degrees Celsius, 4 N HCl in dioxanes (125 mL) was slowly added to the substituted BOC-piperidine from part C (20.6 g, 57 mmol) and stirred
10 for ninety minutes. The reaction was concentrated in vacuo. The residue was dissolved in water (150 mL) and washed two times with ethyl acetate. The aqueous solution was cooled to five degrees Celsius and the pH adjusted to eleven with 5 N sodium hydroxide
15 solution and extracted with ethyl acetate. The ethyl acetate was dried over Na_2SO_4 , filtered, and concentrated in vacuo to provide the substituted piperidine as a beige solid (11.9 g, 80%).

20 Part E: Preparation of 1-(methanesulfonyl)-4-[4-
(trifluoromethoxy)phenoxy]piperidine

The substituted piperidine from part D (11.5 g, 44.1 mmol) was dissolved in dichloromethane (125 mL) with triethylamine (12.3 mL, 88.1 mmol) and at zero degrees Celsius a solution of methane sulfonyl
25 chloride (5.1 mL, 66.1 mmol) in dichloromethane (20 mL) was added. After one hour at ambient temperature, the solvent was stripped in vacuo. The residue was taken up in ethyl acetate, washed with water two times, saturated sodium chloride solution,
30 dried over Na_2SO_4 , filtered, and concentrated in vacuo. The residue was recrystallized (ethyl acetate/hexanes) to give the sulfonamide as an off-white solid (10.77 g, 72%).

Part F: Preparation of methyl [[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxylate

In dry equipment under nitrogen, the sulfonamide
5 from part E (10.77 g, 31.8 mmol) was dissolved in dry tetrahydrofuran (64 mL), chilled to minus seventy-five degrees Celsius, and a 1 M solution of lithium bis(trimethylsilyl)amide (80 mL) was added maintaining the temperature below minus sixty five
10 degrees. After thirty minutes at minus seventy-five degrees Celsius, a solution of methyl chloroformate (2.45 mL, 31.8 mmol) in dry tetrahydrofuran (32 mL) was added maintaining the temperature below minus sixty degrees. After thirty minutes at minus
15 seventy-five degrees Celsius, the reaction was quenched with saturated ammonium chloride solution (125 mL) and extracted with ethyl acetate. The combined extracts were washed with saturated ammonium chloride solution, saturated sodium chloride
20 solution, dried over Na₂SO₄, filtered, and concentrated in vacuo to give the methylene sulfonamide as a yellow oil (12.69 g, 100%).

Part G: Preparation of methyl tetrahydro-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]-
25 sulfonyl]-2H-pyran-4-carboxylate

To a solution of the methylene sulfonamide from part F (5.17 g, 13 mmol) in dimethylformamide (26 mL) was added potassium carbonate (7.2 g, 52.1 mmol), bis-(2-bromoethyl)ether (1.7 mL, 13 mmol) and 18-
30 Crown-6 (500mg). The slurry was stirred at sixty degrees Celsius. After sixteen hours potassium carbonate (2.0 g, 14 mmol) and bis-(2-bromoethyl)ether (0.2 mL, 1.6 mmol) were added and

the reaction stirred at sixty degrees Celsius. After a total of forty hours, the reaction was concentrated *in vacuo*. The residue was taken up in ethyl acetate, washed with water three times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The product was recrystallized (ethyl acetate/hexanes) to give the THP substituted sulfonamide as a white solid (1.85 g, 30%).

10 Part H: Preparation of tetrahydro-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]-sulfonyl]-2H-pyran-4-carboxylic acid

In dry equipment under nitrogen, the THP substituted sulfonamide from part G (1.9 g, 40.7 mmol) was dissolved in dry tetrahydrofuran (80.0 mL) and potassium trimethylsilonate (17.4 g, 122.0 mmol) was added at thirty five degrees Celsius. After two hours water (100 mL) was added and the solution concentrated *in vacuo*. The residue was taken up in water and extracted with ethyl acetate to remove unreacted starting material. The aqueous solution was treated with 6N HCl until pH=1. The slurry was extracted with ethyl acetate and the combined extracts washed with water, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The residue was heated in diethyl ether, the solid filtered and dried to give the carboxylic acid as a white solid (18.5 g, 100 %).

30 Part I: Preparation of tetrahydro-N-[(tetrahydro-2H-pyran-2-yl)oxy]-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]sulfonyl]-2H-pyran-4-carboxamide

In dry equipment under nitrogen, the carboxylic acid from part H (1.63 g, 3.6 mmol) was dissolved in dry dimethylformamide (7.5 mL) and the remaining reagents were added to the solution in the following order: N-hydroxybenzotriazole hydrate (0.97 g, 7.2 mmol), N-methylmorpholine (1.2 mL, 10.8 mmol), O-(tetrahydro-2H-pyran-2-yl)hydroxylamine (1.38 g, 7.2 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.38 g, 7.2 mmol). After one hour at forty five degrees Celsius, the reaction was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water, 5% KHSO₄, saturated NaHCO₃, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo. Chromatography (on silica, ethyl acetate/hexanes) provided the THP hydroxamate as a white foam (1.1 g, 55%).

Part J: Preparation of tetrahydro-N-hydroxy-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]-sulfonyl]-2H-pyran-4-carboxamide


To a solution of the THP hydroxamate from part I (1.0 g, 1.8 mmol) in 1,4-dioxane (5 mL) was added 4N HCl dioxane solution (5 mL) and methanol (5 mL). After thirty minutes at ambient temperature the reaction was diluted with ethyl acetate and washed with water, dried over Na₂SO₄, filtered, and concentrated in vacuo. The product was recrystallized (acetone/hexanes) to give the title compound as a white solid (0.58 g, 68%). HRMS (ES+) M+ H⁺ calculated for C₁₈H₂₃N₂O₇S₁F₃ : 469.1256, found 469.1287.

Example 18: Preparation of N-hydroxy-1-

-400-

(phenylmethyl)-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-sulfonyl]-4-piperidinecarboxamide
monohydrochloride

5



Part A: Preparation of 1,1-dimethylethyl 4-hydroxy-1-
piperidinecarboxylate

10 In dry equipment under nitrogen, 4-hydroxypiperidine (20.2 g, 0.2 mol) was dissolved in tetrahydrofuran (200 mL) and triethylamine (29 mL, 0.21 mol). A solution of di-t-butyl dicarbonate (43.65 g, 0.2 mol) was added at such a rate that the
 15 temperature remained below thirty degrees Celsius. After stirring at ambient temperature for four hours, the reaction was concentrated *in vacuo*. The residue was taken up in ethyl acetate, washed with water, 5% KHSO₄, saturated NaHCO₃, saturated sodium chloride
 20 solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo* to give the BOC piperidine as a white solid (37.7 g, 94%).

Part B: Preparation of 1,1-dimethylethyl 4-[4-(trifluoromethyl)phenoxy]-1-

25 piperidinecarboxylate

To a solution of the BOC piperidine from part A (6.03 g, 30 mmol) in dimethylformamide (60 mL) was added cesium carbonate (9.77 g, 30 mmol) and 4-

-401-

WO 00/46221

fluorobenzotrifluoride (3.8 mL, 30 mmol). The slurry was stirred at ninety degrees Celsius. After nineteen hours cesium carbonate (3.26g, 10 mmol) and 4-fluorobenzotrifluoride (0.95ml mL, 10 mmol) were added and the reaction continued at ninety degrees Celsius. After a total of forty six hours, the reaction was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water three times, saturated sodium chloride solution, dried over Na_2SO_4 , filtered, and concentrated in vacuo. Chromatography (on silica, ethyl acetate/hexanes) provided the substituted BOC piperidine as a white solid (6.0 g, 58%).

Part C: Preparation of 4-[4-(trifluoromethyl)-phenoxy]piperidine

To a slurry of the substituted BOC piperidine from part B (5.95 g, 17.2 mmol) in 1,4-dioxane (10 mL) was added 4N HCl dioxane solution (17 mL). After one hour at ambient temperature, the reaction was concentrated in vacuo. The residue was slurried in diethyl ether and vacuum filtration of the resulting precipitate provided the hydrochloride salt as a white solid (4.6 g, 100%).

Part D: Preparation of 1-(methylsulfonyl)-4-[4-(trifluoromethyl)phenoxy]piperidine

To a solution of the hydrochloride salt from part C (4.6 g, 16.9 mmol) and triethylamine (5.9 mL, 42.4 mmol) in dichloromethane (45 mL) at zero degrees Celsius was added a solution of methane sulfonyl chloride (1.97 mL, 25.4 mmol) in dichloromethane (10 mL). After one hour at ambient temperature, the solvent was stripped in vacuo. The residue was taken

up in ethyl acetate, washed with water two times, saturated sodium chloride solution, dried over Na_2SO_4 , filtered, and concentrated *in vacuo*. The residue was slurried in diethyl ether and vacuum filtration of
5 the resulting precipitate provided the sulfonamide as an off-white solid (5.25 g, 96%).

Part E: Preparation of methyl [[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-
sulfonyl]acetate

10 In dry equipment under nitrogen, the sulfonamide from part D (4.2 g, 13 mmol) was dissolved in dry tetrahydrofuran (26 mL), chilled to minus seventy-five degrees Celsius, and a 1 M solution of lithium bis(trimethylsilyl)amide (26 mL) was added
15 maintaining the temperature below minus sixty five degrees. After thirty minutes at minus seventy-five degrees Celsius, a solution of methyl chloroformate (1.0 mL, 13 mmol) in dry tetrahydrofuran (13 mL) was added maintaining the temperature below minus sixty
20 degrees. After thirty minutes at minus seventy-five degrees Celsius, the reaction was quenched with saturated ammonium chloride solution (100 mL) and extracted with ethyl acetate. The combined extracts were washed with saturated ammonium chloride
25 solution, saturated sodium chloride solution, dried over Na_2SO_4 , filtered, and concentrated *in vacuo* to give the methylene sulfonamide as an yellow oil (4.95 g, 100%).

Part F: Preparation of methyl 1-(phenylmethyl)-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-
30 sulfonyl]-4-piperidinecarboxylate

To a solution of the methylene sulfonamide from part E (1.14 g, 3 mmol) in dimethylformamide (6 mL)

were added potassium carbonate (1.24 g, 9 mmol), bis-(2-chloroethyl)benzyl amine (0.7 g, 3 mmol, and 18-Crown-6 (500mg). The slurry was stirred at sixty degrees Celsius. After sixteen hours the reaction
5 was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water three times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo. Chromatography (on silica, ethyl acetate/hexanes) provided the
10 piperidine sulfonamide as a white solid (950 mg, 59%).

Part G: Preparation of 1-(phenylmethyl)-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-sulfonyl]-4-piperidinecarboxylic acid

15 In dry equipment under nitrogen, the piperidine sulfonamide from part F (7.2 g, 13.3 mmol) was dissolved in dry tetrahydrofuran (26 mL) and potassium trimethylsilonate (5.7 g, 40 mmol) was added and stirred at forty degrees Celsius. After
20 twenty four hours water (50 mL) was added and at 5 degrees Celsius, 6N HCl was added until pH=1. The slurry was filtered, washed with water and dried in vacuo to give the carboxylic acid as a white solid (6.5 g, 93%).

25 Part H: Preparation of 1-(phenylmethyl)-N-[[4-[[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-sulfonyl]-4-piperidinecarboxamide

In dry equipment under nitrogen, the carboxylic
30 acid from part G (6.46 g, 12.2 mmol) was dissolved in dry dimethylformamide (25 mL) and the remaining reagents were added to the solution in the following order: N-hydroxybenzotriazole hydrate (2.0 g, 14.7

mmol), N-methylmorpholine (4.0 mL, 36.8 mmol), O-(tetrahydro-2H-pyran-2-yl)hydroxylamine (4.3 g, 36.8 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (3.3 g, 17.2 mmol).

5 The reaction was stirred at forty five degrees Celsius. After three hours, the reaction was concentrated *in vacuo*. The residue was taken up in ethyl acetate, washed with water, 5% KHSO₄, saturated NaHCO₃, saturated sodium chloride solution, dried over
10 Na₂SO₄, filtered, and concentrated *in vacuo*. Chromatography (on silica, ethyl acetate/hexanes) provided the THP hydroxamate as a white solid (6.1 g, 80%).


Part I: Preparation of N-hydroxy-1-(phenylmethyl)-4-

15 [[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide
monohydrochloride

To a solution of the THP hydroxamate from part H (6.0 g, 9.6 mmol) in 1,4-dioxane (2.4 mL) was added 4
20 N HCl dioxane solution (2.4 mL) and methanol (2.4 mL). After thirty minutes at ambient temperature the reaction was poured into diethyl ether (100 mL). The slurry was filtered under nitrogen, washed with diethyl ether, and dried *in vacuo* to give the title
25 compound as an off white solid (5.4 g, 97%). HRMS (ES+) M+ H⁺ calculated for C₂₅H₃₀F₃N₃O₅S₁: 542.1937, found 542.1938.

Example 19: Preparation of N-hydroxy-1-(2-

30 pyridinylmethyl)-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]sulfonyl]-4-
piperidinecarboxamide, monohydrochloride



Part A: Preparation of 1,1-fimethylethyl 4-hydroxy-1-
piperidinecarboxylate

In dry equipment under nitrogen, 4-hydroxypiperidine (20.2 g, 0.2 mol) was dissolved in tetrahydrofuran (200 mL) and triethylamine (29 mL, 0.21 mol). A solution of di-*t*-butyldicarbonate (43.65 g, 0.2 mol) was added at such a rate that the temperature remained below thirty degrees Celsius. After stirring at ambient temperature for four hours, the reaction was concentrated *in vacuo*. The residue was taken up in ethyl acetate, washed with water, 5% KHSO_4 , saturated NaHCO_3 , saturated sodium chloride solution, dried over Na_2SO_4 , filtered, and concentrated *in vacuo* to give the BOC piperidine as a white solid (37.7 g, 94%).

Part B: Preparation of 4-(methylsulfonyl)hydroxy-1-
20 piperidinecarboxylic acid, 1,1-dimethylethyl
ester

To a solution of the BOC piperidine of part A (5.00 g, 24.84 mmol) in dichloromethane (50 mL) at zero degrees Celsius, was added triethylamine (3.81 mL, 27.32 mmol) followed by methane sulfonyl chloride (2.02 mL, 26.08 mmol). Once the addition was complete the cooling bath was removed. After stirring for two hours the reaction mixture was

concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water two times, saturated sodium chloride solution, dried over Na_2SO_4 , filtered, and concentrated in vacuo to provide the mesylate as
5 an off-white solid (7.34 g, >100%).

Part C: Preparation of 4-[4-(trifluoromethoxy)-
phenoxy]-1-piperidinecarboxylic acid, 1,1-
dimethylethyl ester

In dry equipment under nitrogen, 4-
10 trifluoromethoxyphenol (10.15 g, 57 mmol) was
dissolved in dry dimethylformamide (125 mL) and at
minus five degrees Celsius sodium hydride (2.74g,
68.4 mmol of the 60% oil dispersion) was added and
the ice bath was removed. After one hour at ambient
15 temperature, the mesylate from part B (15.9 g, 57
mmol) was added and the reaction stirred at eighty
degrees Celsius. After stirring at ambient
temperature for four hours, the reaction was
concentrated in vacuo. The residue was taken up in
20 diethyl ether, washed with water, saturated sodium
chloride solution, dried over Na_2SO_4 , filtered, and
concentrated in vacuo to provide the substituted BOC-
piperidine as a beige solid (20.6 g, 100%).

Part D: Preparation of 4-[4-(trifluoromethoxy)-
25 phenoxy]piperidine

At fifteen degrees Celsius, 4 N HCl in dioxanes
(125 mL) was slowly added to the substituted BOC-
piperidine from part C (20.6 g, 57 mmol) and stirred
for ninety minutes. The reaction was concentrated in
30 vacuo. The residue was dissolved in water (150 mL)
and washed two times with ethyl acetate. The aqueous
solution was cooled to five degrees Celsius and the
pH adjusted to eleven with 5 N sodium hydroxide

solution and extracted with ethyl acetate. The ethyl acetate was dried over Na_2SO_4 , filtered, and concentrated *in vacuo* to provide the substituted piperidine as a beige solid (11.9 g, 80%).

5 Part E: Preparation of 1-(methylsulfonyl)-4-[4-(trifluoromethoxy)phenoxy]piperidine

The substituted piperidine from part D (11.5 g, 44.1 mmol) was dissolved in dichloromethane (125 mL) with triethylamine (12.3 mL, 88.1 mmol) and at zero
10 degrees Celsius a solution of methane sulfonyl chloride (5.1 mL, 66.1 mmol) in dichloromethane (20 mL) was added. After one hour at ambient temperature, the solvent was stripped *in vacuo*. The residue was taken up in ethyl acetate, washed with
15 water two times, saturated sodium chloride solution, dried over Na_2SO_4 , filtered, and concentrated *in vacuo*. The residue was recrystallized (ethyl acetate/hexanes) to give the sulfonamide as an off-white solid (10.77 g, 72%).

20 Part F: Preparation of methyl [[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]-sulfonyl]-4-piperidinecarboxylate

In dry equipment under nitrogen, the sulfonamide from part E (10.77 g, 31.8 mmol) was dissolved in dry
25 tetrahydrofuran (64 mL), chilled to minus seventy-five degrees Celsius, and a 1M solution of lithium bis(trimethylsilyl)amide (80 mL) was added maintaining the temperature below minus sixty five degrees. After thirty minutes at minus seventy-five
30 degrees Celsius, a solution of methyl chloroformate (2.45 mL, 31.8 mmol) in dry tetrahydrofuran (32 mL) was added maintaining the temperature below minus sixty degrees. After thirty minutes at minus

seventy-five degrees Celsius, the reaction was quenched with saturated ammonium chloride solution (125 mL) and extracted with ethyl acetate. The combined extracts were washed with saturated ammonium chloride solution, saturated sodium chloride solution, dried over Na_2SO_4 , filtered, and concentrated *in vacuo* to give the methylene sulfonamide as a yellow oil (12.69 g, 100%).

Part G: Preparation of methyl 1-(phenylmethyl)-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxylate

To a solution of the methylene sulfonamide from part F (20.46 g, 51.5 mmol) in dimethylformamide (90 mL) was added potassium carbonate (21.3 g, 154.7 mmol), bis-(2-chloroethyl)benzyl amine (12.0 g, 51.5 mmol; and 18-Crown-6 (700mg). The slurry was stirred at sixty degrees Celsius. After twenty four hours the reaction was concentrated *in vacuo*. The residue was taken up in ethyl acetate, washed with water three times, saturated sodium chloride solution, dried over Na_2SO_4 , filtered, and concentrated *in vacuo*. The residue was recrystallized (methanol) to give the N-benzyl piperidine sulfonamide as a white solid (15 g, 52%).

Part H: Preparation of 4-[[4-[4-(trifluoromethoxy)-phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxylic acid, methyl ester

To a slurry of the N-benzyl piperidine sulfonamide from part G (4.17 g, 7.5 mmol) in methanol (50 mL) was added ammonium formate (1.46 g, 22.5 mmol). The system was purged with nitrogen for 10 minutes. The nitrogen stream was removed and

palladium on carbon (1.5 g of 10 weight % on activated carbon, 50%water) was added. The reaction was refluxed for thirty minutes, cooled, filtered through Celite under nitrogen, and concentrated in vacuo to give the unsubstituted piperidine sulfonamide as a beige solid (3.15 g, 90%).

Part I: Preparation of methyl 1-(2-pyridinylmethyl)-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxylate

10 The unsubstituted piperidine sulfonamide from part H (7.9 g, 16.95 mmol) was dissolved in dry dimethylformamide (35 mL) and potassium carbonate (7.05 g, 51.1 mmol) and 2-picolyl chloride hydrochloride (4.21 g, 25.67 mmol) were added. The reaction was stirred at thirty five degrees Celsius for sixteen hours and then concentrated in vacuo. The residue was taken up in ethyl acetate and filtered through Celite. The filtrate washed with water two times, saturated sodium chloride solution, 15 dried over Na₂SO₄, filtered, and concentrated in vacuo. Chromatography (on silica, ethyl acetate/hexanes) provided the N-picolyl piperidine sulfonamide as a white solid (6.8 g, 72%).

Part J: Preparation of 1-(2-pyridinylmethyl)-N-25 [(tetrahydro-2H-pyran-2-yl)oxy]-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide

In dry equipment under nitrogen, the N-picolyl piperidine sulfonamide from part I (8.5g, 15.3mmol) was dissolved in dry tetrahydrofuran (30 mL) and potassium trimethylsilonate (6.52 g, 45.8 mmol) was added at ambient temperature. After eighteen hours water (50 mL) was added and at 5 degrees Celsius, 6 N


-410-

HCl was added until pH=7. The slurry was filtered, washed with water and dried in vacuo to give the carboxylic acid as a off-white solid (6.9 g, 84%). In dry equipment under nitrogen, the carboxylic acid (6.8 g, 12.5 mmol) was dissolved in dry dimethylformamide (25 mL) and the remaining reagents were added to the solution in the following order: N-hydroxybenzotriazole hydrate (2.03 g, 15.0 mmol), N-methylmorpholine (4.1 mL, 37.6 mmol), O- (tetrahydro-2H-pyran-2-yl)hydroxylamine (4.4 g, 37.6 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (3.36 g, 17.5 mmol). The reaction was stirred at forty degrees Celsius. After twelve hours, the reaction was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water, 5% KHSO₄, saturated NaHCO₃, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo. Chromatography (on silica, ethyl acetate/hexanes) provided the THP hydroxamate as a light yellow foam (7.55 g, 94%).

Part K: Preparation of N-hydroxy-1-(2-pyridinylmethyl)-4-[[4-[4-(trifluoromethoxy)-phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide, monohydrochloride

To a solution of the THP hydroxamate from part J (7.4 g, 11.5 mmol) in 1,4-dioxane (2 mL) was added 4N HCl dioxane solution (29 mL, 115.2 mmol) and methanol (2.9 mL). After fifteen minutes, the reaction was diluted with diethyl ether, the solids filtered under nitrogen and dried in vacuo to give the title compound as a white solid (1.15 g, 96%). HRMS (ES+) M+ H⁺ calculated for C₂₄H₂₉F₃N₄O₆S₁: 559.1838, found 559.1864.

Example 20: Preparation of N-hydroxy-1-(2-
pyrimidinyl)-4-[[4-[4-(trifluoromethyl)-
5 phenoxy]-1-piperidinyl]sulfonyl]-4-
piperidinecarboxamide, monohydrochloride



Part A: Preparation of 1,1-dimethylethyl 4-hydroxy-1-
10 piperidinecarboxylate

In dry equipment under nitrogen, 4-
hydroxypiperidine (20.2 g, 0.2 mol) was dissolved in
tetrahydrofuran (200 mL) and triethylamine (29 mL,
0.21 mol). A solution of di-t-butyldicarbonate
15 (43.65 g, 0.2 mol) was added at such a rate that the
temperature remained below thirty degrees Celsius.
After stirring at ambient temperature for four hours,
the reaction was concentrated *in vacuo*. The residue
was taken up in ethyl acetate, washed with water, 5%
20 KHSO₄, saturated NaHCO₃, saturated sodium chloride
solution, dried over Na₂SO₄, filtered, and
concentrated *in vacuo* to give the BOC piperidine as a
white solid (37.7 g, 94%).

Part B: Preparation of 1,1-dimethylethyl 4-[4-
25 (trifluoromethyl)phenoxy]-1-
piperidinecarboxylate

To a solution of the BOC piperidine from part A
(6.03 g, 30 mmol) in dimethylformamide (60 mL) was

added cesium carbonate (9.77 g, 30 mmol) and 4-fluorobenzotrifluoride (3.8 mL, 30 mmol). The slurry was stirred at ninety degrees Celsius. After nineteen hours cesium carbonate (3.26g, 10 mmol) and
5 4-fluorobenzotrifluoride (0.95ml mL, 10 mmol) were added and the reaction continued at ninety degrees Celsius. After a total of forty six hours, the reaction was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water three
10 times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo. Chromatography (on silica, ethyl acetate/hexanes) provided the substituted BOC piperidine as a white solid (6.0 g, 58%).

15 Part C: Preparation of 4-[4-(trifluoromethyl)-
phenoxy]piperidine

To a slurry of the substituted BOC piperidine from part B (5.95 g, 17.2 mmol) in 1,4-dioxane (10 mL) was added 4N HCl dioxane solution (17 mL). After
20 one hour at ambient temperature the reaction was concentrated in vacuo. The residue was slurried in diethyl ether and vacuum filtration of the resulting precipitate provided the hydrochloride salt as a white solid (4.6 g, 100%).

25 Part D: Preparation of 1-(methylsulfonyl)-4-[4-
(trifluoromethyl)phenoxy]piperidine

To a solution of the hydrochloride salt from part C (4.6 g, 16.9 mmol) and triethylamine (5.9 mL, 42.4 mmol) in dichloromethane (45 mL) at zero degrees
30 Celsius was added a solution of methane sulfonyl chloride (1.97 mL, 25.4 mmol) in dichloromethane (10 mL). After one hour at ambient temperature, the solvent was stripped in vacuo. The residue was taken

up in ethyl acetate, washed with water two times, saturated sodium chloride solution, dried over Na_2SO_4 , filtered, and concentrated in vacuo. The residue was slurried in diethyl ether and vacuum filtration of the resulting precipitate provided the sulfonamide as an off-white solid (5.25 g, 96%).

Part E: Preparation of methyl [[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-sulfonyllacetate

10 In dry equipment under nitrogen, the sulfonamide from part D (4.2 g, 13 mmol) was dissolved in dry tetrahydrofuran (26 mL), chilled to minus seventy-five degrees Celsius, and a 1 M solution of lithium bis(trimethylsilyl)amide (26 mL) was added
15 maintaining the temperature below minus sixty five degrees. After thirty minutes at minus seventy-five degrees Celsius, a solution of methyl chloroformate (1.0mL, 13 mmol) in dry tetrahydrofuran (13 mL) was added maintaining the temperature below minus sixty
20 degrees. After thirty minutes at minus seventy-five degrees Celsius, the reaction was quenched with saturated ammonium chloride solution (100 mL) and extracted with ethyl acetate. The combined extracts were washed with saturated ammonium chloride
25 solution, saturated sodium chloride solution, dried over Na_2SO_4 , filtered, and concentrated in vacuo to give the methylene sulfonamide as an yellow oil (4.95 g, 100%).

Part F: Preparation of methyl 1-(phenylmethyl)-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-sulfonyl]-4-piperidinecarboxylate
30

To a solution of the methylene sulfonamide from part E (1.14 g, 3 mmol) in dimethylformamide (6 mL)

-414-

was added potassium carbonate (1.24 g, 9 mmol), bis-(2-chloroethyl)benzyl amine (0.7 g, 3 mmol); and 18-Crown-6 (500mg). The slurry was stirred at sixty degrees Celsius. After sixteen hours the reaction
5 was concentrated *in vacuo*. The residue was taken up in ethyl acetate, washed with water three times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Chromatography (on silica, ethyl acetate/hexanes) provided the
10 piperidine sulfonamide as a white solid (950 mg, 59%).

Part G: Preparation of methyl 1-(2-pyrimidinyl)-4-
[[4-[4-(trifluoromethyl)phenoxy]-1-
piperidinylsulfonyl]-4-piperidinecarboxylate

15

To a slurry of the piperidine sulfonamide from part F (6.3 g, 12.0 mmol) in methanol (25 mL) was added ammonium formate (2.2 g, 34.5 mmol). The system was purged with nitrogen for 10 minutes. The
20 nitrogen stream was removed and palladium on carbon (1.2 g of 10 weight % on activated carbon, 50%water) was added. The reaction was refluxed for forty five minutes, cooled, filtered through Celite under nitrogen, and concentrated *in vacuo*. The residue
25 (6.21 g, 13.8 mmol) was dissolved in dry dimethylformamide (28 mL) and potassium carbonate (5.7 g, 41.4 mmol) and 2-chloropyrimidine (3.16 g, 27.6 mmol) were added. The reaction was stirred at eighty five degrees Celsius for one hour and then
30 concentrated *in vacuo*. The residue was taken up in ethyl acetate and filtered through Celite. The filtrate washed with water two times, saturated sodium chloride solution, dried over Na₂SO₄, filtered,

-415-

and concentrated *in vacuo*. The product was recrystallized (methanol) to give the N- pyrimidinyl piperidine sulfonamide as a white solid (4.94 g, 68%).

- 5 Part H: Preparation of 1-(2-pyrimidinyl)-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-sulfonyl]-4-piperidinecarboxylic acid

In dry equipment under nitrogen, the N-
10 pyrimidinyl piperidine sulfonamide from part G (4.9 g, 9.28 mmol) was dissolved in dry tetrahydrofuran (35 mL) and potassium trimethylsilonate (3.97 g, 27.8 mmol) was added at thirty five degrees Celsius. After sixteen hours water (50 mL) was added and at 5
15 degrees Celsius, 6 N HCl was added until pH=2. The aqueous slurry was extracted with ethyl acetate. The ethyl acetate solution was washed with water, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The solids were
20 slurried in acetonitrile, filtered under nitrogen, washed with hexanes and dried *in vacuo* to give the carboxylic acid as a white solid (4.7 g, 98%).

- Part I: Preparation of 1-(2-pyrimidinyl)-N-
[(tetrahydro-2H-pyran-2-yl)oxy]-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-
25 sulfonyl]-4-piperidinecarboxamide

In dry equipment under nitrogen, the carboxylic acid from part H (4.6 g, 8.95 mmol) was dissolved in dry dimethylformamide (18 mL) and the remaining
30 reagents were added to the solution in the following order: N-hydroxybenzotriazole hydrate (1.45 g, 10.7mmol), N-methylmorpholine (2.95 mL, 26.8 mmol), O- (tetrahydro-2H-pyran-2-yl)hydroxylamine (3.14 g,

26.8 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (2.4 g, 12.5 mmol). The reaction was stirred at thirty degrees Celsius. After sixteen hours, the reaction was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water, 5% KHSO₄, saturated NaHCO₃, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo. The product was recrystallized (methanol) to give the THP hydroxamate as a white solid (5.0 g, 91%).

Part J: Preparation of N-hydroxy-1-(2-pyrimidinyl)-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide, monohydrochloride


15

To a solution of the THP hydroxamate from part I (4.97 g, 8.1 mmol) in 1,4-dioxane (3 mL) was added 4N HCl dioxane solution (20 mL, 81.1 mmol) and methanol (2 mL). After thirty minutes at ambient temperature the reaction was poured into 100 mL acetonitrile. The slurry was filtered under nitrogen, washed with acetonitrile and dried in vacuo to give the title compound as a white solid (3.3 g, 72%). HRMS (ES+) M+ H⁺ calculated for C₂₂H₂₆F₃N₅O₅S₁: 530.1685, found 530.1696.

25

Example 21: Preparation of N-hydroxy-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-sulfonyl]-1-[4-(trifluoromethyl)-2-pyrimidinyl]-4-piperidinecarboxamide, monohydrochloride

30



Part A: Preparation of 1,1-dimethylethyl 4-hydroxy-1-
piperidinecarboxylate

5 In dry equipment under nitrogen, 4-hydroxypiperidine (20.2 g, 0.2 mol) was dissolved in tetrahydrofuran (200 mL) and triethylamine (29 mL, 0.21 mol). A solution of di-*t*-butyldicarbonate (43.65 g, 0.2 mol) was added at such a rate that the
10 temperature remained below thirty degrees Celsius. After stirring at ambient temperature for four hours, the reaction was concentrated *in vacuo*. The residue was taken up in ethyl acetate, washed with water, 5% KHSO₄, saturated NaHCO₃, saturated sodium chloride
15 solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo* to give the BOC piperidine as a white solid (37.7 g, 94%).

Part B: Preparation of 1,1-dimethylethyl 4-[4-(trifluoromethyl)phenoxy]-1-

20 piperidinecarboxylate

To a solution of the BOC piperidine from part A (6.03 g, 30 mmol) in dimethylformamide (60 mL) was added cesium carbonate (9.77 g, 30 mmol) and 4-fluorobenzotrifluoride (3.8 mL, 30 mmol). The slurry
25 was stirred at ninety degrees Celsius. After nineteen hours cesium carbonate (3.26g, 10 mmol) and 4-fluorobenzotrifluoride (0.95mL, 10 mmol) were added and the reaction continued at ninety degrees

Celsius. After a total of forty six hours, the reaction was concentrated *in vacuo*. The residue was taken up in ethyl acetate, washed with water three times, saturated sodium chloride solution, dried over
5 Na_2SO_4 , filtered, and concentrated *in vacuo*.

Chromatography (on silica, ethyl acetate/hexanes) provided the substituted BOC piperidine as a white solid (6.0 g, 58%).

Part C: Preparation of 4-[4-(trifluoromethyl)-
10 phenoxy]piperidine

To a slurry of the substituted BOC piperidine from part B (5.95 g, 17.2 mmol) in 1,4-dioxane (10 mL) was added 4 N HCl dioxane solution (17 mL). After one hour at ambient temperature the reaction
15 was concentrated *in vacuo*. The residue was slurried in diethyl ether and vacuum filtration of the resulting precipitate provided the hydrochloride salt as a white solid (4.6 g, 100%).

Part D: Preparation of 1-(methanesulfonyl)-4-[4-
20 (trifluoromethyl)phenoxy]piperidine

To a solution of the hydrochloride salt from part C (4.6 g, 16.9 mmol) and triethylamine (5.9 mL, 42.4 mmol) in dichloromethane (45 mL) at zero degrees Celsius was added a solution of methane sulfonyl
25 chloride (1.97 mL, 25.4 mmol) in dichloromethane (10 mL). After one hour at ambient temperature, the solvent was stripped *in vacuo*. The residue was taken up in ethyl acetate, washed with water two times, saturated sodium chloride solution, dried over Na_2SO_4 ,
30 filtered, and concentrated *in vacuo*. The residue was slurried in diethyl ether and vacuum filtration of the resulting precipitate provided the sulfonamide as an off-white solid (5.25 g, 96%).

Part E: Preparation of methyl [[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-sulfonyl]acetate

In dry equipment under nitrogen, the sulfonamide
5 from part D (4.2 g, 13 mmol) was dissolved in dry
tetrahydrofuran (26 mL), chilled to minus seventy-
five degrees Celsius, and a 1 M solution of lithium
bis(trimethylsilyl)amide (26 mL) was added
maintaining the temperature below minus sixty five
10 degrees. After thirty minutes at minus seventy-five
degrees Celsius, a solution of methyl chloroformate
(1.0 mL, 13 mmol) in dry tetrahydrofuran (13 mL) was
added maintaining the temperature below minus sixty
degrees. After thirty minutes at minus seventy-five
15 degrees Celsius, the reaction was quenched with
saturated ammonium chloride solution (100 mL) and
extracted with ethyl acetate. The combined extracts
were washed with saturated ammonium chloride
solution, saturated sodium chloride solution, dried
20 over Na₂SO₄, filtered, and concentrated in vacuo to
give the methylene sulfonamide as a yellow oil
(4.95 g, 100%).

Part F: Preparation of methyl 1-(phenylmethyl)-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-sulfonyl]-4-piperidinecarboxylate

To a solution of the methylene sulfonamide from
part E (1.14 g, 3 mmol) in dimethylformamide (6 mL)
was added potassium carbonate (1.24 g, 9 mmol), bis-
(2-chloroethyl)benzyl amine (0.7 g, 3 mmol; and 18-
30 Crown-6 (500 mg). The slurry was stirred at sixty
degrees Celsius. After sixteen hours the reaction
was concentrated in vacuo. The residue was taken up
in ethyl acetate, washed with water three times,

saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Chromatography (on silica, ethyl acetate/hexanes) provided the piperidine sulfonamide as a white solid (950 mg, 59%).

Part G: Preparation of methyl 4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-sulfonyl]-1-[4-(trifluoromethyl)-2-pyrimidinyl]-4-piperidinecarboxylate

10

To a slurry of the piperidine sulfonamide from part F (6.3 g, 12.0 mmol) in methanol (25 mL) was added ammonium formate (2.2 g, 34.5 mmol). The system was purged with nitrogen for 10 minutes. The nitrogen stream was removed and palladium on carbon (1.2 g of 10 weight % on activated carbon, 50%water) was added. The reaction was refluxed for forty five minutes, cooled, filtered through Celite under nitrogen, and concentrated *in vacuo*. The residue (4.5 g, 10 mmol) was dissolved in dry dimethylformamide (20 mL) and potassium carbonate (4.14 g, 30 mmol) and 2-chloro-4-trifluoromethylpyrimidine (2.4 mL, 20 mmol) were added. The reaction was stirred at seventy degrees Celsius for three hour and then concentrated *in vacuo*. The residue was taken up in ethyl acetate and filtered through Celite. The filtrate washed with water two times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The product was recrystallized (methanol) to give the N-pyrimidinyl piperidine sulfonamide as a white solid (4.51 g, 76%).

Part H: Preparation of 4-[[4-[4-(trifluoromethyl)-

phenoxy]-1-piperidinyl)sulfonyl]-1-[4-(trifluoromethyl)-2-pyrimidinyl]-4-
piperidinecarboxylate

In dry equipment under nitrogen, the N-
5 pyrimidinyl piperidine sulfonamide from part G (4.9 g, 9.28 mmol) was dissolved in dry tetrahydrofuran (35 mL) and potassium trimethylsilonate (3.2 g, 22.45 mmol) was added at forty five degrees Celsius. After four hours water (100 mL) was added and at 5 degrees
10 Celsius, 6 N HCl was added until pH=2. The slurry was filtered under nitrogen, the solids washed with hexanes and dried *in vacuo* to give the carboxylic acid as a white solid (4.13 g, 95%).

Part I: Preparation of N-[(tetrahydro-2H-pyran-2-
15 yl)oxy]-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl)sulfonyl]-1-[4-(trifluoromethyl)-
2-pyrimidinyl]-4-piperidinecarboxamide


In dry equipment under nitrogen, the carboxylic acid from part H (4.0 g, 6.87 mmol) was dissolved in
20 dry dimethylformamide (20 mL) and the remaining reagents were added to the solution in the following order: N-hydroxybenzotriazole hydrate (1.1 g, 8.25 mmol), N-methylmorpholine (2.2 mL, 20.6 mmol), O-(tetrahydro-2H-pyran-2-yl)hydroxylamine (2.4 g, 20.6
25 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.84 g, 9.62 mmol). The reaction was stirred at forty five degrees Celsius. After sixteen hours, the reaction was concentrated *in vacuo*. The residue was taken up in
30 ethyl acetate, washed with water, 5% KHSO₄, saturated NaHCO₃, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The

product was recrystallized (methanol) to give the THP hydroxamate as a white solid (4.06 g, 87%).

Part J: Preparation of N-hydroxy-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-sulfonyl]-1-[4-(trifluoromethyl)-2-pyrimidinyl]-4-piperidinecarboxamide, monohydrochloride

To a solution of the THP hydroxamate from part I (5.0 g, 5.87 mmol) in 1,4-dioxane (3 mL) was added 4N HCl dioxane solution (14.7 mL, 58.7 mmol) and methanol (1.5 mL). After thirty minutes at ambient temperature the reaction was poured into 100 mL acetonitrile. The slurry was filtered under nitrogen, washed with acetonitrile and dried *in vacuo* to give the title compound as a white solid (2.95 g, 80%). HRMS (ES+) $M + H^+$ calculated for $C_{23}H_{25}F_6N_5O_5S_1$: 598.1559, found 598.1531.

Example 22: Preparation of 1-(5-ethyl-2-pyrimidinyl)-N-hydroxy-4-[[4-[4-(trifluoromethyl)-phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide, monohydrochloride



Part A: Preparation of 1,1-dimethylethyl 4-hydroxy-1-

piperidinecarboxylate

In dry equipment under nitrogen, 4-hydroxypiperidine (20.2 g, 0.2 mol) was dissolved in tetrahydrofuran (200 mL) and triethylamine (29 mL, 0.21 mol). A solution of di-t-butylidicarbonate (43.65 g, 0.2 mol) was added at such a rate that the temperature remained below thirty degrees Celsius. After stirring at ambient temperature for four hours, the reaction was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water, 5% KHSO₄, saturated NaHCO₃, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo to give the BOC piperidine as a white solid (37.7 g, 94%).

Part B: Preparation of 1,1-dimethylethyl 4-[4-(trifluoromethyl)phenoxy]-1-

piperidinecarboxylate

To a solution of the BOC piperidine from part A (6.03 g, 30 mmol) in dimethylformamide (60 mL) was added cesium carbonate (9.77 g, 30 mmol) and 4-fluorobenzotrifluoride (3.8 mL, 30 mmol). The slurry was stirred at ninety degrees Celsius. After nineteen hours cesium carbonate (3.26g, 10 mmol) and 4-fluorobenzotrifluoride (0.95ml mL, 10 mmol) were added and the reaction continued at ninety degrees Celsius. After a total of forty six hours, the reaction was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water three times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo. Chromatography (on silica, ethyl acetate/hexanes) provided the substituted BOC piperidine as a white solid (6.0 g, 58%).

Part C: Preparation of 4-[4-(trifluoromethyl)-
phenoxy]piperidine

To a slurry of the substituted BOC piperidine from part B (5.95 g, 17.2 mmol) in 1,4-dioxane (10 mL) was added 4N HCl dioxane solution (17 mL). After one hour at ambient temperature the reaction was concentrated *in vacuo*. The residue was slurried in diethyl ether and vacuum filtration of the resulting precipitate provided the hydrochloride salt as a white solid (4.6 g, 100%).

Part D: Preparation of 1-(methylsulfonyl)-4-[4-(
trifluoromethyl)phenoxy]piperidine

To a solution of the hydrochloride salt from part C (4.6 g, 16.9 mmol) and triethylamine (5.9 mL, 42.4 mmol) in dichloromethane (45 mL) at zero degrees Celsius was added a solution of methane sulfonyl chloride (1.97 mL, 25.4 mmol) in dichloromethane (10 mL). After one hour at ambient temperature, the solvent was stripped *in vacuo*. The residue was taken up in ethyl acetate, washed with water two times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The residue was slurried in diethyl ether and vacuum filtration of the resulting precipitate provided the sulfonamide as an off-white solid (5.25 g, 96%).

Part E: Preparation of methyl [[4-[4-(
trifluoromethyl)phenoxy]-1-piperidinyl]-
sulfonyl]acetate

In dry equipment under nitrogen, the sulfonamide from part D (4.2 g, 13 mmol) was dissolved in dry tetrahydrofuran (26 mL), chilled to minus seventy-five degrees Celsius, and a 1 M solution of lithium bis(trimethylsilyl)amide (26 mL) was added

maintaining the temperature below minus sixty five degrees. After thirty minutes at minus seventy-five degrees Celsius, a solution of methyl chloroformate (1.0mL, 13 mmol) in dry tetrahydrofuran (13 mL) was added maintaining the temperature below minus sixty degrees. After thirty minutes at minus seventy-five degrees Celsius, the reaction was quenched with saturated ammonium chloride solution (100 mL) and extracted with ethyl acetate. The combined extracts were washed with saturated ammonium chloride solution, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo* to give the methylene sulfonamide as an yellow oil (4.95 g, 100%).

Part F: Preparation of methyl 1-(phenylmethyl)-4-[[4-(4-(trifluoromethyl)phenoxy]-1-piperidinyl)sulfonyl]-4-piperidinecarboxylate

To a solution of the methylene sulfonamide from part E (1.14 g, 3 mmol) in dimethylformamide (6 mL) was added potassium carbonate (1.24 g, 9 mmol), bis-(2-chloroethyl)benzyl amine (0.7 g, 3 mmol and 18-Crown-6 (500 mg). The slurry was stirred at sixty degrees Celsius. After sixteen hours the reaction was concentrated *in vacuo*. The residue was taken up in ethyl acetate, washed with water three times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Chromatography (on silica, ethyl acetate/hexanes) provided the piperidine sulfonamide as a white solid (950 mg, 59%).

Part G: Preparation of methyl 1-(5-ethyl-2-pyrimidinyl)-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl)sulfonyl]-4-

piperidinecarboxylate

To a slurry of the piperidine sulfonamide from part F (6.3 g, 12.0 mmol) in methanol (25 mL) was added ammonium formate (2.2 g, 34.5 mmol). The system was purged with nitrogen for 10 minutes. The nitrogen stream was removed and palladium on carbon (1.2 g of 10 weight % on activated carbon, 50% water) was added. The reaction was refluxed for forty five minutes, cooled, filtered through Celite under nitrogen, and concentrated *in vacuo*. The residue (4.5 g, 10 mmol) was dissolved in dry dimethylformamide (20 mL) and potassium carbonate (3.45 g, 325mmol) and 2-chloro-5-ethylpyrimidine (1.82 mL, 15 mmol) were added. The reaction was stirred at eighty degrees Celsius for four hour and then concentrated *in vacuo*. The residue was taken up in ethyl acetate and filtered through Celite. The filtrate washed with water two times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The product was recrystallized (methanol) to give the N-pyrimidinyl piperidine sulfonamide as a white solid (3.33 g, 60%).

Part H: Preparation of 1-(5-ethyl-2-pyrimidinyl)-4-
[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxylic
acid

In dry equipment under nitrogen, the N-pyrimidinyl piperidine sulfonamide from part G (3.3 g, 5.94 mmol) was dissolved in dry tetrahydrofuran (12 mL) and potassium trimethylsilonate (2.54 g, 17.8 mmol) was added at forty five degrees Celsius. After two hours water (100 mL) was added and at 5 degrees

Celsius, 6N HCl was added until pH=2. The slurry was filtered under nitrogen, the solids washed with hexanes and dried in vacuo to give the carboxylic acid as a white solid (3.11 g, 97%).

5 Part I: Preparation of 1-(5-ethyl-2-pyrimidinyl)-N-
[(tetrahydro-2H-pyran-2-yl)oxy]-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-
sulfonyl]-4-piperidinecarboxamide


In dry equipment under nitrogen, the carboxylic
10 acid from part H (3.0 g, 5.5mmol) was dissolved in dry dimethylformamide (15 mL) and the remaining reagents were added to the solution in the following order: N-hydroxybenzotriazole hydrate (0.9 g, 6.64 mmol), N-methylmorpholine (1.8 mL, 16.6 mmol), O-
15 (tetrahydro-2H-pyran-2-yl)hydroxylamine (1.9 g, 16.6 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.49 g, 7.75 mmol). The reaction was stirred at forty five degrees Celsius. After three hours, the reaction was
20 concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water, saturated NaHCO₃, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo. The product was recrystallized (methanol) to give the THP hydroxamate
25 as a white solid (3.34 g, 96%).

Part J: Preparation of 1-(5-ethyl-2-pyrimidinyl)-N-
hydroxy-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-4-
piperidinecarboxamide, monohydrochloride

30 To a solution of the THP hydroxamate from part I (3.3 g, 5.15 mmol) in 1,4-dioxane (3 mL) was added 4 N HCl dioxane solution (13 mL, 51.5 mmol) and methanol (1.3 mL). After thirty minutes at ambient

temperature the reaction was poured into 100 mL acetonitrile. The slurry was filtered under nitrogen, washed with acetonitrile and dried in vacuo to give the title compound as a white solid (2.59 g, 85%). HRMS (ES+) $M+H^+$ calculated for $C_{24}H_{30}F_3N_5O_5S_1$: 558.1998, found 558.1982.

Example 23: Preparation of tetrahydro-*N*-hydroxy-4-
[[4-[4-(trifluoromethoxy)phenoxy]-1-
piperidinyl]sulfonyl]-2*H*-thiopyran-4-
carboxamide



Part A: Preparation of 1,1-dimethylethyl 4-hydroxy-1-
piperidinecarboxylate

In dry equipment under nitrogen, 4-hydroxypiperidine (20.2 g, 0.2 mol) was dissolved in tetrahydrofuran (200 mL) and triethylamine (29 mL, 0.21 mol). A solution of di-*t*-butyldicarbonate (43.65 g, 0.2 mol) was added at such a rate that the temperature remained below thirty degrees Celsius. After stirring at ambient temperature for four hours, the reaction was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water, 5% $KHSO_4$, saturated $NaHCO_3$, saturated sodium chloride solution, dried over Na_2SO_4 , filtered, and concentrated in vacuo to give the BOC piperidine as a white solid (37.7 g, 94%).

Part B: Preparation of 4-(methylsulfonyl)hydroxy-1-

piperidinecarboxylic acid, 1,1-dimethylethyl
ester

To a solution of the BOC piperidine of part A (5.00 g, 24.84 mmol) in dichloromethane (50 mL) at zero degrees Celsius, was added triethylamine (3.81 mL, 27.32 mmol) followed by methane sulfonyl chloride (2.02 mL, 26.08 mmol). Once the addition was complete the cooling bath was removed. After stirring for two hours the reaction mixture was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water two times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo to provide the mesylate as an off-white solid (7.34 g, >100%).

Part C: Preparation of 4-[4-(trifluoromethoxy)-phenoxy]-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester

In dry equipment under nitrogen, 4-trifluoromethoxyphenol (10.15 g, 57 mmol) was dissolved in dry dimethylformamide (125 mL) and at minus five degrees Celsius sodium hydride (2.74g, 68.4 mmol of the 60% oil dispersion) was added and the ice bath was removed. After one hour at ambient temperature, the mesylate from part B (15.9 g, 57 mmol) was added and the reaction stirred at eighty degrees Celsius. After stirring at ambient temperature for four hours, the reaction was concentrated in vacuo. The residue was taken up in diethyl ether, washed with water, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo to provide the substituted BOC-piperidine as a beige solid (20.6 g, 100%).

Part D: Preparation of 4-[4-(trifluoromethoxy)-

phenoxy)piperidine

At fifteen degrees Celsius, 4 N HCl in dioxanes (125 mL) was slowly added to the substituted BOC-piperidine from part C (20.6 g, 57 mmol) and stirred
5 for ninety minutes. The reaction was concentrated in vacuo. The residue was dissolved in water (150 mL) and washed two times with ethyl acetate. The aqueous solution was cooled to five degrees Celsius and the pH adjusted to eleven with 5 N sodium hydroxide
10 solution and extracted with ethyl acetate. The ethyl acetate was dried over Na₂SO₄, filtered, and concentrated in vacuo to provide the substituted piperidine as a beige solid (11.9 g, 80%).

Part E: Preparation of 1-(methylsulfonyl)-4-[4-
15 (trifluoromethoxy)phenoxy)piperidine

The substituted piperidine from part D (11.5 g, 44.1 mmol) was dissolved in dichloromethane (125 mL) with triethylamine (12.3 mL, 88.1 mmol) and at zero
20 degrees Celsius a solution of methane sulfonyl chloride (5.1 mL, 66.1 mmol) in dichloromethane (20 mL) was added. After one hour at ambient temperature, the solvent was stripped in vacuo. The residue was taken up in ethyl acetate, washed with
25 water two times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo. The residue was recrystallized (ethyl acetate/hexanes) to give the sulfonamide as an off-white solid (10.77 g, 72%).

30 Part F: Preparation of methyl [[4-[4-
(trifluoromethoxy)phenoxy]-1-piperidinyl]-
sulfonyl]-4-piperidinecarboxylate

-431-

WO 00/46221

- In dry equipment under nitrogen, the sulfonamide from part E (10.77 g, 31.8 mmol) was dissolved in dry tetrahydrofuran (64 mL), chilled to minus seventy-five degrees Celsius, and a 1 M solution of lithium bis(trimethylsilyl)amide (80 mL) was added maintaining the temperature below minus sixty five degrees. After thirty minutes at minus sixty five degrees Celsius, a solution of methyl chloroformate (2.45 mL, 31.8 mmol) in dry tetrahydrofuran (32 mL) was added maintaining the temperature below minus sixty degrees. After thirty minutes at minus seventy-five degrees Celsius, the reaction was quenched with saturated ammonium chloride solution (125 mL) and extracted with ethyl acetate. The combined extracts were washed with saturated sodium chloride solution, saturated sodium sulfate, filtered, and concentrated in vacuo to give the methylene sulfonamide as an yellow oil (12.69 g, 100%).
- 20 Part G: Preparation of tetrahydro-4-[[4-[[4-sulfonyl]-2H-thiopyran-4-carboxylic acid (trifluoromethoxy)phenoxy]-1-piperidinyl]-
- To a solution of the methylene sulfonamide from part F (6.0 g, 15 mmol) in dimethylformamide (30 mL) were added potassium carbonate (6.2 g, 45 mmol), bis(2-bromoethyl)sulfide (3.72 g, 15 mmol; J.Chem.Soc.,1948;37) and 18-Crown-6 (500mg). The slurry was stirred at forty degrees Celsius. After twenty hours the reaction was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water three times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo to give impure

tetrahydrothiopyran substituted sulfonamide (4.68 g, 65%). In dry equipment under nitrogen, the tetrahydrothiopyran substituted sulfonamide (4.6 g, 9.52 mmol) was dissolved in dry tetrahydrofuran (20 mL) and potassium trimethylsilonate (4.07 g, 28.6 mmol) was added at fifty degrees Celsius. After four hours water (100 mL) was added and the solution concentrated *in vacuo*. The residue was taken up in water and extracted with ethyl acetate to remove impurities. The aqueous solution was treated with 6N HCl until pH=1. The slurry was extracted with ethyl acetate and the combined extracts washed with water, dried over Na₂SO₄, filtered, and concentrated in *vacuo*. The residue was recrystallized (acetone/hexanes) to give the carboxylic acid as a white solid (2.62 g, 59%).

Part H: Preparation of tetrahydro-*N*-[(tetrahydro-2H-pyran-2-yl)oxy]-4-[[4-[4-(trifluoromethoxy)-phenoxy]-1-piperidinyl]sulfonyl]-2H-thiopyran-4-carboxamide

In dry equipment under nitrogen, the carboxylic acid from part G (2.6 g, 5.54 mmol) was dissolved in dry dimethylformamide (11 mL) and the remaining reagents were added to the solution in the following order: N-hydroxybenzotriazole hydrate (0.9 g, 6.65 mmol), N-methylmorpholine (1.83 mL, 16.6 mmol), O-(tetrahydro-2H-pyran-2-yl)hydroxylamine (1.95 g, 16.6 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.49 g, 7.72 mmol). After two hour at thirty five degrees Celsius, the reaction was concentrated in *vacuo*. The residue was taken up in ethyl acetate, washed with water, saturated NaHCO₃, saturated sodium chloride solution,


-433-

dried over Na₂SO₄, filtered, and concentrated in vacuo. Chromatography (on silica, ethyl acetate/hexanes) provided the THP hydroxamate as a white foam (2.69 g, 85%).

5 Part I: Preparation of tetrahydro-*N*-hydroxy-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]-sulfonyl]-2*H*-thiopyran-4-carboxamide

To a solution of the THP hydroxamate from part H (0.99 g, 1.75 mmol) in 1,4-dioxane (3.5 mL) was added
10 4*N* HCl dioxane solution (0.9 mL, 3.5 mmol) and methanol (0.14 mL). After thirty minutes at ambient temperature the reaction was diluted with ethyl acetate and washed with water, dried over Na₂SO₄, filtered, and concentrated in vacuo. The product was
15 slurried in diethyl ether, filtered under nitrogen and dried to give the title compound as a white solid (0.65 g, 77%). HRMS (ES+) *M*+ *H*⁺ calculated for C₁₈H₂₃N₂O₆S₂F₃ : 485.1028, found 485.1034.

20 Example 24: Preparation of tetrahydro-*N*-hydroxy-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]sulfonyl]-2*H*-thiopyran-4-carboxamide, 1,1-dioxide



25

Part A: Preparation of 1,1-dimethylethyl 4-hydroxy-1-piperidinecarboxylate

In dry equipment under nitrogen, 4-
30 hydroxypiperidine (20.2 g, 0.2 mol) was dissolved in

tetrahydrofuran (200 mL) and triethylamine (29 mL, 0.21 mol). A solution of di-t-butylidicarbonate (43.65 g, 0.2 mol) was added at such a rate that the temperature remained below thirty degrees Celsius.

- 5 After stirring at ambient temperature for four hours, the reaction was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water, 5% KHSO₄, saturated NaHCO₃, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and
- 10 concentrated in vacuo to give the BOC piperidine as a white solid (37.7 g, 94%).

Part B: Preparation of 4-(methylsulfonyl)hydroxy-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester

- 15 To a solution of the BOC piperidine of part A (5.00 g, 24.84 mmol) in dichloromethane (50 mL) at zero degrees Celsius, was added triethylamine (3.81 mL, 27.32 mmol) followed by methane sulfonyl chloride (2.02 mL, 26.08 mmol). Once the addition was
- 20 complete the cooling bath was removed. After stirring for two hours, the reaction mixture was concentrated in vacuo. The residue was taken up in ethyl acetate, washed with water two times, saturated sodium chloride solution, dried over Na₂SO₄, filtered,
- 25 and concentrated in vacuo to provide the mesylate as an off-white solid (7.34 g, >100%).

Part C: Preparation of 4-[4-(trifluoromethoxy)-phenoxy]-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester

- 30 In dry equipment under nitrogen, 4-trifluoromethoxyphenol (10.15 g, 57 mmol) was dissolved in dry dimethylformamide (125 mL), and at minus five degrees Celsius sodium hydride (2.74g,

68.4 mmol of the 60% oil dispersion) was added and the ice bath was removed. After one hour at ambient temperature, the mesylate from part B (15.9 g, 57 mmol) was added and the reaction stirred at eighty
5 degrees Celsius. After stirring at ambient temperature for four hours, the reaction was concentrated in vacuo. The residue was taken up in diethyl ether, washed with water, saturated sodium chloride solution, dried over Na_2SO_4 , filtered, and
10 concentrated in vacuo to provide the substituted BOC-piperidine as a beige solid (20.6 g, 100%).

Part D: Preparation of 4-[4-(trifluoromethoxy)-
phenoxy]piperidine

At fifteen degrees Celsius, 4 N HCl in dioxanes
15 (125 mL) was slowly added to the substituted BOC-piperidine from part C (20.6 g, 57 mmol) and stirred for ninety minutes. The reaction was concentrated in vacuo. The residue was dissolved in water (150 mL) and washed two times with ethyl acetate. The aqueous
20 solution was cooled to five degrees Celsius and the pH adjusted to eleven with 5 N sodium hydroxide solution and extracted with ethyl acetate. The ethyl acetate was dried over Na_2SO_4 , filtered, and concentrated in vacuo to provide the substituted
25 piperidine as a beige solid (11.9 g, 80%).

Part E: Preparation of 1-(methylsulfonyl)-4-[4-
(trifluoromethoxy)phenoxy]piperidine

The substituted piperidine from part D (11.5 g, 44.1 mmol) was dissolved in dichloromethane (125 mL)
30 with triethylamine (12.3 mL, 88.1 mmol) and at zero degrees Celsius a solution of methane sulfonyl chloride (5.1 mL, 66.1 mmol) in dichloromethane (20 mL) was added. After one hour at ambient

temperature, the solvent was stripped in vacuo. The residue was taken up in ethyl acetate, washed with water two times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in
5 vacuo. The residue was recrystallized (ethyl acetate/hexanes) to give the sulfonamide as an off-white solid (10.77 g, 72%).

Part F: Preparation of methyl [[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]-
10 sulfonyl]-4-piperidinecarboxylate

In dry equipment under nitrogen, the sulfonamide from part E (10.77 g, 31.8 mmol) was dissolved in dry tetrahydrofuran (64 mL), chilled to minus seventy-five degrees Celsius, and a 1 M solution of lithium
15 bis(trimethylsilyl)amide (80 mL) was added maintaining the temperature below minus sixty five degrees. After thirty minutes at minus seventy-five degrees Celsius, a solution of methyl chloroformate (2.45 mL, 31.8 mmol) in dry tetrahydrofuran (32 mL)
20 was added maintaining the temperature below minus sixty degrees. After thirty minutes at minus seventy-five degrees Celsius, the reaction was quenched with saturated ammonium chloride solution (125 mL) and extracted with ethyl acetate. The
25 combined extracts were washed with saturated ammonium chloride solution, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo to give the methylene sulfonamide as an yellow oil (12.69 g, 100%).
30 Part G: Preparation of tetrahydro-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]-sulfonyl]-2H-thiopyran-4-carboxylic acid

To a solution of the methylene sulfonamide from part F (6.0 g, 15 mmol) in dimethylformamide (30 mL) was added potassium carbonate (6.2 g, 45 mmol), bis-(2-bromoethyl)sulfide (3.72 g, 15 mmol;
5 *J.Chem.Soc.*;1948;37) and 18-Crown-6 (500 mg). The slurry was stirred at forty degrees Celsius. After twenty hours the reaction was concentrated *in vacuo*. The residue was taken up in ethyl acetate, washed with water three times, saturated sodium chloride
10 solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo* to give impure tetrahydrothiopyran substituted sulfonamide (4.68 g, 65%). In dry equipment under nitrogen, the tetrahydrothiopyran substituted sulfonamide (4.6 g,
15 9.52 mmol) was dissolved in dry tetrahydrofuran (20 mL) and potassium trimethylsilonate (4.07 g, 28.6 mmol) was added at fifty degrees Celsius. After four hours, water (100 mL) was added and the solution concentrated *in vacuo*. The residue was taken up in
20 water and extracted with ethyl acetate to remove impurities. The aqueous solution was treated with 6N HCl until pH=1. The slurry was extracted with ethyl acetate and the combined extracts washed with water, dried over Na₂SO₄, filtered, and concentrated in
25 *vacuo*. The residue was recrystallized (acetone/hexanes) to give the carboxylic acid as a white solid (2.62 g, 59%).

Part H: Preparation of tetrahydro-*N*-[(tetrahydro-2*H*-
pyran-2-yl)oxy]-4-[[4-[4-(trifluoromethoxy)-
30 phenoxy]-1-piperidinyl]sulfonyl]-2*H*-
thiopyran-4-carboxamide

In dry equipment under nitrogen, the carboxylic acid from part G (2.6 g, 5.54 mmol) was dissolved in

dry dimethylformamide (11 mL) and the remaining reagents were added to the solution in the following order: N-hydroxybenzotriazole hydrate (0.9 g, 6.65 mmol), N-methylmorpholine (1.83 mL, 16.6 mmol), O-
5 (tetrahydro-2H-pyran-2-yl)hydroxylamine (1.95 g, 16.6 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.49 g, 7.72 mmol). After two hour at thirty five degrees Celsius, the reaction was concentrated *in vacuo*. The residue was
10 taken up in ethyl acetate, washed with water, saturated NaHCO₃, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in *vacuo*. Chromatography (on silica, ethyl acetate/hexanes) provided the THP hydroxamate as a
15 white foam (2.69 g, 85%).

Part I: Preparation of tetrahydro-N-[(tetrahydro-2H-pyran-2-yl)oxy]-4-[[4-[4-(trifluoromethoxy)-phenoxy]-1-piperidinyl]sulfonyl]-2H-
thiopyran-4-carboxamide, 1,1-dioxide

20

The THP hydroxamate from part H (1.0 g, 1.76 mmol) was dissolved in dichloromethane (10 mL) and 3-chloroperoxybenzoic acid (1.33g, 4.4 mmol, 57-86%) was added at twenty five degrees Celsius. After two
25 hours, a solution of saturated NaHCO₃ with 5% sodium thiosulfate (10 mL) was added and the mixture was stirred for ten minutes. The layers were separated and washed with saturated NaHCO₃, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and
30 concentrated *in vacuo*. Chromatography (on silica, ethyl acetate/hexanes) provided the sulfone THP hydroxamate as a white solid (0.907 g, 92%).

Part J: Preparation of tetrahydro-N-hydroxy-4-[[4-[4-

(trifluoromethoxy)phenoxy]-1-piperidinyl]-
sulfonyl]-2*H*-thiopyran-4-carboxamide, 1,1-
dioxide

To a solution of the sulfone THP hydroxamate
5 from part I (0.9 g, 1.5 mmol) in 1,4-dioxane (3. mL)
was added 4*N* HCl dioxane solution (1.9 mL, 7.5 mmol)
and methanol (0.5 mL). After ten minutes at ambient
temperature the reaction was diluted with ethyl
acetate and washed with water, dried over Na₂SO₄,
10 filtered, and concentrated *in vacuo*. The product was
slurried in diethyl ether, filtered under nitrogen
and dried to give the title compound as a white
solid (0.70 g, 90%). HRMS (ES+) M+ NH₄⁺ calculated
for C₁₈H₂₃N₂O₈S₂F₃ : 534.1192, found 534.1231.

15

Example 25: Preparation of tetrahydro-*N*-hydroxy-4-
[[4-[4-(trifluoromethyl)phenoxy]-1-
piperidinyl]sulfonyl]-2*H*-thiopyran-4-
carboxamide



20

Part A: Preparation of 1,1-dimethylethyl 4-hydroxy-1-
piperidinecarboxylate

In dry equipment under nitrogen, 4-
25 hydroxypiperidine (20.2 g, 0.2 mol) was dissolved in
tetrahydrofuran (200 mL) and triethylamine (29 mL,
0.21 mol). A solution of di-*t*-butyldicarbonate
(43.65 g, 0.2 mol) was added at such a rate that the
temperature remained below thirty degrees Celsius.
30 After stirring at ambient temperature for four hours,

-440-

the reaction was concentrated *in vacuo*. The residue was taken up in ethyl acetate, washed with water, 5% KHSO₄, saturated NaHCO₃, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and

5 concentrated *in vacuo* to give the BOC piperidine as a white solid (37.7 g, 94%).

Part B: Preparation of 1,1-dimethylethyl 4-[4-(trifluoromethyl)phenoxy]-1-
piperidinecarboxylate

10 To a solution of the BOC piperidine from part A (6.03 g, 30 mmol) in dimethylformamide (60 mL) was added cesium carbonate (9.77 g, 30 mmol) and 4-fluorobenzotrifluoride (3.8 mL, 30 mmol). The slurry was stirred at ninety degrees Celsius. After
15 nineteen hours cesium carbonate (3.26g, 10 mmol) and 4-fluorobenzotrifluoride (0.95ml mL, 10 mmol) were added and the reaction continued at ninety degrees Celsius. After a total of forty six hours, the reaction was concentrated *in vacuo*. The residue was
20 taken up in ethyl acetate, washed with water three times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Chromatography (on silica, ethyl acetate/hexanes) provided the substituted BOC piperidine as a white
25 solid (6.0 g, 58%).

Part C: Preparation of 4-[4-(trifluoromethyl)-
phenoxy]piperidine

To a slurry of the substituted BOC piperidine from part B (5.95 g, 17.2 mmol) in 1,4-dioxane (10
30 mL) was added 4 N HCl dioxane solution (17 mL). After one hour at ambient temperature the reaction was concentrated *in vacuo*. The residue was slurried in diethyl ether and vacuum filtration of the

resulting precipitate provided the hydrochloride salt as a white solid (4.6 g, 100%).

Part D: Preparation of 1-(methylsulfonyl)-4-[4-(trifluoromethyl)phenoxy]piperidine

5

To a solution of the hydrochloride salt from part C (4.6 g, 16.9 mmol) and triethylamine (5.9 mL, 42.4 mmol) in dichloromethane (45 mL) at zero degrees Celsius was added a solution of methane sulfonyl chloride (1.97 mL, 25.4 mmol) in dichloromethane (10 mL). After one hour at ambient temperature, the solvent was stripped in vacuo. The residue was taken up in ethyl acetate, washed with water two times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated in vacuo. The residue was slurried in diethyl ether and vacuum filtration of the resulting precipitate provided the sulfonamide as an off-white solid (5.25 g, 96%).

Part E: Preparation of methyl [[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-sulfonyl]acetate

In dry equipment under nitrogen, the sulfonamide from part D (4.2 g, 13 mmol) was dissolved in dry tetrahydrofuran (26 mL), chilled to minus seventy-five degrees Celsius, and a 1 M solution of lithium bis(trimethylsilyl)amide (26 mL) was added maintaining the temperature below minus sixty five degrees. After thirty minutes at minus seventy-five degrees Celsius, a solution of methyl chloroformate (1.0mL, 13 mmol) in dry tetrahydrofuran (13 mL) was added maintaining the temperature below minus sixty degrees. After thirty minutes at minus seventy-five degrees Celsius, the reaction was quenched with

saturated ammonium chloride solution (100 mL) and extracted with ethyl acetate. The combined extracts were washed with saturated ammonium chloride solution, saturated sodium chloride solution, dried
5 over Na₂SO₄, filtered, and concentrated *in vacuo* to give the methylene sulfonamide as an yellow oil (4.95 g, 100%).

Part F: Preparation of tetrahydro-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-
10 sulfonyl]-2H-thiopyran-4-carboxylic acid

To a solution of the methylene sulfonamide from part E (5.7 g, 15 mmol) in dimethylformamide (30 mL) was added potassium carbonate (6.2 g, 45 mmol), bis-
15 (2-bromoethyl)sulfide (3.72 g, 15 mmol; *J.Chem.Soc.*;1948;37) and 18-Crown-6 (500 mg). The slurry was stirred at forty degrees Celsius. After sixty hours the reaction was concentrated *in vacuo*. The residue was taken up in ethyl acetate, washed
20 with water three times, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo* to give impure tetrahydrothiopyran substituted sulfonamide (3.7 g, 53%). In dry equipment under nitrogen, the
25 tetrahydrothiopyran substituted sulfonamide (3.68 g, 7.88 mmol) was dissolved in dry tetrahydrofuran (15 mL) and potassium trimethylsilonate (3.37 g, 23.6 mmol) was added at fifty degrees Celsius. After ninety minutes water (100 mL) was added and the
30 solution concentrated *in vacuo*. The residue was taken up in water and extracted with ethyl acetate to remove impurities. The aqueous solution was treated with 6 N HCl until pH=1. The slurry was extracted

with ethyl acetate and the combined extracts washed with water, dried over Na₂SO₄, filtered, and concentrated *in vacuo* to give the carboxylic acid as a white foam (1.66 g, 46%).

5 Part G: Preparation of tetrahydro-*N*-[(tetrahydro-2*H*-pyran-2-yl)oxy]-4-[[4-[4-(trifluoromethyl)-phenoxy]-1-piperidinyl]sulfonyl]-2*H*-thiopyran-4-carboxamide

In dry equipment under nitrogen, the carboxylic
10 acid from part F (1.5 g, 3.31 mmol) was dissolved in dry dimethylformamide (7 mL) and the remaining reagents were added to the solution in the following order: *N*-hydroxybenzotriazole hydrate (0.54 g, 3.97 mmol), *N*-methylmorpholine (1.1 mL, 9.93 mmol), *O*-
15 (tetrahydro-2*H*-pyran-2-yl)hydroxylamine (1.16 g, 9.93 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.89 g, 4.64 mmol). After ninety minutes at thirty five degrees Celsius, the reaction was concentrated *in vacuo*. The residue
20 was taken up in ethyl acetate, washed with water, saturated NaHCO₃, saturated sodium chloride solution, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Chromatography (on silica, ethyl acetate/hexanes) provided the THP hydroxamate as a
25 white foam (0.85 g, 47%).

Part H: Preparation of tetrahydro-*N*-hydroxy-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-sulfonyl]-2*H*-thiopyran-4-carboxamide


30 To a solution of the THP hydroxamate from part G (0.7 g, 1.27 mmol) in 1,4-dioxane (2.5 mL) was added 4 N HCl dioxane solution (1.6 mL, 6.34 mmol) and methanol (0.4 mL). After ten minutes at ambient

- 444 -

temperature the reaction was diluted with ethyl acetate and washed with water, dried over Na₂SO₄, filtered, and concentrated in vacuo. Dichloromethane (20 mL) was added and the solution was stripped. The product was slurried in diethyl ether, filtered under nitrogen and dried to give the title compound as a white solid (0.56 g, 94%). HRMS (ES+) M+ H⁺ calculated for C₁₈H₂₃N₂O₅S₂F₃ : 469.1079, found 469.1061.


10

Example 26: Preparation of *N*-hydroxy-4[[1'-(*n*-pentyl)[4,4'-bipiperidin]-1-yl]-sulfonyl]-tetrahydro-2*H*-pyran-4-carboxamide



15

Part 1: Preparation of




20

To a solution of the *N*-(*t*-butoxycarbonyl)-4,4'-bipiperidine (prepared using Preparation 22 in patent application WO 94/14776) (32.3 g, 120.0 mmol) and triethylamine (30.1 mL, 216.0 mmol) in dichloromethane (330 mL) at zero degrees Celsius was

added a solution of methane sulfonyl chloride (16.2 mL, 209.0 mmol) in dichloromethane (100 mL). After 2.5 hours at ambient temperature, the solvent was removed in vacuo. The residue was partitioned
5 between ethyl acetate (1 L) and water (0.5 L). The aqueous layer was extracted twice with ethyl acetate (300 mL), then the combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated in vacuo. The residue was triturated
10 with diethyl ether to give the methyl sulfonamide N'-t-butoxycarbamate as a white solid (33.31 g, 80%). ¹H NMR and mass spectrum were consistent with the desired compound.

Part 2: Preparation of

15




In dry equipment under nitrogen, the methyl sulfonamide N'-t-butoxycarbamate from part 1 (28.0 g, 81.0 mmol) was dissolved in dry tetrahydrofuran (160 mL) and cooled to minus seventy-five degrees Celsius.
20 The resulting solution was then treated with a 1 M solution of lithium bis(trimethylsilyl)amide (210 mL, 210.0 mmol) at a rate such that the temperature remained below minus sixty five degrees. After
25 addition was complete, the reaction mixture was allowed to warm to zero degrees Celsius. After 1 hour, the solution was cooled to minus seventy-five degrees Celsius and treated with a solution of methyl chloroformate (8.2 mL, 97.0 mmol) in dry

-446-

tetrahydrofuran (50 mL), while maintaining the temperature below minus seventy degrees. After 1 hour at minus seventy-five degrees Celsius, the reaction was quenched with saturated ammonium chloride aqueous solution (60 mL), warmed to ambient temperature and concentrated *in vacuo*. The resulting residue was partitioned between ethyl acetate (500 mL) and 5% aqueous KHSO₄ (500 mL). The aqueous layer was extracted with ethyl acetate (250 mL), then the combined organic layers were washed with water (250 mL), twice with brine (2 x 50 mL), dried over Na₂SO₄, filtered, and concentrated *in vacuo* to give the methylene sulfonamide N'-t-butoxycarbamate as a yellow solid (32.6 g, 99.9%). ¹H NMR and mass spectrum were consistent with the desired compound.


Part 3: Preparation of



To a solution of the methylene sulfonamide N'-t-butoxycarbamate from part 2 (15.0 g, 37.0 mmol) in dimethylformamide (75 mL) was added bis-(2-bromoethyl)ether (10.3 g, 44 mmol), 18-Crown-6 (400 mg), followed by potassium carbonate (15.4 g, 111.0 mmol). The heterogeneous mixture was stirred at sixty degrees Celsius. After 48 hours, the reaction was concentrated *in vacuo*, and the resulting oil was partitioned between ethyl acetate (300 mL) and water (200 mL). The organic layer was washed with 5% aqueous KHSO₄ (50 mL), twice with water (2 x 50 mL),


brine (50 mL), dried over Na_2SO_4 , filtered, and concentrated in vacuo to give the pyran sulfonamide N'-t-butoxycarbamate as a white solid. (18 g, 97%). ^1H NMR and mass spectrum were consistent with the
5 desired compound.

Part 4: Preparation of



The pyran sulfonamide N'-t-butoxycarbamate (17 g, 36 mmol) from part 3 was dissolved in 4 N HCl 1,4-dioxane solution (90 mL). After 2 hours at ambient
10 temperature the clear yellow solution began to form a precipitate. After 4 hours, the reaction was diluted with diethyl ether and vacuum filtration of the resulting white suspension provided the pyran
15 sulfonamide hydrochloride salt as a white solid (13.4 g, 91%). ^1H NMR and mass spectrum were consistent with the desired compound.

Part 5: Preparation of




20

A suspension of the pyran sulfonamide hydrochloride salt from part 4 (3.5 g, 8.5 mmol) in tetrahydrofuran (26 mL) was treated with sodium acetate (0.7 g, 8.9 mmol), valeraldehyde (0.9 mL, 8.5
25 mmol), followed by sodium triacetoxyborohydride (2.2 g, 10.0 mmol). After 96 hours at ambient temperature, the reaction mixture was concentrated in


vacuo. The resulting residue was partitioned between ethyl acetate (100 mL) and water (70 mL) and treated with saturated sodium carbonate aqueous solution until pH = 8. The aqueous layer was extracted with ethyl acetate (25 mL) and the combined organic layers were washed with water (20 mL), brine (30 mL), dried over Na₂SO₄, filtered, and concentrated *in vacuo* to give the pyran sulfonamide N'-n-pentyl amine as a light yellow solid (3.7 g, 97%).¹H NMR and mass spectrum were consistent with the desired compound.

Part 6: Preparation of



In dry equipment under nitrogen, the pyran sulfonamide N'-n-pentyl amine from part 5 (3.2 g, 7.2 mmol) was dissolved in dry tetrahydrofuran (36 mL) and potassium trimethylsilanolate (3.1 g, 22.0 mmol) was added at ambient temperature. After 21 hours, the reaction was concentrated *in vacuo*, and the resulting residue was dissolved in water (20 mL) and treated with 2N HCl until pH = 7. The white suspension was vacuum filtered, washed with water and dried *in vacuo* to give the carboxylic acid as a white solid (2.36 g, 76%).¹H NMR and mass spectrum were consistent with the desired compound.


Part 7: Preparation of



In dry equipment under nitrogen, the carboxylic acid from part 6 (2.2 g, 5.1 mmol) was dissolved in dry dimethylformamide (17 mL) and treated with 1-(3-dimethylamino-propyl)-3-ethylcarbodiimide hydrochloride (1.5 g, 7.63 mmol) and N-hydroxybenzo-triazole hydrate (1.03 g, 7.63 mmol). The resulting suspension became a clear amber solution after stirring at fifty degrees Celsius for 1.5 hours. The reaction was then treated with O-(tetrahydro-2H-pyran-2-yl)hydroxylamine (0.9 g, 7.63 mmol), followed by N-methylmorpholine (1.7 mL, 15 mmol) three minutes later. The reaction was stirred at fifty degrees Celsius. After 64 hours, the reaction was treated again with 1-(3-dimethylamino-propyl)-3-ethylcarbodiimide hydrochloride (0.49 g, 2.6 mmol), N-hydroxybenzo-triazole hydrate (0.34 g, 2.6 mmol), O-(tetrahydro-2H-pyran-2-yl)hydroxylamine (0.30 g, 2.6 mmol), followed by N-methylmorpholine (0.28 mL, 2.6 mmol). After 98 hours, the reaction was concentrated in vacuo. The resulting residue was partitioned between ethyl acetate (100 mL) and water (40 mL). The organic layer was washed three times with saturated sodium bicarbonate aqueous solution (3 x 25 mL), twice with brine (2 x 25 mL), dried over Na₂SO₄, filtered, and concentrated in vacuo. The resulting oil was recrystallized in methanol (4 mL) to give the THP hydroxamate as a white solid (1.28 g, 48%). ¹H NMR and mass spectrum were consistent with the desired compound.


Part 8: Preparation of

-450-




The THP hydroxamate from part 7 (1.1 g, 2.08 mmol) was suspended in methanol (0.5 mL) and the treated with 4 N HCl in 1,4-dioxane solution (5.2 mL, 21 mmol). After 2 hours at ambient temperature, the reaction was concentrated to half the reaction volume and then diluted with diethyl ether (200 mL). The white suspension was filtered under nitrogen and dried in vacuo to give the title compound as a white solid (0.94 g, 94%). MS(ES+) m/z calculated for C₂₁H₃₉N₃O₅S: 445, found (M+1) 446.

Example 27: Preparation of *N*-hydroxy-4[[1'-(4-methoxybenzoyl)[4,4'-bipiperidin]-1-yl]sulfonyl]-tetrahydro-2*H*-pyran-4-carboxamide



Part 9: Preparation of




A suspension of the pyran sulfonamide hydrochloride salt from part 4 of Example 26 (3.0 g, 7.3 mmol), in dichloromethane (15 mL) and

-451-

triethylamine (1.12 mL, 8.0 mmol) was cooled to zero degrees Celsius and treated with 4-(dimethylamino)pyridine (0.1g) followed by *p*-anisoyl chloride (1.37 g, 8.0 mmol). The cooling bath was removed and the reaction was stirred at ambient temperature for 22 hours. The reaction was again treated with triethylamine (0.51 mL, 3.65 mmol) *p*-anisoyl chloride (0.62 g, 3.65 mmol). After stirring for 8 days, the reaction was diluted with dichloromethane (75 mL) and partitioned with water (50 mL). The aqueous layer was extracted twice with dichloromethane (2 x 25 mL). The combined organic layers were washed twice with 5% aqueous KHSO₄ (2 x 30 mL), water (20 mL), saturated sodium bicarbonate aqueous solution (25 mL), brine (10 mL), dried over Na₂SO₄, filtered, and concentrated *in vacuo* to give the amide methyl ester as a white solid. (3.7 g, crude).

The amide methyl ester was suspended in dry tetrahydrofuran (42 mL) and treated with potassium trimethylsilanolate (2.73 g, 21.0 mmol) and stirred at forty degrees Celsius. After 4.5 days, the reaction was diluted with water (50 mL) and treated with 2 N HCl until pH = 1. The resulting white, yellow suspension was vacuum filtered, washed with water and dried *in vacuo* to give the carboxylic acid as a light yellow solid (2.55 g, 82% over 2 steps). ¹H NMR and mass spectrum were consistent with the desired compound.


Part 10: Preparation of



In dry equipment under nitrogen, the carboxylic acid from part 9 (2.2 g, 4.6 mmol) was dissolved in dry dimethylformamide (15 mL) and treated with 1-(3-dimethylamino-propyl)-3-ethylcarbodiimide hydrochloride (1.3 g, 6.7 mmol) and N-hydroxybenzotriazole hydrate (0.9 g, 6.7 mmol). The resulting suspension became a clear amber solution after stirring at fifty degrees Celsius for 50 minutes.


The reaction was then treated with O-(tetrahydro-2H-pyran-2-yl)hydroxylamine (0.8 g, 6.7 mmol), followed by N-methylmorpholine (1.47 mL, 13 mmol) three minutes later. The reaction was stirred at fifty degrees Celsius. After 25 hours, the reaction was concentrated *in vacuo*. The resulting residue was partitioned between ethyl acetate (100 mL) and water (50 mL). The aqueous layer was extracted with ethyl acetate (25 mL), and the combined organic layers were washed twice with saturated sodium bicarbonate aqueous solution (2 x 25 mL), brine (25 mL), dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The resulting yellow oil was recrystallized in methanol (4 mL) to give the THP hydroxamate as a white solid (1.81 g, 69%). ¹H NMR and mass spectrum were consistent with the desired compound.

Part 11: Preparation of




The THP hydroxamate from part 10 (1.44 g, 2.43 mmol) was dissolved in acetonitrile (25 mL), diluted with water (15 mL), and then treated with aqueous 2N HCl (2.5 mL, 4.85 mmol). After 2 hours at ambient temperature, the acetonitrile and excess hydrochloric acid were removed with a stream of nitrogen. The resulting white suspension was filtered under nitrogen, washed with water and dried in vacuo to give the title compound as a white solid (0.67 g, 56%). MS(ES+) m/z calculated for C₂₄H₃₅N₃O₇S: 509; found (M+1) 510.

Example 28: Preparation of N-hydroxy-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]-sulfonyl]-4-piperidinecarboxamide, monohydrochloride




Part 12: Preparation of



-454-

In dry equipment under nitrogen, the compound of Example 20, part F (16.51 g, 30.5 mmol) was dissolved in dry tetrahydrofuran (61 mL) and potassium trimethylsilanolate (11.8 g, 91.6 mmol) was added at ambient temperature. After 21 hours, the reaction was concentrated *in vacuo*, and the resulting residue was dissolved in water (100 mL) and treated with 2N HCl until pH = 7. The white suspension was vacuum filtered, washed with water and dried *in vacuo* to give the carboxylic acid as a white solid (15.45 g, 96%). ¹H NMR and mass spectrum were consistent with the desired compound.


Part 13: Preparation of



In dry equipment under nitrogen, the carboxylic acid from part 12 (15.45 g, 29.3 mmol) was dissolved in dry dimethylformamide (147 mL) and treated with 1-(3-dimethylamino-propyl)-3-ethylcarbodiimide hydrochloride (8.44 g, 44 mmol) and N-hydroxybenzotriazole hydrate (5.95 g, 44 mmol). After 1 hour and 20 minutes at ambient temperature, the suspension was treated with O-(tetrahydro-2H-pyran-2-yl)hydroxylamine (5.15 g, 44 mmol), followed by N-methylmorpholine (9.7 mL, 88 mmol) three minutes later. After stirring for 16 hours at ambient temperature, the reaction was heated to fifty degrees Celsius for 3 hours, and then treated with 1-(3-dimethylamino-propyl)-3-ethylcarbodiimide hydrochloride (1.41 g, 7.4 mmol), N-hydroxybenzo-

triazole hydrate (0.99 g, 7.3 mmol), O-(tetrahydro-2H-pyran-2-yl)hydroxylamine (0.86 g, 7.3 mmol), and N-methylmorpholine (1.61 mL, 14.6 mmol). After 20 hours, the reaction was concentrated *in vacuo*, and the resulting residue was partitioned between ethyl acetate (400 mL) and water (400 mL). The aqueous layer was extracted with ethyl acetate (200 mL), and the combined organic layers were washed with saturated sodium bicarbonate aqueous solution (100 mL), water (100 mL), brine (100 mL), dried over Na₂SO₄, filtered, and concentrated *in vacuo* to give the THP hydroxamate piperidine N-benzyl as a brown oil (18.4 g, 100%). ¹H NMR and mass spectrum were consistent with the desired compound.


Part 14: Preparation of



The THP hydroxamate piperidine N-benzyl from part 13 (1.0 g, 1.6 mmol) was dissolved in methanol (4.5 mL) and then treated with ammonium formate (302 mg, 4.8 mmol) followed by palladium on carbon (Degussa catalyst, 400 mg of 10 weight % on activated carbon, 50% water). The black heterogeneous mixture was stirred at ambient temperature for 30 minutes, diluted with methanol (4.5 mL), and then stirred for another 3 hours. The reaction was then filtered through a methanol washed pad of Celite under nitrogen, and the filtrate was concentrated *in vacuo* to give the THP hydroxamate piperidine as a white glassy foam (0.73 g, 85%). ¹H NMR and mass spectrum were consistent with the desired compound.


-456-

Part 15: Preparation of



The THP hydroxamate piperidine from part 14 (0.34 g, 0.64 mmol) was treated with methanol (0.2 mL) followed by 4 N HCl in 1,4-dioxane solution (2 mL, 8 mmol), which formed a white precipitate. After 1 hour at ambient temperature, the reaction diluted with diethyl ether (5 mL), and the white suspension was filtered under nitrogen, washed with diethyl ether and dried *in vacuo* to give the title compound as a white solid (0.18 g, 58%). MS(FABMS) *m/z* calculated for $C_{18}H_{24}N_3O_5SF_3$: 451, found (*M*+1) 452.

Example 29: Preparation of bis(2-chloroethyl)-
benzylamine (Part 16)




A suspension of bis(2-chloroethyl)amine hydrochloride (Aldrich, 500 g, 2.8 mol), sodium acetate (229.8 g, 2.8 mol), and benzaldehyde (270.5 mL, 2.66 mol), in tetrahydrofuran (2.5 L), was cooled to ten degrees Celsius, and treated with sodium triacetoxyborohydride (712.4 g, 3.36 mol) at a rate such that the temperature did not exceed eighteen degrees Celsius. After stirring the white suspension at ambient temperature for 25 hours, the reaction was quenched with the addition of ethyl acetate (4 L) followed by 2.5 M sodium hydroxide (3.5 L), which

-457-

made the mixture pH = 9. The mixture was partitioned with the addition of water (3 L). The aqueous layer was extracted with ethyl acetate (1.5 L). The combined organic layers were washed twice with brine (2 x 1 L), dried over MgSO₄, filtered, and concentrated *in vacuo*. The crude yellow oil was purified on silica gel (hexanes) to give a clear, colorless oil (482.6 g, 78%). ¹H NMR and mass spectrum were consistent with the desired compound.


10

Example 30: Preparation of 1-(2-furanylmethyl)-N-hydroxy-4-[[4-[4-(trifluoromethyl)-phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide



15

Part 17: Preparation of




To a solution of the piperidine methyl ester from Part 1 of Example 42 (3.91 g, 8.7 mmol) and 2-furaldehyde (0.79 mL, 9.6 mmol) in dichloroethane (58 mL) was added glacial acetic acid (0.5 mL, 8.7 mmol) followed by sodium triacetoxyborohydride (2.4 g, 11.3 mmol). After 64 hours, the reaction was concentrated *in vacuo* and partitioned between ethyl acetate (75

25

-458-

mL) and water (75 mL). The aqueous layer was made basic (pH = 8) with saturated sodium bicarbonate aqueous solution, then extracted three times with ethyl acetate (3 x 50 mL). The combined organic
5 layers were washed with water (25 mL), twice with brine (2 x 25 mL), dried with MgSO_4 , filtered, and concentrated *in vacuo*. Recrystallization of the crude oil in methanol yielded a brown, tan solid, which was suspended in tetrahydrofuran (17.4 mL) and
10 treated with potassium trimethylsilanolate (3.34 g, 26.0 mmol). After 20 hours at ambient temperature, the reaction was diluted with tetrahydrofuran (10 mL) and charged again with potassium trimethylsilanolate (2.22 g, 17.3 mmol). After 7 hours, the reaction was
15 concentrated *in vacuo*, and the resulting residue was dissolved in water (100 mL) and treated with 2 N HCl until pH = 7. The white suspension was vacuum filtered, washed with water and dried *in vacuo* to give the carboxylic acid as a white solid (4.4 g,
20 97%). ^1H NMR and mass spectrum were consistent with the desired compound.


Part 18: Preparation of



In dry equipment under nitrogen, the carboxylic
25 acid from part 17 (3.93 g, 7.61 mmol) was dissolved in dry dimethylformamide (38 mL) and treated with 1-(3-dimethylamino-propyl)-3-ethylcarbodiimide hydrochloride (2.2 g, 11.4 mmol), N-hydroxybenzotriazole hydrate (1.54 g, 11.4 mmol), O-(tetrahydro-

2H-pyran-2-yl)hydroxylamine (1.34 g, 11.4 mmol), followed by N-methylmorpholine (2.51 mL, 22.8 mmol). After 17 hours at ambient temperature, the reaction was concentrated *in vacuo*, and the resulting residue was partitioned between ethyl acetate (175 mL) and water (175 mL). The aqueous layer was extracted with ethyl acetate (100 mL), and the combined organic layers were washed with saturated sodium bicarbonate aqueous solution (100 mL), water (100 mL), brine (100 mL), dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The resulting yellow oil was recrystallized in methanol (6 mL) to give the THP-hydroxamate as a white solid (2.48 g, 53%). ¹H NMR and mass spectrum were consistent with the desired compound.


Part 19: Preparation of




The THP-hydroxamate from part 18 (2.41 g, 3.91 mmol) was dissolved in methanol (1.0 mL) and 1,4-dioxane (15 mL), then treated with 4 N HCl in 1,4-dioxane solution (10 mL, 39 mmol), which formed a tan precipitate. After 1 hour at ambient temperature, the reaction was diluted with acetonitrile (3 mL), and the suspension was filtered under nitrogen, washed with acetonitrile and dried *in vacuo* to give the title compound as a white solid (1.68 g, 81%). MS(EI) m/z calculated for C₂₃H₂₈N₃O₆SF₃: 531, found (M+1) 532.

Example 31: Preparation of 4-[[4-[4-[4-

(trifluoromethyl)phenoxy]phenoxy]-1-
piperidinyl]sulfonyl]tetrahydro-*N*-
hydroxy-2*H*-pyran-4-carboxamide



Part 20: Preparation of




In dry equipment under nitrogen, 4-[4-(trifluoromethyl)phenoxy]phenol (Aldrich, 6.0 g, 23.6 mmol) was added to a dimethylformamide (53 mL) suspension of sodium hydride (60% dispersion in mineral oil, 0.95 g, 23.6 mmol), which was pre-washed in hexanes, while at zero degrees Celsius. After addition, the cold bath was removed and the reaction was warmed to ambient temperature. After 30 minutes, the reaction was cooled to zero degrees Celsius, charged with 4-(methylsulfonyl)hydroxy-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester (Example 23, part B) (5.5 g, 19.7 mmol), then heated to eighty degrees Celsius. After 17 hours, the reaction was again treated with sodium hydride (60% dispersion in mineral oil, 0.95 g, 23.6 mmol), and then 10 minutes later with 4-(methylsulfonyl)hydroxy-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester (5.5 g, 19.7 mmol). After 47 hours, the reaction was cooled to ambient temperature, quenched with water (10 mL), concentrated in vacuo, and partitioned between diethyl ether (100 mL) and water (100 mL).

The aqueous layer was extracted twice with diethyl ether (2 x 25 mL) and then with ethyl acetate (50 mL). The combined organic layers were washed with water (25 mL), twice with brine (2 x 25 mL), dried
5 over Na_2SO_4 , filtered, concentrated and dried *in vacuo* to give a brown oil.

The crude oil was treated with 4N HCl in 1,4-dioxane solution (47 mL, 188 mmol) at ambient temperature. After 2 hours, the reaction was
10 concentrated *in vacuo*, and partitioned between ethyl acetate (100 mL) and water (50 mL). The aqueous layer was made pH = 8 with the addition of saturated sodium bicarbonate aqueous solution, then extracted twice with ethyl acetate (2 x 25 mL). The combined
15 organic layers were washed with saturated sodium bicarbonate aqueous solution (50 mL), water (50 mL), brine (25 mL), dried over Na_2SO_4 , filtered, concentrated and dried *in vacuo* to give a brown oil. The crude product was purified on silica gel
20 (hexanes/ ethyl acetate) to give the amine as a yellow, orange solid (5.5 g, 69% over 2 steps). ^1H NMR and mass spectrum were consistent with the desired compound.


Part 21: Preparation of



To a solution of the amine from part 20 (4.5 g, 13.3 mmol) and triethylamine (3.72 mL, 26.7 mmol) in dichloromethane (12 mL) at zero degrees Celsius was added a solution of methane sulfonyl chloride (1.55
30 mL, 20.0 mmol) in dichloromethane (15 mL). After 2.5 hours at ambient temperature, the solvent was removed

in vacuo. The residue was partitioned between ethyl acetate (100 mL), water (100 mL), and 10% aqueous hydrochloric acid until pH = 4. The organic layer was washed with saturated sodium bicarbonate aqueous solution (50 mL), brine (25 mL), dried over Na₂SO₄, filtered, and concentrated *in vacuo* to give the methyl sulfonamide as a white, yellow solid (5.6 g, 100%). ¹H NMR and mass spectrum were consistent with the desired compound.

10 Part 22: Preparation of




In dry equipment under nitrogen, the methyl sulfonamide from part 21 (5.62 g, 13.5 mmol) was dissolved in dry tetrahydrofuran (27 mL) and cooled to minus seventy-five degrees Celsius. The resulting solution was then treated with a 1M solution of lithium bis(trimethylsilyl)amide (40.6 mL, 41.0 mmol) at a rate such that the temperature remained below minus sixty five degrees. After 30 minutes, the reaction was treated with a solution of methyl chloroformate (1.05 mL, 13.5 mmol) in dry tetrahydrofuran (13 mL), while maintaining the temperature below minus seventy degrees. After 3 hours at minus sixty degrees Celsius, the reaction was quenched with saturated ammonium chloride aqueous solution (100 mL), warmed to ambient temperature and concentrated *in vacuo*. The resulting residue was partitioned between ethyl acetate (150 mL) and water (100 mL).

30 The organic layer was washed with water (50 mL), brine (50 mL), dried over Na₂SO₄, filtered, and

-463-


concentrated *in vacuo* to give the methylene sulfonamide as a yellow oil (6.32 g, 98.6%). ¹H NMR and mass spectrum were consistent with the desired compound.

5 Part 23: Preparation of



To a solution of the methylene sulfonamide from part 22 (2.56 g, 5.41 mmol) in dimethylformamide (14 mL) was added bis-(2-bromoethyl)ether (1.38 g, 5.95 mmol), 18-Crown-6 (250 mg), followed by potassium carbonate (2.24 g, 16.2 mmol). After the heterogeneous mixture was stirred at sixty degrees Celsius for 22 hours, more potassium carbonate (0.75 g, 5.4 mmol) was added to the reaction. After another 17 hours, the reaction was charged with more potassium carbonate (0.75 g, 5.4 mmol). The reaction was concentrated *in vacuo* after 30 hours, and the resulting oil was partitioned between ethyl acetate (150 mL) and water (150 mL). The aqueous layer was extracted twice with ethyl acetate (2 x 50 mL), then the combined organic layers were washed with water (50 mL), brine (25 mL), dried over Na₂SO₄, filtered, and concentrated *in vacuo* to give the pyran methyl ester as a white solid. (2.84 g, 97%). ¹H NMR and mass spectrum were consistent with the desired compound.


 Part 24: Preparation of



-464-

In dry equipment under nitrogen, the pyran methyl ester from part 23 (2.55 g, 4.69 mmol) was suspended in dry tetrahydrofuran (10 mL) and potassium trimethylsilanolate (1.81 g, 14.1 mmol) was added at ambient temperature. After 28 hours, the reaction was charged with more potassium trimethylsilanolate (0.3 g, 2.3 mmol). After 21 hours, the reaction was again charged with more potassium trimethylsilanolate (0.3 g, 2.3 mmol). After 3 hours, the reaction was concentrated in vacuo, and the resulting residue was partitioned between ethyl acetate (50 mL) and water (50 mL). The aqueous layer was extracted twice with ethyl acetate (2 x 25 mL), then the combined organic layers were washed with brine (30 mL), dried over Na₂SO₄, filtered, concentrated and dried in vacuo to give the carboxylic acid as a tan solid (1.71 g, 69%). ¹H NMR and mass spectrum were consistent with the desired compound.


Part 25: Preparation of



In dry equipment under nitrogen, the carboxylic acid from part 24 (1.44 g, 2.72 mmol) was dissolved in dry dimethylformamide (14 mL) and treated with 1-(3-dimethylamino-propyl)-3-ethylcarbodiimide hydrochloride (0.78 g, 4.1 mmol), N-hydroxybenzotriazole hydrate (0.55 g, 4.1 mmol), O-(tetrahydro-2H-pyran-2-yl)hydroxylamine (0.48 g, 4.1 mmol), followed by N-methylmorpholine (0.90 mL, 8.16 mmol). After 26 hours at ambient temperature, the reaction

was concentrated *in vacuo*, and the resulting residue was partitioned between ethyl acetate (50 mL) and water (50 mL). The aqueous layer was extracted with ethyl acetate (20 mL), and the combined organic
5 layers were washed with 5% KHSO₄ aqueous solution (30 mL), brine (30 mL), dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The resulting yellow oil was purified on silica gel (1:1, hexanes:ethyl acetate) to give the THP hydroxamate as a yellow oil (0.96 g,
10 56%). ¹H NMR and mass spectrum were consistent with the desired compound.


Part 26: Preparation of



The THP hydroxamate from part 25 (0.82 g, 1.3
15 mmol) was dissolved in acetonitrile (10 mL) and then treated with aqueous 10% aqueous hydrochloric acid solution (10 mL, 12 mmol). After 25 hours at ambient temperature, the acetonitrile and excess hydrochloric acid were removed with a stream of nitrogen. The
20 resulting white suspension was filtered under nitrogen, washed with water and dried *in vacuo* to give the title compound as a white solid (0.39 g, 55%). HRMS *m/z* calculated for C₂₄H₃₅N₃O₇S: 545.1569, observed 545.1586.

25

Example 32: Preparation of tetrahydro-N-hydroxy-4-
[[4-(4-pentylphenyl)-1-piperazinyl]-
sulfonyl]-2H-pyran-4-carboxamide,
monohydrochloride



Part A: To a solution of tert-butyl-piperazine (5.00 g, 26.84 mmol) in toluene (50 mL) was added sodium tert-butoxide (3.01 g, 31.32 mmol). After
5 stirring at ambient temperature for 5 minutes, 1-bromo-4-n-pentylbenzene (5.08 g, 22.37 mmol), 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (BINAP) (0.418 g, 0.671 mmol) and tris(dibenzylideneacetone)-
dipalladium (0) (0.205 g, 0.224 mmol) were added to
10 the reaction mixture. The resulting mixture was heated to eighty degrees Celsius for 22 hours. After cooling to ambient temperature the reaction mixture was filtered through a pad of Celite®, washing with tetrahydrofuran and methanol. The filtrate was
15 concentrated *in vacuo* to give the aryl Boc-piperazine as an orange oily solid (9.60 g, >100%).

Part B: The aryl Boc-piperazine of part A (9.60 g, ~ 22.37 mmol) was treated with a solution of 4N HCl in dioxane (56 mL). The resulting mixture was
20 stirred at ambient temperature for 2 hours then the reaction was concentrated *in vacuo*. Diethyl ether (50 mL) was added and the precipitate was collected by filtration to give the aryl piperazine as a tan solid (8.49 g, 100 %).

25 Part C: To a solution of the aryl piperazine of part B (4.00 g, 13.10 mmol) in dichloromethane (50 mL), cooled to zero degrees Celsius, was added

triethylamine (5.48 mL, 39.30 mmol) followed by methanesulfonyl chloride (1.22 mL, 15.72 mmol). Once the addition was complete the cooling bath was removed and the resulting mixture was stirred for 2 hours. The reaction mixture was then concentrated in vacuo and the residue was partitioned between H₂O and ethyl acetate. An emulsion formed and the solids were collected by filtration. The solids were triturated with ethyl acetate to give the methyl sulfonamide as a tan solid (3.99 g, 98 %).

Part D: To a suspension of the methyl sulfonamide of part C (3.35 g, 10.79 mmol) in tetrahydrofuran (50 mL), cooled to minus seventy-eight degrees Celsius, was added lithium bis(trimethylsilyl)amide (24.00 mL, 1.0 M in tetrahydrofuran, 24.00 mmol) at such a rate that the temperature of the reaction mixture never exceeded minus seventy degrees Celsius. Once the addition was complete, the cooling bath was removed. After 30 minutes, the cooling bath was replaced and a solution of dimethyl carbonate (1.09 mL, 12.95 mmol) in tetrahydrofuran (5.0 mL) was added. After 30 minutes additional lithium bis(trimethylsilyl)amide (5.40 mL, 1.0 M in tetrahydrofuran, 5.40 mmol) was added followed by additional dimethyl carbonate (0.273 mL, 3.24 mmol). After stirring at minus seventy-eight degrees Celsius for 1 hour the reaction was quenched by the addition saturated NH₄Cl and concentrated in vacuo. The residue was diluted with H₂O and extracted with ethyl acetate. The combined organic layers were washed with 5% KHSO₄, saturated NaCl and dried over Na₂SO₄. Chromatography (on silica, ethyl

-468-

WO 00/46221

acetate/hexanes) provided the sulfonamide ester as a tan solid (1.20 g, 30%).

Part E: To a solution of the sulfonamide ester of part D (1.20 g, 3.26 mmol) in N,N-dimethylformamide (10 mL) was added K_2CO_3 (1.35 g, 9.78 mmol) and dibromoethyl ether (0.430 mL, 3.42 mmol) and the resulting mixture was heated to forty degrees Celsius. After 21 hours additional K_2CO_3 (0.450 g, 3.26 mmol) were added and the resulting mixture was heated to forty degrees Celsius for 4 hours. After cooling to ambient temperature the reaction was diluted with H_2O and extracted with ethyl acetate. The combined organic layers were washed with saturated NaCl and dried over Na_2SO_4 . Chromatography (on silica, ethyl acetate/hexanes) provided the ester as a pale yellow solid (1.23 g, 86%).

Part F: To a solution of the ester of part E (1.23 g, 2.80 mmol) in tetrahydrofuran (10 mL) was added potassium trimethylsilanolate (0.718 g, 5.60 mmol). The resulting mixture was stirred at ambient temperature overnight (about 18 hours) and then diluted with H_2O and acidified (pH=7.0) with 1 N HCl. The precipitate was collected by filtration. The solids were suspended in acetonitrile and then concentrated in vacuo to give the acid as an off-white solid (0.910 g, 76%).


Part G: To a suspension of the acid of part F (0.910 g, 2.14 mmol) in N,N-dimethylformamide (10 mL) was added 1-hydroxybenzotriazole (0.347 g, 2.57 mmol), N-methylmorpholine (0.701 mL, 6.42 mmol), O-(tetrahydropuranyl) hydroxylamine (0.752 g, 6.42 mmol) and 1-3-[(dimethylamino)propyl]-3-

ethylcarbodiimide hydrochloride (0.574 g, 3.00 mmol). The resulting mixture was stirred at ambient temperature for 22 hours. Then the reaction mixture was diluted with H₂O and extracted with ethyl acetate.

5 The combined organic layers were washed with H₂O, saturated NaHCO₃, saturated NaCl and dried over Na₂SO₄. Chromatography (on silica, ethyl acetate/hexanes) provided the protected hydroxamate as a clear oil (1.11 g, 99%).

10 Part H: The protected hydroxamate of part G (1.10 g, 2.10 mmol) was treated with a solution of 4N HCl in dioxane (5.25 mL) and methanol (0.851 mL, 21.00 mmol) and the resulting mixture was stirred at ambient temperature for 2 hours. Diethyl ether (20
15 mL) was added and the precipitate was collected by filtration to provide the title compound as an off-white solid (0.972 g, 97%). MS MH⁺ calculated for C₂₁H₃₄O₅N₃S: 440, found 440.

20 Example 33: Preparation of tetrahydro-N-hydroxy-4-[(4-phenyl-1-piperazinyl)sulfonyl]-2H-pyran-4-carboxamide



Part A: To a solution of bis(2-bromoethyl)ether
25 (1.55 g, 6.70 mmol) in acetone (30 mL) was added K₂CO₃ (9.26 g, 67.00 mmol), 18-crown-6 (500 mg) and the sulfonamide ester of part B for SC-81434A (2.00 g, 6.70 mmol). The reaction mixture was heated at reflux for 15 hours then filtered through a pad of

Celite® and the filtrate was concentrated *in vacuo*. The residue was dissolved in N,N-dimethylformamide (30 mL) and treated with K₂CO₃ (9.26 g, 67.00 mmol), 18-crown-6 (500 mg) and bis(2-bromoethyl)ether (1.55 g, 6.70 mmol) and the resulting mixture was heated at sixty degrees Celsius for 6 hours. The reaction mixture was concentrated *in vacuo* and the residue was partitioned between H₂O and chloroform. The aqueous layer was further extracted with chloroform. The combined organic layers were washed with H₂O and dried over Na₂SO₄. Chromatography (on silica, ethyl acetate with 5% acetonitrile/hexanes) provided the cyclized ester as a pale yellow oil (1.34 g, 54%).

Part B: To a solution of the cyclized ester of part A (1.34 g, 3.64 mmol) in tetrahydrofuran (15 mL) was added potassium trimethylsilanolate (1.40 g, 10.92 mmol). The resulting mixture was stirred at ambient temperature for 24 hours and then the tetrahydrofuran was removed by blowing N₂ over the reaction mixture. The residue was dissolved in H₂O, washed with diethyl ether, then acidified (pH-3.0) with 1N HCl and extracted with ethyl acetate. The combined organic layers were washed with saturated NaCl and dried over Na₂SO₄. Concentration *in vacuo* provided the acid as an off-white solid (1.20 g, 79%).


Part C: To a solution of the acid of part B (1.00 g, 2.82 mmol) in N,N-dimethylformamide (6.0 mL) was added 1-hydroxybenzotriazole (0.457 g, 3.38 mmol), N-methylmorpholine (0.924 mL, 8.46 mmol), O-(tetrahydropuranyl) hydroxylamine (0.469 g, 4.23 mmol) and 1-3-[(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (0.757 g, 3.95 mmol).

-471-

The resulting mixture was stirred at ambient temperature for 23 hours and then concentrated *in vacuo*. The residue was partitioned between ethyl acetate and H₂O. The aqueous layer was further
5 extracted with ethyl acetate. The organic layers were washed with saturated NaHCO₃, saturated NaCl and dried over Na₂SO₄. The resulting solids were washed with diethyl ether to provide the protected hydroxamate as an off-white solid. (1.08 g, 84%).
10 Part D: To a solution of the protected hydroxamate of part C (1.08 g, 2.38 mmol) in acetonitrile (8.0 mL) and H₂O (4.0 mL) was added 10% HCl (2.0 mL). After stirring at ambient temperature for 20 hours the acetonitrile was removed by blowing
15 N₂ over the reaction mixture. The aqueous reaction mixture was diluted with H₂O and extracted with ethyl acetate. The combined organic layers were washed with saturated NaCl and dried over Na₂SO₄. After concentration *in vacuo*, the solids were washed with
20 diethyl ether to give the title compound as a pale pink solid (0.503 g, 57%). MS MH⁺ calculated for C₁₆H₂₄O₅N₃S₁: 370, found 370.

Example 34: Preparation of N-hydroxy-1-(2-
25 methoxyethyl)-4-[[4-[[4-(trifluoromethyl)benzoyl]amino]-1-piperidinyl]sulfonyl]-4-
piperidinecarboxamide, monohydrochloride

-472-



Part A: To a solution of 4-amino-1-benzylpiperidine (10.0 g, 45.82 mmol) in tetrahydrofuran (100 mL) was added di-tert-butyl dicarbonate (30.0 g, 137.46 mmol) and a catalytic amount of N,N-dimethylaminopyridine. The resulting mixture was heated at reflux for 5 hours. After cooling to ambient temperature the reaction mixture was concentrated *in vacuo*. The solids were washed with hexanes to provide the carbamate as a white crystalline solid (13.7 g, >100%).

Part B: To a solution of the carbamate of part A (10.0 g, 34.43 mmol) in methanol (200 mL) was added ammonium formate (6.51 g, 103.29 mmol) and 4% Pd/C. The resulting mixture was heated at reflux for 1.5 hours. After cooling to ambient temperature the reaction mixture was filtered through a pad of Celite®, washing with methanol. The filtrate was concentrated *in vacuo* to provide the piperidine as an off-white solid (6.90 g, 100%).

Part C: To a solution of the piperidine of part B (6.90 g, 34.43 mmol) in dichloromethane (100 mL), cooled to zero degrees Celsius, was added triethylamine (5.28 mL, 37.87 mmol) followed by methanesulfonyl chloride (2.79 mL, 36.12 mmol). Once the addition was complete the cooling bath was removed and the resulting mixture was stirred for 15

hours. After concentration *in vacuo* the residue was partitioned between H₂O and ethyl acetate. The combined organic layers were washed with saturated NaCl and dried over Na₂SO₄. The resulting solids were
5 washed with hexanes to give the sulfonamide as an off-white solid (9.12 g, 95%).

Part D: To a solution of lithium bis(trimethylsilyl)amide (50.0 mL, 1.0M in tetrahydrofuran, 50.00 mmol), cooled to minus
10 seventy-eight degrees Celsius, was added a suspension of the sulfonamide of part C (4.49 g, 16.13 mmol) in tetrahydrofuran (40 mL). Once the addition was complete, the cooling bath was removed and replaced after 0.5 hours. To the resulting mixture was
15 quickly added methyl chloroformate (1.37 mL, 17.74 mmol). After 0.5 hours, the reaction mixture was quenched by the addition of saturated NH₄Cl and then the tetrahydrofuran was concentrated *in vacuo*. The reaction mixture was diluted with H₂O and extracted
20 with ethyl acetate. The combined organic layers were washed with saturated NaCl and dried over Na₂SO₄. Concentration *in vacuo* provided the sulfonamide ester as an off-white solid (5.29 g, 97%).

Part E: The sulfonamide ester of part D (5.27
25 g, 5.67 mmol) was treated with a solution of 4 N HCl in dioxane (40 mL). After stirring at ambient temperature for 1 hour the reaction mixture was concentrated *in vacuo*. The resulting solids were washed with hexanes to provided the amine as a tan
30 solid (4.19 g, 98%).

Part F: To a solution of the amine of part E (1.50 g, 5.50 mmol) in dichloromethane (10 mL), cooled to zero degrees Celsius, was added

-474-

triethylamine (1.61 mL, 11.55 mmol) followed by 4-(trifluoromethyl)benzoyl chloride (0.858 mL, 5.78 mmol). Once the addition was complete the cooling bath was removed and after stirring at ambient
5 temperature for 3.5 hours the reaction mixture was concentrated *in vacuo*. The solids were washed with H₂O and diethyl ether to provide the amide as a tan solid (1.79 g, 80%).

Part G: To a solution of the amide of part F
10 (1.52 g, 3.72 mmol) in N,N-dimethylformamide (10.0 mL) was added K₂CO₃ (1.54 g, 11.16 mmol), 18-crown-6 (0.50 g) and bis(2-chloroethyl)benzyl amine (0.864 g, 3.72 mmol). The resulting mixture was heated to sixty degrees Celsius of 22 hours, at which time
15 additional K₂CO₃ (0.514 g, 3.72 mmol) and bis(2-chloroethyl)benzyl amine (0.216 g, 0.93 mmol) were added. The resulting mixture was heated to sixty degrees Celsius for 14.5 hours at which time additional K₂CO₃ (0.514 g, 3.72 mmol) was added. The
20 resulting mixture was heated to sixty degrees Celsius of 14 hours, at which time additional K₂CO₃ (0.514 g, 3.72 mmol) and bis(2-chloroethyl)benzyl amine (0.216 g, 0.93 mmol Prepared by Darren Kassib) were added. The resulting mixture was heated to sixty degrees
25 Celsius for 24 hours. After cooling to ambient temperature the reaction mixture was diluted with H₂O and extracted with chloroform. The combined organic layers were washed with saturated NaCl and dried over Na₂SO₄. Chromatography (on silica, ethyl acetate with
30 5% methanol/hexanes) followed by trituration with diethyl ether provided the cyclized ester as an off-white solid (0.950 g, 45%).

Part H: To a solution of the cyclized ester of part G (0.950 g, 1.67 mmol) in methanol (10 mL) was added ammonium formate (0.317 g, 5.02 mmol) and 10% Pd/C (0.320 g). The resulting mixture was heated at reflux for 1.5 hours. After cooling to ambient temperature, the reaction mixture was filtered through a pad of Celite [®], washing with methanol. The filtrate was concentrated *in vacuo* to provide the amine as a gray solid (0.760 g, 95%).

10 Part I: To a solution of the amine of part H (0.760 g, 1.59 mmol) in N,N-dimethylformamide (5.0 mL) was added K₂CO₃ (0.330 g, 2.39 mmol) and 2-bromoethyl methyl ether (0.225 mL, 2.39 mmol). The resulting mixture was stirred at ambient temperature over the weekend then additional K₂CO₃ (0.055 g, 0.398 mmol) and 2-bromoethyl methyl ether (0.037 mL, 0.398 mmol) was added. After stirring at ambient temperature for 24 hours the reaction mixture was diluted with H₂O and extracted with ethyl acetate.

15 The combined organic layers were washed with saturated NaCl and dried over Na₂SO₄. Chromatography (on silica, ethyl acetate with 5% methanol/hexanes) provided the alkylated amine as a white solid (0.550 g, 65%).

25 Part J: To a solution of the alkylated amine of part I (0.540 g, 1.01 mmol) in tetrahydrofuran (5.0 mL) was added potassium trimethylsilanolate (0.388 g, 3.02 mmol). The resulting mixture was stirred at ambient temperature for 18 hours then the tetrahydrofuran was removed by blowing N₂ over the reaction mixture. The reaction mixture was diluted with H₂O and neutralized (pH=7) with 1N HCl. The resulting aqueous reaction mixture was concentrated

30

-476-

WO 00/46221

in vacuo to provide the crude acid as a pink solid (0.610 g, >100%).

Part K: To a solution of the crude acid of part J (0.610 g, ~ 1.01 mmol) in N,N-dimethylformamide (5.0 mL) was added 1-hydroxybenzotriazole (0.164 g, 1.21 mmol), N-methylmorpholine (0.331 mL, 3.03 mmol), O-(tetrahydropuranyl) hydroxylamine (0.177 g, 1.52 mmol) and 1-3-[(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (0.271 g, 1.41 mmol).


10 After stirring at ambient temperature for 0.5 hours, additional N,N-dimethylformamide (5.0 mL) was added. After stirring at ambient temperature overnight (about 18 hours) the reaction mixture was heated to forty-five degrees Celsius for 24 hours. The reaction mixture was then diluted with H₂O and extracted with ethyl acetate. The combined organic layers were washed with saturated NaHCO₃, saturated NaCl and dried over Na₂SO₄. The resulting solids were washed with diethyl ether to give the protected hydroxamate as an off-white solid (0.300 g, 49%).

20 Part L: To a solution of the protected hydroxamate of part K (0.300 g, 0.483 mmol) in dioxane (3.0 mL) and methanol (1.0 mL). After solution of 4N HCl in dioxane (1.2 mL). After stirring at ambient temperature for 1.5 hours the solvent was removed by blowing N₂ over the reaction mixture. The resulting solids were washed with diethyl ether to give the title compound as a pink solid (0.193 g, 70%). MS MH⁺ calculated for C₂₂H₃₂O₆N₄S₁F₃: 537, found 537.

30 Example 35: Preparation of N-hydroxy-1-phenyl-4-[[4-(4-(trifluoromethoxy)phenoxy]-1-

-477-

piperidinyl)sulfonyl]-4-

piperidinecarboxamide, monohydrochloride

5 Part A: To a solution of commercially available
4-hydroxypiperidine (46.3 g, 458 mmol) in
tetrahydrofuran (300 mL) was slowly added
triethylamine (67.0 mL, 481 mmol) followed by a
solution of di-tert-butyl dicarbonate (100 g, 458
10 mmol) in tetrahydrofuran (200 mL). After stirring at
ambient temperature for 17 hours the reaction mixture
was concentrated *in vacuo*. The residue was dissolved
in ethyl acetate (500 mL) and washed with 5% KHSO₄,
saturated NaHCO₃, saturated NaCl and dried over
15 Na₂SO₄. Crystallization with hexanes provided the
carbamate as an off-white solid (87.8 g, 95%).

 Part B: To a solution of the carbamate of part
A (5.00 g, 24.84 mmol) in dichloromethane (50 mL),
precooled in an ice-bath, was added triethylamine
20 (3.81 mL, 27.32 mmol) followed by methanesulfonyl
chloride (2.02 mL, 26.08 mmol). Once the addition
was complete, the cooling bath was removed. After
stirring for 2 hours, the reaction mixture was
concentrated *in vacuo*. The residue was partitioned
25 between ethyl acetate and H₂O. The aqueous layer was
further extracted with ethyl acetate. The combined
organic layers were washed with saturated NaCl and

dried over Na_2SO_4 . Concentration *in vacuo* provided the mesylate as an off-white solid (7.34 g, >100%).

Part C: To a solution of 4-(trifluoromethoxy)-phenol (2.00 g, 11.23 mmol) in N,N-dimethylformamide (25 mL), cooled to zero degrees Celsius, was added sodium hydride (0.449 g, 60% oil dispersion, 11.23 mmol). Once the addition was complete the cooling bath was removed and then replaced after 0.5 hours. To the resulting mixture was added the mesylate of part B (2.61 g, 9.36 mmol). The reaction mixture was then heated to forty degrees Celsius. After stirring at forty degrees Celsius for 15 hours the temperature of the reaction was increased to eighty degrees Celsius. After 8 hours at eighty degrees Celsius the reaction mixture was cooled in an ice-bath and additional sodium hydride (0.225 g, 60% oil dispersion, 5.62 mmol) was added. After 30 minutes additional mesylate of part B (1.31 g, 4.68 mmol) was added and the resulting mixture was heated to eighty degrees Celsius. After 15 hours at eighty degrees Celsius the reaction was cooled to ambient temperature and concentrated *in vacuo*. The residue was partitioned between H_2O and diethyl ether. The organic layer was washed with saturated NaCl and dried over Na_2SO_4 . After concentration *in vacuo* the residue was treated with a solution of 4 N HCl in dioxane (30 mL). After stirring at ambient temperature for 2 hours, the reaction mixture was concentrated *in vacuo*. Water was added and the reaction was extracted with ethyl acetate. The aqueous layer was then made alkaline (pH-10) with 2.5 N NaOH and extracted with ethyl acetate. The combined organic layers were washed with saturated

NaCl and dried over Na_2SO_4 . Concentration *in vacuo* provided the amine as an off-white solid (1.67 g, 68%).

Part D: To a solution of the amine of part C
5 (11.5 g, 44.1 mmol) in dichloromethane (125 mL),
precooled in an ice-bath, was added triethylamine
(12.3 mL, 88.1 mmol) followed by a solution of
methanesulfonyl chloride (5.1 mL, 66.1 mmol) in
dichloromethane (20 mL). Once the addition was
10 complete, the cooling bath was removed. After 1 hour
at ambient temperature, the reaction mixture was
concentrated *in vacuo*. The residue was partitioned
between H_2O and ethyl acetate. The organic layer was
washed with saturated NaCl and dried over Na_2SO_4 .
15 Recrystallization from ethyl acetate/hexanes provided
the sulfonamide as an off-white solid (10.77 g,
72%).

Part E: To a solution of the sulfonamide of
part D (10.77 g, 31.8 mmol) in tetrahydrofuran (64
20 mL), cooled to minus seventy-five degrees Celsius,
was added lithium bis(trimethylsilyl)amide (80 mL, 1M
in tetrahydrofuran, 80.0 mmol), at such a rate that
the temperature of the reaction never exceeded minus
sixty five degrees Celsius. After stirring at minus
25 seventy-five degrees Celsius for 30 minutes, a
solution of methyl chloroformate (2.45 mL, 31.8 mmol)
in tetrahydrofuran (32 mL) was slowly added at such a
rate that the temperature of the reaction never
exceeded minus sixty five degrees Celsius. After
30 stirring at minus seventy-five degrees Celsius for 30
minutes the reaction was quenched by the addition of
saturated NH_4Cl and extracted with ethyl acetate. The
combined organic layers were washed with saturated

-480-

WO 00/46221

NaCl and dried over Na_2SO_4 . Concentration in vacuo gave the sulfonamide ester as a yellow oil (12.69, 100%).

5 Part F: To a suspension of dibromotriphenylphosphorane (50.0 g, 118.45 mmol) in dichloromethane (200 mL), cooled to zero degrees Celsius, was added N-phenyldiethanol amine (10.0 g, 55.18 mmol). Once the addition was complete, the cooling bath was removed. After stirring at ambient temperature for 10 17.5 hours, the reaction mixture was concentrated in vacuo. The residue was then treated with warm ethyl acetate and the resulting precipitate was removed and the filtrate was concentrated in vacuo.

15 Chromatography (on silica, hexanes/dichloromethane) provided the dibromoamine as a light yellow oil (5.85 g, 35%).

Part G: To a solution of the dibromoamine of part F (5.60 g, 18.24 mmol) in N,N-dimethylformamide were added K_2CO_3 (6.87 g, 49.74 mmol), the sulfonamide ester of part E (6.59 g, 16.58 mmol) and 18-crown-6 (1.66 g). The resulting mixture was heated to eighty 20 degrees Celsius for 14.5 hours at which time additional K_2CO_3 (3.44 g, 24.89 mmol) was added. After stirring at eighty degrees Celsius for 7.5 25 hours the reaction mixture was cooled to ambient temperature and then concentrated in vacuo. The residue was dissolved in ethyl acetate, filtered through a pad of Celite[®] and the filtrate was concentrated in vacuo. Chromatography (on silica, 30 ethyl acetate/hexanes) gave a yellow solid that after washing with hot methanol provided the cyclized ester as an off-white solid (3.15 g, 35%).


Part H: To a solution of the cyclized ester of part G (3.15 g, 5.81 mmol) in tetrahydrofuran (25 mL) was added potassium trimethylsilanolate (2.24 g, 17.43 mmol). The resulting mixture was stirred at ambient temperature for 24 hours and then H₂O was added. After neutralization (pH=7) with 1 N HCl, the tetrahydrofuran was removed by concentration in vacuo. After readjusting the pH of the aqueous reaction mixture (pH = 7) the precipitate was collected by filtration of give the acid as an off-white solid (2.97 g, 97%).

Part I: To a solution of the acid of part H (2.97 g, 5.62 mmol) in N,N-dimethylformamide (20 mL) were added 1-hydroxybenzotriazole (0.911 g, 6.74 mmol), N-methylmorpholine (1.85 mL, 16.86 mmol), O-(tetrahydropuranyl) hydroxylamine (1.98 g, 16.86 mmol) and 1-3-[(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (1.51 g, 7.87 mmol). After stirring at ambient temperature for 18 hours the reaction mixture was concentrated in vacuo. Water was added and the aqueous reaction mixture was extracted with ethyl acetate. The combined organic layers were washed with saturated NaHCO₃, saturated NaCl and dried over Na₂SO₄. The resulting solids were washed with hot methanol to give the protected hydroxamate as a white solid (3.09 g, 88%).

Part J: The protected hydroxamate of part I (3.09 g, 4.92 mmol) was treated with a solution of 4 N HCl in dioxane (12.0 mL) and methanol (1.99 mL, 49.23 mmol). After stirring at ambient temperature for several minutes additional dioxane (10 mL) was added. After stirring at ambient temperature for 1.5 hours diethyl ether was added and the precipitate was

collected by filtration to give the title compound as an off-white solid (2.67 g, 94%). MS MH^+ calculated for $C_{24}H_{29}O_6N_3S_1F_3$: 544, found 544.

- 5 Example 36: Preparation of N-hydroxy-1-phenyl-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-4-
piperidinecarboxamide, monohydrochloride



10

Part A: To a solution of the carbamate of part A for Example 35 (6.03 g, 30.0 mmol) in N,N-dimethylformamide (60 mL), were added Cs_2CO_3 (9.77 g, 30.0 mmol) and 4-fluorobenzotrifluoride (3.8 mL, 30.0 mmol). After stirring at ninety degrees Celsius for 19 hours, additional Cs_2CO_3 (3.26 g, 10.0 mmol) and 4-fluorobenzotrifluoride (0.95 mL, 10.0 mmol) were added. After stirring at ninety degrees Celsius for 46 hours, the reaction mixture was concentrated *in vacuo*. The residue was dissolved in ethyl acetate and the organic layer was washed with H_2O , saturated NaCl and dried over Na_2SO_4 . Chromatography (on silica, ethyl acetate/hexanes) provided the Boc-amine as a white solid (6.0 g, 58%).

Part B: The Boc-amine of part A (4.10 g, 11.87 mmol) was treated with a solution of 4N HCl in dioxane (20.0 mL). After stirring at ambient

temperature for 1.5 hours, the reaction mixture was concentrated in vacuo to provide the amine as a white solid (3.54 g, >100%).

Part C: To a solution of the amine of part B
5 (3.34 g, 11.87 mmol) in dichloromethane (40 mL), cooled in an ice-bath, was added triethylamine (3.31 mL, 23.74 mmol) followed by methanesulfonyl chloride (1.38 mL, 17.81 mmol). Once the addition was complete the cooling bath was removed. After
10 stirring at ambient temperature for 2 hours, the reaction mixture was concentrated in vacuo. The residue was partitioned between H₂O and ethyl acetate. The organic layer was washed with 5% KHSO₄, saturated NaCl and dried over Na₂SO₄. Concentration in vacuo
15 provided the sulfonamide as a pale yellow solid (4.25 g, >100%).

Part D: To a solution of the sulfonamide of part C (5.00 g, 15.46 mmol) in tetrahydrofuran (30 mL), cooled to minus forty degrees Celsius, was added
20 lithium bis(trimethylsilyl)amide (31.0 mL, 1 M in tetrahydrofuran, 31.0 mmol) at such a rate that the temperature of the reaction never exceeded minus thirty five degrees Celsius. After stirring at minus forty degrees Celsius for 30 minutes, a solution of
25 dimethyl carbonate (1.56 mL, 18.55 mmol) in tetrahydrofuran (10 mL) was added at such a rate that the temperature of the reaction never exceeded minus thirty five degrees Celsius. After stirring at minus forty degrees Celsius for 30 minutes, the reaction
30 was quenched by the addition of saturated NH₄Cl. The resulting mixture was slowly permitted to warm to ambient temperature and then the tetrahydrofuran was removed in vacuo. The aqueous reaction mixture was

diluted with H₂O and extracted with ethyl acetate and diethyl ether. The combined organic layers were washed with 5% KHSO₄, saturated NaCl and dried over Na₂SO₄. Concentration in vacuo provided the
5 sulfonamide ester as a thick yellow oil (5.75 g, 97%).

Part E: To a solution of the dibromoamine of part F of Example 35 (7.00 g, 22.80 mmol) in N,N-dimethylformamide (45 mL), was added K₂CO₃ (9.45 g,
10 68.40 mmol), the sulfonamide ester of part D (8.70 g, 22.80 mmol) and 18-crown-6 (2.28 g). The resulting mixture was heated to eighty degrees Celsius for 15 hours and then additional K₂CO₃ (4.73 g, 34.22 mmol) was added. The resulting mixture was then heated to
15 eighty degrees Celsius for 6 hours. After cooling to ambient temperature, the reaction mixture was diluted with H₂O and extracted with ethyl acetate. The combined organic layers were washed with saturated NaCl and dried over Na₂SO₄. Chromatography (on
20 silica, ethyl acetate/hexanes) followed by washing of the resulting solids with boiling methanol provided the cyclized ester as an off-white solid (4.50 g, 37%).

Part F: To a solution of the cyclized ester of
25 part E (4.50 g, 8.55 mmol) in tetrahydrofuran (40 mL) was added potassium trimethylsilanolate (3.29 g, 25.64 mmol). The resulting mixture was stirred at ambient temperature for 22 hours. The reaction mixture was diluted with H₂O and neutralized (pH-7)
30 with 1N HCl. The tetrahydrofuran was removed in vacuo and the precipitate was collected by filtration. The solids were suspended in

-485-


acetonitrile and concentrated *in vacuo* to provide the acid as a white solid (4.05 g, 92%).

Part G: To a solution of the acid of part F (4.05 g, 7.90 mmol) in N,N-dimethylformamide were
5 added 1-hydroxybenzotriazole (1.28 g, 9.48 mmol), N-methylmorpholine (2.59 mL, 23.70 mmol), O-(tetrahydropuranyl) hydroxylamine (2.78 g, 23.70 mmol) and 1-3-[(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (2.12 g, 11.06 mmol).
10 After stirring at ambient temperature for 16 hours the reaction mixture was diluted with H₂O and extracted with ethyl acetate. The combined organic layers were washed with saturated NaHCO₃, saturated NaCl and dried over Na₂SO₄. The solids were washed with boiling
15 methanol to provide the protected hydroxamate as a white solid (4.12 g, 85%).

Part H: The protected hydroxamate of part G (4.12 g, 6.74 mmol) was treated with a solution of 4N HCl in dioxane (12.0 mL) and methanol (1.99 mL, 49.23
20 mmol). After stirring at ambient temperature for several minutes, additional dioxane (10 mL) was added. After stirring at ambient temperature for 1.5 hours, diethyl ether was added and the precipitate was collected by filtration to give the title
25 compound as an off-white solid (3.32 g, 87%). MS MH⁺ calculated for C₂₄H₂₉O₅N₃S₁F₃: 528, found 528.

Example 37: Preparation of 4-[[4-[4-(1,1-dimethylethyl)phenyl]-1-piperazinyl]-sulfonyl]-N-hydroxy-1-(2-methoxyethyl)-4-piperidinecarboxamide,
30 monohydrochloride

-486-



Part A: To a suspension of 4-bromopiperidine hydrobromide (30 g, 122.5 mmol) and N-

5 (benzyloxycarbonyloxy)succinimide in tetrahydrofuran (250 mL) was added N-methylmorpholine (15 g, 148.3 mmol) followed by a catalytic amount of N,N-dimethylaminopyridine. After stirring at ambient temperature for 17 hours, the reaction mixture was

10 concentrated in vacuo. The residue was partitioned between ethyl acetate and H₂O. The organic layer was washed with 10% HCl solution, saturated NaCl and dried over Na₂SO₄. Concentration in vacuo provided the carbamate as a clear oil (33.0 g, 90%).

15 Part B: To a solution of the carbamate of part A (30.4 g, 101.96 mmol) in N,N-dimethylformamide (200 mL) was added potassium thioacetate (12.75 g, 111.64 mmol) and the resulting mixture was stirred at ambient temperature. After 23 hours, the reaction

20 mixture was heated to sixty degrees Celsius. After heating at sixty degrees Celsius for 3 hour the reaction mixture was concentrated in vacuo. The residue was dissolved in ethyl acetate and filtered through a pad of Celite®. The filtrate was then

25 concentrated in vacuo. The residue was redissolved in ethyl acetate and the organic layer was washed with saturated NaHCO₃, saturated NaCl and dried over

Na₂SO₄. Chromatography (on silica, ethyl acetate/hexanes) provided the thioacetate as an orange oil (22.8 g, 76%).

Part C: Into a solution of the thioacetate of part B (22.8 g, 77.71 mmol) in carbon tetrachloride (240 mL) and ethanol (60 mL) was bubbled Cl₂. Once the exotherm was complete the bubbling of Cl₂ was discontinued. N₂ was bubbled through the reaction mixture for several minutes then the reaction was concentrated *in vacuo* to give the sulfonyl chloride as a brown oil (25.5 g, >100%).

Part D: To a solution of tert-butyl piperazine (5.24 g, 28.15 mmol) in toluene (50 mL) was added sodium tert-butoxide (3.16 g, 32.84 mmol). The resulting mixture was stirred at ambient temperature for several minutes and then 1-bromo-4-tert-butylbenzene (5.00 g, 23.46 mmol), BINAP (0.438 g, 0.704 mmol) and tris(dibenzylideneacetone)dipalladium (0) (0.215 g, 0.235 mmol) were added. The resulting mixture was heated to eighty degrees Celsius for 16 hours. After cooling to ambient temperature, diethyl ether was added and the reaction mixture was filtered through a pad of Celite® and the filtrate was concentrated *in vacuo*. The residue was dissolved in diethyl ether and washed with H₂O, 5% KHSO₄, saturated NaCl and dried over Na₂SO₄. Concentration *in vacuo* provided the phenyl-piperazine as a brown solid (7.80 g, >100%).

Part E: The phenyl-piperazine of part D (7.80 g, 24.49 mmol) was treated with a solution of 4N HCl in dioxane (61 mL). The resulting reaction mixture was stirred at ambient temperature for 1 hour and then concentrated *in vacuo*. The solids were washed

-488-

with diethyl ether to give the amine hydrochloride as a mustard colored solid (6.10 g, 98%).

Part F: To a suspension of the amine hydrochloride of part D (1.50 g, 5.89 mmol) and
5 triethylamine (1.81 mL, 12.96 mmol) in dichloromethane (10 mL), cooled in an ice-bath, was added the sulfonyl chloride of part C (1.82 g, 5.73 mmol). Once the addition was complete the cooling bath was removed. After stirring at ambient
10 temperature for 2 hours the reaction mixture was concentrated in vacuo. The residue was partitioned between H₂O and ethyl acetate and the aqueous layer was further extracted with ethyl acetate. The combined organic layers were washed with 5% KHSO₄,
15 saturated NaCl and dried over Na₂SO₄. The resulting solids were triturated with methanol to provide the sulfonamide as a light brown solid (1.41 g, 49%).

Part G: To a solution of the sulfonamide of part F (1.40 g, 2.80 mmol) in tetrahydrofuran (10
20 mL), at ambient temperature, was added lithium bis(trimethylsilyl)amide (6.20 mL, 1 M in tetrahydrofuran, 6.20 mmol). After stirring at ambient temperature for 1 hour, dimethyl carbonate (0.283 mL, 3.36 mmol) was added. The resulting
25 mixture was stirred at ambient temperature for 3 hours and then additional lithium bis(trimethylsilyl)amide (3.10 mL, 1 M in tetrahydrofuran, 3.10 mmol) was added. After 2 hours at ambient temperature additional dimethyl carbonate
30 (0.140 mL, 1.66 mmol) was added. After stirring at ambient temperature overnight (about 18 hours), the reaction was quenched by the addition of saturated NH₄Cl. The reaction was diluted with H₂O and

extracted with ethyl acetate. The combined organic layers were washed with 5% KHSO_4 , saturated NaCl and dried over Na_2SO_4 . The resulting solids were triturated with methanol to give the sulfonamide ester as a tan solid (1.29 g, 83%).

Part H: To a solution of the sulfonamide ester of part G (1.29 g, 2.31 mmol) in tetrahydrofuran (20 mL) and 10% Pd/C (0.300 g) was bubbled H_2 . After bubbling H_2 through the reaction for 17 hours, the mixture was purged with N_2 and filtered through a pad of Celite®, washing with tetrahydrofuran. The filtrate was concentrated *in vacuo* to give the amine as a dark brown sticky solid (0.950 g, 97%).

Part I: To a solution of the amine of Part H (0.940 g, 2.22 mmol) in *N,N*-dimethylformamide (7.0 mL) was added K_2CO_3 (0.614 g, 4.44 mmol) and 2-bromoethyl methyl ether (0.313 mL, 3.33 mmol). The resulting mixture was stirred at ambient temperature for 3 days. The reaction mixture was concentrated *in vacuo* and the residue was dissolved in ethyl acetate and filtered through a pad of Celite®. Chromatography (on silica, ethyl acetate/hexanes) provided the alkylated amine as an off-white solid (0.595 g, 56%).


Part J: To a solution of the alkylated amine of part I (0.595 g, 1.24 mmol) in tetrahydrofuran (8.0 mL) was added potassium trimethylsilanolate (0.318 g, 2.48 mmol). After stirring at ambient temperature for 17 hours the tetrahydrofuran was removed by blowing N_2 over the reaction mixture. The residue was dissolved in H_2O and the aqueous reaction mixture was neutralized ($\text{pH} = 7$) with 1 N HCl . The precipitate

was collected by filtration to provide the acid as an off-white solid (0.475 g, 82%).

Part K: To a solution of the acid part J (0.475 g, 1.02 mmol) in N,N-dimethylformamide (5.0 mL) was
5 added 1-hydroxybenzotriazole (0.165 g, 1.22 mmol), N-methylmorpholine (0.334 mL, 3.06 mmol), O-(tetrahydropuranyl) hydroxylamine (0.359 g, 3.06 mmol) and 1-3-[(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (0.274 g, 1.43 mmol).
10 The resulting reaction mixture was stirred at ambient temperature for 17.5 hours and then heated to sixty degrees Celsius for 6 hours. To the reaction mixture was added triethylamine (0.427 mL, 3.06 mmol) and the resulting mixture was heated to sixty degrees Celsius
15 for 21 hours. After cooling to ambient temperature, the reaction mixture was concentrated *in vacuo*. The residue was suspended in H₂O, acidified (pH = 1) with 1 N HCl and then concentrated *in vacuo*. To a solution of the resulting solids in N,N-
20 dimethylformamide (5.0 mL) were added 1-hydroxybenzotriazole (0.119 g, 0.881 mmol) and 1-3-[(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (0.197 g, 1.03 mmol). After stirring at ambient temperature for 1 hour, N-methylmorpholine
25 (0.400 mL, 3.67 mmol), O-(tetrahydropuranyl) hydroxylamine (0.258 g, 2.20 mmol) were added. After stirring at ambient temperature for 2 days the reaction mixture was diluted with H₂O, then neutralized (pH = 7) and extracted with ethyl
30 acetate. The combined organic layers were washed with saturated NaHCO₃, saturated NaCl and dried over Na₂SO₄. The crude protected hydroxamate was treated with a solution of 4 N HCl in dioxane (1.8 mL) and

methanol (0.309 mL). After stirring at ambient temperature for 1.5 hours, diethyl ether was added and the precipitate was collected by filtration. The solids were washed with acetonitrile to give the
5 title compound as a tan solid (0.188 g, 49%). MS MH^+ calculated for $C_{23}H_{39}O_5N_4S_1$: 483, found 483.

10 Example 38: Preparation of 4-[[4-(4-butoxyphenyl)-1-piperazinyl]sulfonyl]-N-hydroxy-1-(2-methoxyethyl)-4-piperidinecarboxamide, dihydrochloride



Part A: To a solution of n-butyloxybromobenzene (5.00 g, 21.82 mmol) in toluene (50 mL) was added
15 tert-butyl piperazine (4.88 g, 26.18 mmol) and sodium tert-butoxide (2.94 g, 30.55 mmol). After stirring at ambient temperature for several minutes, BINAP (0.408 g, 0.655 mmol) and tris(dibenzylideneacetone)-
dipalladium (0) (0.200 g, 0.218 mmol) were added.
20 The resulting mixture was heated to eighty degrees Celsius for 21 hours. After cooling to ambient temperature, the reaction mixture was filtered through a pad of Celite®, washing with diethyl ether and dichloromethane. Chromatography (on silica,
25 ethyl acetate/hexanes) provided the phenyl-piperazine as a tan solid (5.56 g, 76%).

-492-

WO 00/46221

Part B: The phenyl-piperazine of part A (5.56 g, 16.62 mmol) was treated with a solution of 4 N HCl in dioxane (42 mL). The resulting mixture was stirred at ambient temperature for 1.5 hours and then concentrated in vacuo. The resulting solids were washed with diethyl ether to provide the amine hydrochloride as an off-white solid (4.60 g, >100%).

Part C: To a solution of the amine hydrochloride of part B (2.24 g, 8.26 mmol) and triethylamine (2.41 mL, 17.31 mmol) in dichloromethane, cooled to zero degrees Celsius, was slowly added a solution of the sulfonyl chloride of part C of Example 37 (2.50 g, 7.87 mmol) in dichloromethane (20 mL). Once the addition was complete, the cooling bath was removed. After stirring at ambient temperature for 4 hours, additional triethylamine (1.25 mL, 8.97 mmol) was added. The resulting mixture was stirred at ambient temperature for 2.5 hours and then concentrated in vacuo. The residue was partitioned between 5% KHSO₄ solution and ethyl acetate. The organic layer was washed with saturated NaCl and dried over Na₂SO₄. Chromatography (on silica, ethyl acetate/hexanes) provided the sulfonamide as an off-white solid (3.21 g, 75%).

Part D: To a solution of the sulfonamide (3.18 g, 6.17 mmol) in tetrahydrofuran (30 mL), cooled to zero degrees Celsius, was added lithium bis(trimethylsilyl)amide (13.6 mL, 1 M in tetrahydrofuran, 13.6 mmol). Once the addition was complete, the cooling bath was removed and replaced after 1 hour. To the resulting mixture was added dimethyl carbonate (0.624 mL, 7.40 mmol). Once the

addition was complete the cooling bath was removed. After stirring at ambient temperature overnight (about 18 hours), additional dimethyl carbonate (0.260 mL, 3.09 mmol) was added. After stirring at
5 ambient temperature for 2 hours, additional lithium bis(trimethylsilyl)amide (3.09 mL, 1 M in tetrahydrofuran, 3.09 mmol) was added. After stirring at ambient temperature for 2 hours, the reaction mixture was cooled to zero degrees Celsius,
10 and quenched by the addition of saturated NH_4Cl . The tetrahydrofuran was concentrated *in vacuo* and the aqueous reaction mixture was diluted with H_2O and extracted with diethyl ether. The combined organic layers were washed with 5% KHSO_4 , saturated NaCl and
15 dried over Na_2SO_4 . Chromatography (on silica, ethyl acetate/hexanes) provided the sulfonamide ester as a pale yellow gum (1.76 g, 50%).

Part E: To a suspension of the sulfonamide ester of part D (1.76 g, 3.07 mmol) and 10% Pd/C
20 (0.307 g) in tetrahydrofuran (25 mL) was bubbled H_2 . After bubbling H_2 through the reaction mixture for 23 hours the mixture was purged with N_2 and filtered through a pad of Celite®, washing with methanol. The filtrate was concentrated *in vacuo* to give the amine
25 as a pale yellow gum (1.19 g, 88%).

Part F: To a solution of the amine of part E (1.19 g, 2.71 mmol) in N,N-dimethylformamide (10 mL) were added K_2CO_3 (0.749 g, 5.42 mmol) and 2-bromoethyl methyl ether (0.382 mmol, 4.07 mmol). The resulting
30 mixture was stirred at ambient temperature for 21 hours. The reaction mixture was concentrated *in vacuo* and the residue was dissolved in ethyl acetate and filtered through a pad of Celite®.

Chromatography (on silica, ethyl acetate with 5% methanol/hexanes) provided the alkylated amine as a tan solid (0.770 g, 57%).


Part G: To a solution of the alkylated amine of
5 part F (0.770 g, 1.55 mmol) in tetrahydrofuran (5.0 mL) was added potassium trimethylsilanolate (0.398 g, 3.10 mmol). After stirring at ambient temperature for 3 hours additional tetrahydrofuran (5.0 mL) was added. The resulting reaction mixture was stirred at
10 ambient temperature for 24 hours and then the tetrahydrofuran was removed by blowing N₂ over the reaction mixture. To the residue was added acetonitrile (20 mL) and 1 N HCl (5 mL). The resulting mixture was stirred at ambient temperature
15 for several minutes and then concentrated *in vacuo* to provide the crude acid as a tan solid (0.806 g, >100%).

Part H: To a suspension of the crude acid of part G (0.806 g, ~1.55 mmol) in N,N-dimethylformamide
20 (7.0 mL) were added 1-hydroxybenzotriazole (0.251 g, 1.86 mmol) and 1-3-[(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (0.416 g, 2.17 mmol). After stirring at ambient temperature for 1.5 hours, N-methylmorpholine (0.847 mL, 7.75 mmol) and O-
25 (tetrahydropuranyl) hydroxylamine (0.545 g, 4.65mmol) were added. After stirring at ambient temperature overnight (about 18 hours) the reaction mixture was diluted with H₂O and extracted with ethyl acetate. The combined organic layers were washed with
30 saturated NaHCO₃, saturated NaCl and dried over Na₂SO₄. Chromatography (on silica, ethyl acetate with 5% methanol/hexanes) provided the protected hydroxamate as a yellow sticky oil (0.687 g, 76%).

Part I: The protected hydroxamate of part H (0.687 g, 1.18 mmol) was treated with a solution of 4 N HCl in dioxane (2.95 mL) and methanol (0.500 mL). The resulting mixture was stirred at ambient temperature for 1.5 hours and then diethyl ether was added. The precipitate was collected by filtration, washing with diethyl ether, to give the title compound as an off-white solid (0.530 g, 79%). MS MH⁺ calculated for C₂₃H₃₉N₄O₆S₁: 499, found 499.


10

Example 39: Preparation of N-hydroxy-4-[[1'-(2-methoxyphenyl)[4,4'-bipiperidin]-1-yl]sulfonyl]-1-(phenylmethyl)-4-piperidinecarboxamide, dihydrochloride



15

Part 1: Preparation of




To N-(t-butoxycarbonyl)-4,4'-bipiperidine (prepared as described in WO94/14776; preparation 22; (3.5g, 30 mmol) and sodium t-butoxide (Aldrich, 1.8 mL, 110 mmol), 2-bromoanisole (Aldrich; 2 g, mmol), BINAP (Aldrich; 150 mg), palladium chloride (Aldrich,

20

50 mg) were slurried in toluene (22 mL) and heated to 80°C. After the disappearance of the starting material, the solvent was removed and the residue was taken up in ethyl acetate (100 mL) and H₂O (30 mL).
5 Once separated, the organic layer was washed with 5% KHSO₄ (3x-50 mL) and brine (1x- 50 mL). The organic layer was then dried over Na₂SO₄, filtered, and concentrated to afford the N-aryl Boc bipiperidine as an oil (6.5 g). ¹H NMR and mass spectroscopy showed
10 the desired compound.


Part 2: Preparation of



To a solution of the product (6 g) of Part 1 in 1,4-dioxane (10 mL) was added 4 N HCl in dioxane (50
15 mL, 200 mmol). The mixture was stirred at room temperature until starting material was gone by LC (about 1 hour). The solvents were then removed and the residue was slurried in diethyl ether and filtered. The solid was washed with diethyl ether
20 (2x-50 mL) and dried in vacuo to afford the N-aryl bipiperidine as a white solid (6 g). ¹H NMR and mass spectroscopy showed the desired compound as the HCl salt.

Part 3: Preparation of


-497-



The HCl salt of Part 2 (5g, 14 mmol) and triethylamine (Aldrich, 4.4 mL, 42 mmol) were slurried in CH₂Cl₂ (50 mL) and cooled to zero°C. A solution of methane sulfonyl chloride (Aldrich, 2 g, 50 mmol) in CH₂Cl₂ (20 mL) was slowly added, maintaining the temperature below 10°C. After the addition, the ice bath was removed and the reaction stirred 1 hour as it warmed to ambient temperature.

After the disappearance of the starting material, the solvent was removed and the residue was taken up in ethyl acetate (100 mL) and H₂O (30 mL). Once separated, the organic layer was washed with 5% KHSO₄ (3x-50 mL) and brine (1x- 50 mL). The organic layer was then dried over Na₂SO₄, filtered, and concentrated to afford an oily solid that was recrystallized from diethyl ether, affording the N-aryl methylsulfonamide bipiperidine as an off-white solid (4 g). ¹H NMR and mass spectroscopy showed the desired compound.


Part 4: Preparation of



Oven-dried glassware was charged with the compound from Part 3 (3.5 g, 10 mmol) and

tetrahydrofuran (20 mL,) and cooled to -75°C . Lithium bis(trimethylsilyl)amide (Aldrich, 1.0 M in tetrahydrofuran, 30 mL, 33 mmol) was slowly added, keeping temperature less than -60°C . The reaction was stirred for 30 minutes after the addition, and was then charged with a solution of methyl chloroformate (Aldrich, 1.1 mmol, 11 mmol) in tetrahydrofuran (1 mL) again keeping the temperature at less than -60°C . After stirring for 1 hour at -75°C , the reaction was quenched with saturated NH_4Cl , keeping temperature less than -20°C . The aqueous portion froze into a solid chunk of ice. After warming to 5°C , the mixture was extracted via ethyl acetate (3x- 200 mL). Organics were washed with saturated NH_4Cl (2x-100 mL) and brine (1x-100mL), then dried over Na_2SO_4 and concentrated to afford the N-aryl bipiperidine methylene as a tan oil (4.0 g, 90% crude yield). ^1H NMR and mass spectroscopy indicated desired compound.

Part 5: Preparation of




20

To a solution of compound from Part 4 (3 g, 11 mmol) and bis-(2-chloroethyl)benzyl amine (2.7 mL, 15.1 mmol) in dimethylformamide (30 mL) was added 18-Crown-6 (Aldrich, 500 mg, cat.), followed by potassium carbonate (Aldrich, 5 g, 27.4 mmol). The mixture was heated at 60°C for 16 hours. The product was isolated by pouring into water(200 mL) and extraction with ethyl acetate (3x- 300 mL). Organics

were washed with brine (2x- 200 mL), dried over Na_2SO_4 , and concentrated to afford the amino ester an oil (3g) that solidified on standing. ^1H NMR and mass spectroscopy showed the desired compound.


5 Part 6: Preparation of



To a solution of the compound from Part 5 (2 g, 7 mmol) in tetrahydrofuran (20 mL) was added potassium trimethylsilanolate (Aldrich, 2. g, 18 mmol). The reaction stirred overnight (about 18 hours) at room temperature, at which time LC showed less than 3% starting material remained. Work up comprised removing the tetrahydrofuran and taking the residue up in H_2O (100 mL). The solution was washed with diethylether (50 mL). The aqueous was then cooled to zero°C and 10% HCl_{aq} was slowly added until $\text{pH} = 3$. The acidic mixture was then extracted with ethyl acetate (3x-150 mL). The organics were washed with brine (1x- 100 mL), dried over Na_2SO_4 , and concentrated to afford a wet solid. The solid was dried in vacuo with phosphorous pentoxide yielding the amino acid as an orange solid (2.4 g, 72% yield). ^1H NMR and mass spectroscopy showed the desired compound.

25 Part 7: Preparation of

-500-




To a solution of the acid product in Part 6 (3 g, 5.2 mmol) in dimethylacetamide (20 mL) was added N-methylmorpholine (Aldrich, 3.0 mL, 15 mmol) followed by N-hydroxybenzotriazole hydrate (Aldrich, 1.0 g, 5 mmol), O-(tetrahydro-2H-pyran-2-yl) hydroxylamine (1.1 g, 7.5 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (Sigma, 1.5 g, 9.4 mmol). The mixture was stirred overnight (about 18 hours) and was then stripped of solvent. The residue was taken up in ethyl acetate (250 mL) and washed with 5% NaHSO₄ (1x-150 mL), saturated potassium carbonate (1x-150 mL), and brine (1x-150 mL). The organic layer was then dried over Na₂SO₄ and concentrated to afford a viscous oil. ¹H NMR and mass spectroscopy showed the desired compound.

The viscous crude oil (3.0 g, 6.2 mmol) was dissolved in acetonitrile (10 mL) and stirred with 10% HCl_{aq} (15 mL) for 2 hours, after which LC showed no more starting material. The acetonitrile was removed with N₂ stream over the surface of the solution affording a solid that was collected, washed with H₂O (1x-20 mL), and dried *in vacuo* to afford the product as a tan solid (950 mg, 60% yield). ¹H NMR showed the desired compound. Mass spectroscopy showed: C₃₀H₄₂N₄O₅S - 2HCl; M⁺_{found} = 570 (M⁺_{calc} = 570).


-501-

Example 40: Preparation of 4-[[4-(4-bromophenyl)-4-fluoro-1-piperidinyl]sulfonyl]-tetrahydro-N-hydroxy-2H-pyran-4-carboxamide

5




Part 1: Preparation of



10 To 4-(4-Bromophenyl)-4-hydroxypiperidine-N-methylsulfonamide (Aldrich, 3g, 9 mmol) dissolved in CH_2Cl_2 (75 mL) and cooled to -78°C was added DAST (Aldrich, 1.9g 12 mmol). After the addition, the dry ice bath/reaction was left to warm to ambient
15 temperature. After the disappearance of the starting material, aqueous ammonium chloride (100 mL) was added and the layers were separated. The organic layer was dried over sodium sulfate. The solvent was removed under reduced pressure to give the
20 methylsulfonamide as a white solid (3 g). ^1H NMR and mass spectroscopy showed the desired compound.


Part 2: Preparation of

-502-



Oven-dried glassware was charged with the compound from Part 1 (4.0 g, 1.2 mmol) and tetrahydrofuran (25 mL,) and cooled to -75°C. Lithium bis(trimethylsilyl)amide (Aldrich, 1.0 M in tetrahydrofuran, 35 mL, 33 mmol) was slowly added, keeping temperature less than -60°C. The reaction was stirred for 30 minutes after the addition, and was then charged with a solution of methyl chloroformate (Aldrich, 1.3 mmol, 1.3 mmol) in tetrahydrofuran (17 mL), again keeping the temperature at less than -60°C. After stirring for 1 hour at -75°C, the reaction was quenched with saturated NH₄Cl, keeping temperature at less than -20°C. The aqueous portion freezes into a solid chunk of ice. After warming to 5°C, the mixture was extracted with ethyl acetate (3x- 200 mL). Organics were washed with saturated NH₄Cl (2x-100 mL) and brine (1x-100 mL), then dried over Na₂SO₄ and concentrated to afford the methylene sulfonamide as an amber oil (4.6 g, 90% crude yield). ¹H NMR and mass spectroscopy indicated desired compound.

Part 3: Preparation of




To a solution of compound from Part 2 (3.5 g, 1 mmol) and dibromo-diethylether (Lancaster, 3 g, 1.4

-503-

mmol) in dimethylformamide (28 mL) was added 18-Crown-6 (Aldrich, 500 mg, cat.) followed by potassium carbonate (Aldrich, 5 g, 3.6 mmol). The mixture was heated at 60°C for 16 hours. The product was isolated
5 by pouring into stirring 10% HCl_{aq} (200 mL) and extracted with ethyl acetate (3x- 300 mL). Organics were washed with brine (2x- 200 mL), dried over Na₂SO₄, and concentrated to afford an oil. The crystallized to result in 4.8 grams of the ester as
10 a tan solid. ¹H NMR and mass spectroscopy showed the desired compound.


Part 4: Preparation of



To a solution from Part 3 (2 g, 0.4 mmol) in
15 tetrahydrofuran (20 mL) was added potassium trimethylsilonate (Aldrich, 2 g, 1.5 mmol). The reaction was stirred overnight (about 18 hours) at room temperature. LC showed less than 3% starting material remained. Work up comprised removing the
20 tetrahydrofuran and taking the residue up in H₂O (100 mL). The solution was washed with diethyl ether (50 mL). The aqueous was then cooled to zero°C and 10% HCl_{aq} was slowly added until pH = 3. The product was filtered and washed with water to result in the acid
25 as a white solid (1.5 g, 72% yield). ¹H NMR and mass spectroscopy showed the desired compound.

Part 5: Preparation of

-504-




To a solution of the acid product in Part 4 (1 g, 0.2 mmol) in dimethylacetamide (10 mL) was added N-methylmorpholine (Aldrich, 2.0 mL, 2 mmol),
5 followed by N-hydroxybenzotriazole hydrate (Aldrich, 1.0 g, 0.7 mmol), O-(tetrahydro-2H-pyran-2-yl) hydroxylamine (1.1 g, 0.9 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (Sigma, 1.1 g, 0.6 mmol). The mixture
10 was stirred overnight (about 18 hours) and was then stripped of solvent. The residue was taken up in ethyl acetate (250 mL) and washed with 5% NaHSO₄ (1x-150 mL), saturated potassium carbonate (1x-150 mL), and brine (1x-150 mL). The organic was then dried
15 over Na₂SO₄ and concentrated to afford the protected hydroxamate as a viscous oil. ¹H NMR and mass spectroscopy showed the desired compound.

The viscous crude oil from above was dissolved in acetonitrile (10 mL) and stirred with 10% HCl_{aq} (15
20 mL) for 2 hours, after which LC showed no more starting material. The acetonitrile was removed with a N₂ stream over the surface of the solution affording a solid that was collected, washed with H₂O (1x-20 mL), and dried in vacuo to afford the product as a
25 tan solid (755 mg). ¹H NMR showed the desired compound. Mass spectroscopy showed: C₁₇H₂₂FN₂O₅SBrF
 $M^{+H}_{\text{found}} = 465$ ($M^{+H}_{\text{calc}} = 465$).


Example 41: Preparation of 4-[[4-[4-(3,5-

dimethylphenoxy)phenoxy]-1-piperidinyl]-
sulfonyl]tetrahydro-N-hydroxy-2H-pyran-
4-carboxamide



5


Part 1: Preparation of



To a slurry of Cs_2CO_3 (Aldrich, 40 g, 12 mmol) and 3,5-dimethyl phenol (Aldrich, 5 g, 4 mmol) in dimethylformamide (60 mL) at 25°C under N_2 was added
10 4-fluorobenzaldehyde (Aldrich, 5 g, 4 mmol). The mixture was stirred and heated to 90°C for 16 hours. After this time, the solvent was removed by roto-evaporation and taking the residue up in ethyl
15 acetate (150 mL) and H_2O (100 mL). The layers were separated and the aqueous layer was extracted with ethyl acetate (2x- 150 mL). The organics were washed with saturated K_2CO_3 (2x-100 mL), H_2O (1x-150 mL), and brine (1x- 150 mL), then dried over Na_2SO_4 , filtered,
20 and concentrated to afford a crude oil. The oil was purified on silica gel to give 8 g of the aldehyde as a clear oil. ^1H NMR and mass spectroscopy was consistent with the desired compound.


Part 2: Preparation of

-506-



To a solution of aldehyde (8 g, 35 mmol) from part 1 in methylene chloride (100 mL) was added meta-chloroperoxybenzoic acid (8 g, 52 mmol). The
5 reaction mixture was stirred at ambient temperature for 16 hours. After complete reaction, the solid meta-chlorobenzoic acid was removed by filtration. The solvent of the filtrate was removed under reduced pressure to give an oil. This oil was dissolved in
10 methanol (100 mL) to which lithium hydroxide (2 g) was added. After 4 hours, the reaction was complete. The solvent was removed by rotary evaporation to give an oil, which was dissolved in ethyl acetate and washed with 10% aqueous hydrochloric acid, separated
15 and dried over sodium sulfate to result in 5.5 grams of the phenol as an oil. ¹H NMR and mass spectroscopy were consistent with the desired compound.

Part 3: Preparation of




20 Sodium hydride (Aldrich, 2 g, 50 mmol) was added to a solution of 4-(methylsulfonyl)hydroxy-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester of Example 14, B (10 g, 25 mmol) and the phenol from part 2 (5.5 g, 50 mmol) dissolved in
25 dimethylformamide (60 mL) at 25°C and under N₂. The mixture was stirred and heated to 80°C for 16 hours. After this time, the solvent was removed by roto-

-507-


evaporation, followed by taking the residue up in ethyl acetate (150 mL) and H₂O (100 mL). The layers were separated and the aqueous was extracted via ethyl acetate (2x- 150 mL). The organics were washed
5 with saturated K₂CO₃ (2x-100 mL), H₂O (1x-150 mL), and brine (1x- 150 mL), then dried over Na₂SO₄, filtered, and concentrated to afford a crude oil. The oil was purified on silica gel to give 10g of the N-boc
10 piperidine as a clear oil. ¹H NMR and mass spectroscopy was consistent with the desired compound.

Part 4: Preparation of



To a solution of the product (10 g) of Part 3 in
15 1,4-dioxane (10 mL) was added 4 N HCl in dioxane (50 mL, 200 mmol). The mixture was stirred at room temperature until starting material was gone by LC (about 1 hour). The solvents were then removed and the residue was slurried in diethyl ether and
20 filtered. The solid was washed with diethyl ether (2x-50 mL) and dried in vacuo to afford a white solid (3g). ¹H NMR and mass spectroscopy showed the desired compound as the HCl salt.

Part 5: Preparation of




25

The HCl salt of Part 4 (3g, 10 mmol) and triethylamine (Aldrich, 3 mL, 15 mmol) were slurried

-508-

in CH₂Cl₂ (50 mL) and cooled to zero°C. A solution of methane sulfonyl chloride (Aldrich, 1.3 g, 13 mmol) in CH₂Cl₂ (20 mL) was slowly added, maintaining the temperature below 10°C. After the addition, the ice bath was removed and the reaction stirred 1 hour as it warmed to ambient temperature. After the disappearance of the starting material, the solvent was removed and the residue was taken up in ethyl acetate (100 mL) and H₂O (30 mL). Once separated, the organic layer was washed with 5% KHSO₄ (3x-50 mL) and brine (1x- 50 mL). The organic layer was then dried over Na₂SO₄, filtered, and concentrated to afford an oily solid that was recrystallized from diethyl ether, affording the methylsulfonamide as an off-white solid (1.2 g). ¹H NMR and mass spectroscopy showed the desired compound.


Part 6: Preparation of



Oven-dried glassware was charged with the compound from Part 5 (1.2 g, 3.2 mmol) and tetrahydrofuran (25 mL), and cooled to -75°C. Lithium bis(trimethylsilyl)amide (Aldrich, 1.0 M in tetrahydrofuran, 9 mL, 6 mmol) was slowly added, keeping temperature less than -60°C. The reaction was stirred for 30 minutes after the addition, and was then charged with a solution of methyl chloroformate (Aldrich, 350mg, 3.5 mmol) in tetrahydrofuran (1 mL) again keeping the temperature at less than -60°C. After stirring for 1 hour at -75°C, the reaction was quenched with saturated NH₄Cl, keeping temperature at

less than -20°C . The aqueous phase froze into a solid chunk of ice. After warming to 5°C , the mixture was extracted with ethyl acetate (3x- 200 mL). Organics were washed with saturated NH_4Cl (2x-100 mL) and brine (1x-100 mL), then dried over Na_2SO_4 and concentrated to afford the methylene sulfonamide as a tan oil (2.0 g, 90% crude yield). ^1H NMR and mass spectroscopy indicated desired compound.

Part 7: Preparation of




10

To a solution of compound from Part 6 (2 g, 11 mmol) and dibromo-diethylether (Lancaster, 1.8 mL, 15.1 mmol) in dimethylformamide (28 mL) was added 18-Crown-6 (Aldrich, 500mg, cat.) followed by potassium carbonate (Aldrich, 3.8 g, 27.4 mmol). The mixture was heated at 60°C for 16 hours. The product was isolated by pouring the reaction mixture into stirring 10% HCl_{aq} (200 mL), followed by extraction with ethyl acetate (3x- 300 mL). Organics were washed with brine (2x- 200 mL), dried over Na_2SO_4 , and concentrated to afford the ester as an oil. The oil was crystallized from diethyl ether (2 g). ^1H NMR and mass spectroscopy showed the desired compound.

20

Part 8: Preparation of




25

To a solution from Part 7 (2 g, 7 mmol) in tetrahydrofuran (20 mL) was added potassium

trimethylsilonate (Aldrich, 2. g, 18 mmol). The reaction was stirred overnight (about 18 hours) at room temperature. LC showed less than 3% starting material remained. Work up comprised removing the tetrahydrofuran and taking the residue up in H₂O (100 mL). The solution was washed with diethyl ether (50 mL). The aqueous layer was then cooled to zero°C and 10% HCl_{aq} was slowly added until pH = 3. The acidic mixture was then extracted via ethyl acetate (3x-150 mL). The organics were washed with brine (1x- 100 mL), dried over Na₂SO₄, and concentrated to afford a wet solid. The solid was dried in vacuo with phosphorous pentoxide yielding the acid as an orange solid (2 g, 92% yield). ¹H NMR and mass spectroscopy showed the desired compound.

Part 9: Preparation of




To a solution of the acid product in Part 8 (2 g, 6.2 mmol) in dimethylacetamide (10 mL) was added N-methylmorpholine (Aldrich, 2.0 mL, 18.6 mmol), followed by N-hydroxybenzotriazole hydrate (Aldrich, 1.0 g, 7.4 mmol), O-(tetrahydro-2H-pyran-2-yl) hydroxylamine (1.1 g, 9.4 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (Sigma, 1.8 g, 9.4 mmol). The mixture stirred overnight (about 18 hours), and was then stripped of solvent. The residue was taken up in ethyl acetate (250 mL) and washed with 5% NaHSO₄ (1x-150 mL), saturated potassium carbonate (1x-150 mL),

-511-


and brine (1x-150 mL). The organic layer was then dried over Na_2SO_4 and concentrated to afford a viscous oil. ^1H NMR and MS showed the desired compound.

The viscous crude oil (3.0 g, 6.2 mmol) was dissolved in acetonitrile (10 mL) and stirred with 10% HCl_{aq} (15 mL) for 2 hours, after which, LC showed no more starting material. The acetonitrile was removed with a N_2 stream over the surface of the solution affording a solid that was collected, washed with H_2O (1x-20 mL), and dried *in vacuo* to afford the product as a tan solid (230 mg, 64% yield). ^1H NMR showed the desired compound. Mass spectroscopy showed: $\text{C}_{25}\text{H}_{32}\text{F}_3\text{N}_2\text{O}_7\text{S}$ $M^+_{\text{found}} = 504$ ($M^+_{\text{calc}} = 504$).

Example 42: Preparation of 1-cyclopropyl-N-hydroxy-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide, monohydrochloride



Part 1: Preparation of




To a methanol (160 mL) solution of methyl 1-(phenylmethyl)-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxylate of

-512-

Example 12, Part F (40 g), and ammonium formate (15 g) was added 10%Pd on carbon (16 g, Degussa type catalyst). The black mixture was refluxed for 30-45 minutes. After complete reaction, the mixture was cooled and filtered through a Celite® pad. The solvent was removed under reduced pressure to give the piperidine methyl ester as an oil that solidified on standing (34 g). ¹H NMR and mass spectroscopy were consistent with the desired structure.


10 Part 2: Preparation of



To a solution of compound from Part 1 (5 g, 1 mmol) in methanol (35 mL) were added [(1-ethoxycyclopropyl)oxy]-trimethylsilane (Aldrich, 7 g, 4 mmol), acetic acid (6 g, 10 mmol), sodium cyanoborohydride (1.8 g, 3 mmol) and molecular sieves (2.5 g). The reaction mixture was stirred and heated for 8 hours. The progress of the reaction was monitored by RPHPLC. Work up comprised filtering the reaction mixture through a pad of Celite®, concentrating the methanol and partitioning the residue between in H₂O (50 mL) and ethyl acetate (500 mL). The organics were washed with brine (1x- 100 mL), dried over Na₂SO₄, and concentrated to afford the amino methyl ester as a semi-solid (3 g). ¹H NMR and mass spectroscopy were consistent with the desired structure.


25 Part 3: Preparation of

-513-



To a solution from Part 2 (3 g, 5 mmol) in tetrahydrofuran (20 mL) was added potassium trimethylsilanolate (Aldrich, 1.5 g, 10 mmol). The reaction was complete overnight (about 18 hours) at room temperature. A nitrogen stream was blown over the surface of the solution to concentrate the mixture. Then water was added (20 mL) followed by aqueous 10% HCl until pH = 7. The zwitterion was filtered washed with water (10 mL) and dried in vacuo with over phosphorous pentoxide yielding a solid (3 g). ¹H NMR and mass spectroscopy were consistent with the desired structure.

Part 4: Preparation of



15


To a solution of the zwitterion product in Part 3 (3 g, 5 mmol) in dimethylacetamide (20 mL) were added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (Sigma, 1.8 g, 9.4 mmol) and N-hydroxybenzotriazole hydrate (Aldrich, 1.0 g, 7.4 mmol), followed by heating to 50°C for 15 minutes. N-Methylmorpholine (Aldrich, 2.0 mL, 18.6 mmol) was added, followed by, O-(tetrahydro-2H-pyran-2-yl)

hydroxylamine (1.1 g, 9.4 mmol). The mixture was stirred and heated for 1 hour then left to stir at room temperature overnight (about 18 hours). After complete reaction, the reaction mixture was

5 concentrated under reduced pressure. The residue was taken up in ethyl acetate (250 mL) and washed with saturated potassium carbonate (1x-150 mL), and brine (1x-150 mL). The organic layer was dried over Na₂SO₄ and concentrated to afford a viscous oil. The oil

10 was purified on SiO₂ using ethyl acetate in hexane to give 1.8 g of the THP-protected hydroxamate as a clear oil the solidified on standing. ¹H NMR and mass spectroscopy were consistent with the desired structure.

15 Part 5: Preparation of



The solid from Part 4 (1.8 g) was slurried in methanol (600 mg) diethyl ether (10 mL). To this was added 4 N HCl in dioxane (10 mL) and stirred for 2


20 hours, after which, RPHPLC showed complete reaction. The dioxane was concentrated by half, diethyl ether was added (100 mL) and the white solid (1.6 g, 64% yield) filtered and dried under vacuum. ¹H NMR showed the desired compound. Mass spectroscopy showed:

25 C₂₁H₂₈F₃N₃O₅S-HCl M⁺_{found} = 527 (M⁺_{calc} = 527).


Example 43: Preparation of N-hydroxy-1-(iminophenylmethyl)-4-[[4-[4-(trifluoromethyl)phenoxy]-1-

- 515 -

piperidinyl)sulfonyl]-4-
piperidinecarboxamide, monohydrochloride




5 Part 1: Preparation of



To a methanol (160 mL) solution of methyl
 1-(phenylmethyl)-4-[[4-[[4-(trifluoromethyl)phenoxy]-
 1-piperidinyl]sulfonyl]-4-piperidinecarboxylate of
 10 Example 12, Part F (40 g), and ammonium formate (15
 g) was added 10%Pd on carbon (16 g, Degussa type
 catalyst). The black mixture was refluxed for 30-45
 minutes. After complete reaction the mixture is
 cooled and filtered through a Celite® pad. The
 15 solvent was removed under reduced pressure to give
 34 g of the amino ester as an oil that solidified on
 standing. ¹H NMR and mass spectroscopy were consistent
 with the desired structure.


Part 2: Preparation of



-516-

To a solution of compound from Part 1 (5 g, 1.0 mmol) in acetonitrile (50 mL) was added methyl benzimidate HCl (Aldrich, 2.5 g, 1.3 mmol) followed DMAP (100 mg). The reaction mixture was stirred and
5 heated to 60 degrees Celcius for 8 hours. The progress of the reaction was monitored by RPHPLC. Work up comprised removing the solvent under reduced pressure. The resulting solid was triturated with water and filtered to give 7 grams of the amidino
10 ester. ^1H NMR and mass spectroscopy were consistent with the desired structure.


Part 3: Preparation of



To a solution of the compound from Part 2 (5 g, 1 mmol) in tetrahydrofuran (45 mL) was added
15 potassium trimethylsilonate (Aldrich, 2.5 g, 3 mmol). The reaction was complete overnight (about 18 hours) at room temperature. A nitrogen stream was blown over the surface of the solution to concentrate the mixture. Then water was added (20 mL) followed by
20 aqueous 10% HCl until pH = 7. The zwitterion was filtered washed with water (10 mL) and dried in vacuo with over phosphorous pentoxide yielding the amidino acid as a solid (3 g). ^1H NMR and mass spectroscopy
25 were consistent with the desired structure.

Part 4: Preparation of


-517-



To a solution of the zwitterion product in Part 3 (2.7 g, 5 mmol) in dimethylacetamide (20 mL) were added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (Sigma, 2.5 g, 9.4 mmol), N-hydroxybenzotriazole hydrate (Aldrich, 1.9 g, 7.4 mmol) and heated to 50°C for 15 minutes. Thereafter, N-methylmorpholine (Aldrich, 4.0 mL, 18.6 mmol) was added, followed by O-(tetrahydro-2H-pyran-2-yl) hydroxylamine (2 g, 9.4 mmol). The mixture was stirred and heated for 1 hour, then left to stir at room temperature overnight (about 18 hours). After complete reaction, the reaction mixture was concentrated under reduced pressure. The residue was taken up in ethyl acetate (250 mL) and washed with saturated potassium carbonate (1x-150 mL), and brine (1x-150 mL). The organic layer was then dried over Na₂SO₄ and concentrated to afford the THP-protected amidino hydroxamate as a pink solid (1.7 g). ¹H NMR and mass spectroscopy were consistent with the desired structure.


Part 5: Preparation of

-518-




The solid from Part 4 (2.5 g) was slurried in methanol (800 mg) diethyl ether (30 mL). To this was added 4 N HCl in dioxane (10 mL) and the reaction mixture was stirred for 2 hours, after which RPHPLC showed complete reaction. The dioxane was concentrated by half, diethyl ether was added (100 mL) and the white solid (1 g, 70% yield) filtered and dried under vacuum. ¹H NMR showed the desired compound. Mass spectroscopy showed: C₂₅H₂₉F₃N₄O₅S - HClM⁺_{found} = 591 (M⁺_{calc} = 591).

Example 44: Preparation of N-hydroxy-1-[(4-hydroxyphenyl)iminomethyl]-4-[[4-(4-(trifluoromethyl)phenoxy)-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide, monohydrochloride




Part 1: Preparation of

-519-



To a methanol (160 mL) solution of methyl 1-
 (phenylmethyl)-4-[[4-[4-(trifluoromethyl)phenoxy]-1-
 piperidinyl]sulfonyl]-4-piperidinecarboxylate of
 5 Example 12, Part F (40 g), and ammonium formate (15
 g) was added 10%Pd on carbon (16 g, Degussa type
 catalyst). The black mixture was refluxed for 30-45
 minutes. After complete reaction, the mixture is
 cooled and filtered through a Celite® pad. The
 10 solvent was removed under reduced pressure to give 34
 g of the amino ester as an oil that solidified on
 standing. ¹H NMR and mass spectroscopy were
 consistent with the desired structure.

Part 2: Preparation of




15

To a solution of compound from Part 1 (6 g, 1.2
 mmol) in Dimethylformamide (20 mL) was added ethyl 4-
 hydroxybenzimidate HCl (Aldrich, 3 g, 1.4 mmol)
 followed by DMAP (150 mg). The reaction mixture was
 20 stirred and heated to 50 degrees Celcius for 8 hours.
 The progress of the reaction was monitored by
 RPHPLC. Work up comprised removing the solvent under
 reduced pressure. The resulting solid was triturated
 with water (50 mL) and filtered to give 5 grams of
 25 the amidino ester product. ¹H NMR and mass

-520-


spectroscopy were consistent with the desired structure.

Part 3: Preparation of



5 To a solution from Part 2 (3 g, 5 mmol) in tetrahydrofuran (20 mL) was added potassium trimethylsilonate (Aldrich, 1.5 g, 10 mmol). The reaction was complete overnight (about 18 hours) at room temperature. A nitrogen stream was blown over
10 the surface of the solution to concentrate the mixture. Then water was added (20 mL) followed by aqueous 10% HCl until pH = 7. The zwitterion was filtered, washed with water (10 mL) and dried in vacuo with over phosphorous pentoxide yielding the
15 amidino acid as a solid (2.4 g). ¹H NMR and mass spectroscopy were consistent with the desired structure.

Part 4: Preparation of




20 To a solution of the zwitterion product in Part 3 (1.1 g, 5 mmol) in dimethylacetamide (20 mL) were

-521-

added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (Sigma, 1.8 g, 9.4 mmol) and N-hydroxybenzotriazole hydrate (Aldrich, 1.0 g, 7.4 mmol), and the mixture was heated to 50°C for 15 minutes. N-methylmorpholine (Aldrich, 2.0 mL, 18.6 mmol) was then added, followed by O-(tetrahydro-2H-pyran-2-yl) hydroxylamine (1.1 g, 9.4 mmol). The mixture was stirred and heated for 1 hour, then left to stir at room temperature overnight (about 18 hours). After complete reaction, the reaction mixture was concentrated under reduced pressure. The residue was taken up in ethyl acetate (250 mL) and washed with saturated potassium carbonate (1x-150 mL), and brine (1x-150 mL). The organic layer was then dried over Na₂SO₄ and concentrated to afford the THP-protected hydroxamate as a viscous oil. The oil solidified (700 mg) on standing. ¹H NMR and mass spectroscopy were consistent with the desired structure.

20 Part 5: Preparation of




The solid from Part 4 (700 mg) was slurried in methanol (800 mg) and diethyl ether (30 mL). To this was added 4 N HCl in dioxane (10 mL) and stirred for 2 hours, after which RPHPLC showed complete reaction. The dioxane was concentrated by half, diethyl ether was added (100 mL) and the white solid (500 mg, 70% yield) filtered and dried under vacuum. ¹H NMR showed

-522-


the desired compound. Mass spectroscopy showed:

$C_{25}H_{29}F_3N_4O_6S \cdot HCl$; $M^+_{found} = 607$ ($M^+_{calc} = 607$).

Example 45: Preparation of 1-(2-furanylcarbonyl)-N-
hydroxy-4-[[4-[4-(trifluoromethyl)-
phenoxy]-1-piperidinyl]sulfonyl]-4-
piperidinecarboxamide




Part 1: Preparation of



To a methanol (160 mL) solution of methyl 1-(phenylmethyl)-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxylate of Example 12, Part F (40 g), and ammonium formate (15 g) was added 10%Pd on carbon (16 g, Degussa type catalyst). The black mixture was refluxed for 30-45 minutes. After complete reaction the mixture was cooled and filtered through a Celite® pad. The solvent was removed under reduced pressure to give 34 g of the amino ester as an oil that solidified on standing. 1H NMR and mass spectroscopy were consistent with the desired structure.


Part 2: Preparation of

- 523 -



To a solution of the compound from Part 1 (4 g, 0.8 mmol) in methylene chloride (75 mL) was added furfuryl chloride (Aldrich, 1.2 g, 0.8 mmol), followed by N-methylmorpholine (4 mL, 2.3 mmol). The reaction mixture was stirred for 1 hour. The progress of the reaction was monitored by RPHPLC. Work up comprised removing the solvent under reduced pressure. Water (50 mL) was added to the resulting solid and the product as extracted with ethyl acetate (100 mL). The ethyl acetate was washed with brine and dried over sodium sulfate to give the amide ester as a solid (6 grams of product). ^1H NMR and mass spectroscopy were consistent with the desired structure.


Part 3: Preparation of



To a solution of the compound from Part 2 (6 g, 5 mmol) in tetrahydrofuran (20 mL) was added potassium trimethylsilonate (Aldrich, 5 g, 15 mmol). The reaction was complete overnight (about 18 hours) at room temperature. A nitrogen stream was blown over the surface of the solution to concentrate the

mixture. Then water was added (20 mL), followed by aqueous 10% HCl until pH = 7. The product was extracted with methylene chloride (100 mL). The methylene chloride layer was dried over sodium sulfate and the solvent removed under reduced pressure to result in the amide acid as a tan solid (3g). ¹H NMR and mass spectroscopy were consistent with the desired structure.

Part 4: Preparation of



10


To a solution of the product in Part 3 (3 g, 5 mmol) in dimethylformamide (50 mL) were added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (Sigma, 1.8 g, 9.4 mmol) and N-hydroxybenzotriazole hydrate (Aldrich, 1.0 g, 7.4 mmol), and the mixture was heated to 50°C for 15 minutes. N-methylmorpholine (Aldrich, 2.0 mL, 18.6 mmol) was added, followed by O-(tetrahydro-2H-pyran-2-yl) hydroxylamine (1.1 g, 9.4 mmol). The mixture was stirred and heated for 1 hour then left to stir at room temperature overnight (about 18 hours). After complete reaction, the reaction mixture was concentrated under reduced pressure. The residue was taken up in ethyl acetate (250 mL) and washed with saturated potassium carbonate (1x-150 mL), and brine (1x-150 mL). The organic layer was then dried over

15
20
25

-525-

Na₂SO₄ and concentrated to afford THP-protected hydroxamate as a viscous oil. The oil solidified (2.1 g) on standing. ¹H NMR and mass spectroscopy were consistent with the desired structure.


5 Part 5: Preparation of




The solid from Part 4 (70 mg) was dissolved in acetonitrile (25mL). To this was added 10% aq. HCl (25 mL), and stirring for 2 hours, after which RPHPLC
10 showed complete reaction. A stream of nitrogen was blown of the surface on the reaction and concentrated by half to result in a precipitate. The white solid was filtered. The solid was crystallized in methanol to give (2.5 g) of a white solid and dried under
15 vacuum. ¹H NMR showed the desired compound. Mass spectroscopy showed: C₂₃H₂₆F₃N₃O₇S ; M⁺_{found} = 545 (M⁺_{calc} = 545).

Example 46: Preparation of N-hydroxy-1-[2-
20 (methylthio)-4-pyrimidinyl]-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-4-
piperidinecarboxamide, monohydrochloride

- 526 -




Part 1: Preparation of



To a methanol (160 mL) solution of methyl 1-
 5 (phenylmethyl)-4-[[4-[(trifluoromethyl)phenoxy]-1-
 piperidinyl]sulfonyl]-4-piperidinecarboxylate of
 Example 12, Part F (40 g), and ammonium formate (15
 g) was added 10%Pd on carbon (16 g, Degussa type
 catalyst). The black mixture was refluxed for 30-45
 10 minuets. After complete reaction the mixture is
 cooled and filtered through a Celite® pad. The
 solvent is removed under reduced pressure to give 34
 g of the amino ester as an oil that solidified on
 standing. ¹H NMR and mass spectroscopy were
 15 consistent with the desired structure.


Part 2: Preparation of



-527-

To a solution of the compound from Part 1 (5 g, 0.8 mmol) in dimethylformamide (50 mL) was added chloro-1-methylthiopiperazine (Aldrich, 1.2 g, 1.2 mmol) followed by potassium carbonate (4 g, 2.3 mmol). The reaction mixture was heated to 80 degrees Celsius and stirred for 1 hour, then stirred at room temperature for 12 hours. The progress of the reaction was monitored by RPHPLC. Work up comprised removing the solvent under reduced pressure. Water (50 mL) was added to the resulting solid and the product was filtered to give the N-heteroaryl ester as a solid (6.7 grams of product). ¹H NMR and mass spectroscopy were consistent with the desired structure.


Part 3: Preparation of



To a solution of the compound from Part 2 (6.5 g, 5 mmol) in tetrahydrofuran (50 mL) was added potassium trimethylsilonate (Aldrich, 5 g, 15 mmol). The reaction was complete overnight (about 18 hours) at room temperature. A nitrogen stream was blown over the surface of the solution to concentrate the mixture. Then water was added (20 mL) followed by aqueous 10% HCl until pH = 7. The product was extracted with methylene chloride (100 mL). The methylene chloride layer was dried over sodium sulfate and the solvent removed under reduced pressure to result in N-heteroaryl acid as a tan

solid (4.2g). ^1H NMR and mass spectroscopy were consistent with the desired structure.


Part 4: Preparation of



5 To a solution of the product in Part 3 (4.6 g, 5 mmol) in dimethylformamide (50 mL) were added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (Sigma, 5 g, 9.4 mmol) and N-hydroxybenzotriazole hydrate (Aldrich, 1.0 g, 7.4
10 mmol), and the mixture was heated to 50°C for 15 minutes. N-methylmorpholine (Aldrich, 4.0 mL, 18.6 mmol) was added, followed by O-(tetrahydro-2H-pyran-2-yl) hydroxylamine (6.1 g, 9.4 mmol). The mixture stirred and heated for 1 hour then left to stir at
15 room temperature overnight (about 18 hours). After complete reaction, the reaction mixture was concentrated under reduced pressure. The residue was taken up in water (75 mL) and filtered to result in the N-heteroaryl THP-protected hydroxamate as a pink
20 solid (4.1 g). ^1H NMR and mass spectroscopy were consistent with the desired structure.


Part 5: Preparation of

-529-



The solid from Part 4 (5.1 g) was dissolved in acetonitrile (25mL). To this was added 10% aq. HCl (25 mL), and the mixture was stirred for 2 hours, after which RPHPLC showed complete reaction. A stream of nitrogen was blown off the surface of the reaction and the volume concentrated by half to result in a precipitate. The white solid was filtered. The solid was crystallized in methanol to give (2.5 g) of a white solid and dried under vacuum. ^1H NMR showed the desired compound. Mass spectroscopy showed: $\text{C}_{23}\text{H}_{28}\text{F}_3\text{N}_5\text{O}_5\text{S}_2 \cdot \text{HCl}$; $M^+_{\text{found}} = 612$ ($M^+_{\text{calc}} = 612$).


Example 47: Preparation of 1-cyclopropyl-N-hydroxy-4-[[4-[4-(tri-fluoromethoxy)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide, monohydrochloride



20


Part 1: Preparation of

-530-



To a solution of 4-[[4-[4-(trifluoromethoxy)-
phenoxy]-1-piperidinyl]sulfonyl]-4-piperidine-
carboxylic acid, methyl ester from Part H of Example
5 15 (14 g, 30 mmol) in methanol (80 mL) were added
[(1-ethoxycyclopropyl)oxy]trimethylsilane (Aldrich,
21 g, 120 mmol), acetic acid (18g, 300 mmol), sodium
cyanoborohydride (5.5 g, 90 mmol) and molecular
sieves (7 g). The reaction mixture was stirred and
10 heated for 8 hours. The progress of the reaction was
monitored by RPHPLC. Work up comprised filtering the
reaction mixture through a pad of Celite® then
concentrating the methanol and partitioning the
residue between in H₂O (50 mL) and ethyl acetate (500
15 mL). The organics were washed with brine (1x- 100
mL), dried over Na₂SO₄, and concentrated to afford the
amino ester as a solid (8 g) after crystallization in
methanol. ¹H NMR and mass spectroscopy were
consistent with the desired structure.


20 Part 2: Preparation of



To a solution of the compound from Part 1 (8 g,
15 mmol) in tetrahydrofuran (20 mL) was added
potassium trimethylsilonate (Aldrich, 4 g, 30 mmol).
25 The reaction was complete overnight (about 18 hours)

at room temperature. A nitrogen stream was blown over the surface of the solution to concentrate the mixture. Then water was added (20 mL) followed by aqueous 10% HCl until pH = 7. The zwitterion was
5 filtered washed with water (10 mL) and dried *in vacuo* with over phosphorous pentoxide yielding the amino acid as a solid (6.8 g). ¹H NMR and mass spectroscopy were consistent with the desired structure.

Part 3: Preparation of




10

To a solution of the zwitterion product in Part 2 (6.8 g, 14 mmol) in dimethylformamide (50 mL) were added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (Sigma, 5.2 g, 28 mmol) and N-
15 hydroxybenzotriazole hydrate (Aldrich, 3.7 g, 28 mmol), and the mixture was heated to 50°C for 15 minutes. N-methylmorpholine (Aldrich, 2.0 mL, 28 mmol) was added followed by O-(tetrahydro-2H-pyran-2-yl) hydroxylamine (3.2 g, 28 mmol). The mixture
20 stirred and heated for 1 hour then left to stir at room temperature overnight (about 18 hours). After complete reaction, the reaction mixture was concentrated under reduced pressure. The residue was taken up in ethyl acetate (250 mL) and washed with
25 saturated potassium carbonate (1x-150 mL), and brine (1x-150 mL). The organic layer was then dried over Na₂SO₄ and concentrated to afford a viscous oil. The

-532-


oil was purified on SiO₂ using ethyl acetate in hexane to give 6.5 g of the THP-protected hydroxamate as a clear oil the solidified with addition of methanol. ¹H NMR and mass spectroscopy were consistent with the
5 desired structure.

Part 4: Preparation of



The solid from Part 3 (6.8 g) was slurried in methanol (4.5 mL). To this was added 4 N HCl in
10 dioxane (60 mL), and the mixture stirred for 2 hours, after which RPHPLC showed complete reaction. The dioxane was concentrated by half, diethyl ether was added (100 mL) and the white solid (5.5 g, 90% yield) filtered and dried under vacuum. ¹H NMR showed the
15 desired compound. Mass spectroscopy showed:
C₂₁H₂₈F₃N₃O₆S -HCl; M⁺_{found} = 543 (M⁺_{calc} = 543).

Example 48: Preparation of 4-(1,4-dioxo-8-azaspiro-
[4.5]dec-8-ylsulfonyl)tetrahydro-N-
20 hydroxy-2H-pyran-4-carboxamide



Part A: 1,4-Dioxo-8-azaspiro[4.5]-decane (Aldrich, 10.0 g, 69.8 mmol) and triethylamine
25 (Aldrich, 14.6 mL, 105 mmol) were slurried in dichloromethane (150 mL) and cooled to zero°C. A

solution of methane sulfonyl chloride (Aldrich, 8.1 mL, 105 mmol) in dichloromethane (60 mL) was slowly added, maintaining the temperature at less than 10°C. After the addition, the ice bath was removed and the reaction stirred 1 hour as it came to ambient temperature. After the disappearance of the starting material, the solvent was removed and the residue was taken up in ethyl acetate (200 mL) and H₂O (50 mL). Once separated, the organic layer was washed with 5% KHSO₄ aq (3x-50 mL) and brine (1x- 50 mL). The organic layer was then dried over Na₂SO₄, filtered, and concentrated to afford the sulfonamide as a yellow solid (15.3 g, 99% yield). ¹H NMR showed the desired compound.

Part B: Oven-dried glassware was charged with the sulfonamide of Part A (15.3 g, 69.1 mmol) and tetrahydrofuran (70 mL) and cooled to -75°C. Lithium bis(trimethylsilyl)amide (Aldrich, 1.0 M in tetrahydrofuran, 140 mL, 140 mmol) was slowly added, keeping temperature less than -60°C. The reaction was stirred for 30 minutes after the addition and was then charged with a solution of methyl chloroformate (Aldrich, 5.4 mL, 69.1 mmol) in tetrahydrofuran (30 mL), again keeping the temperature at less than -60°C. After stirring for 1 hour at -75°C, the reaction was quenched with saturated NH₄Cl_{aq}, keeping temperature below -20°C. The aqueous phase froze into a solid chunk of ice. After warming to 5°C, the mixture was extracted via ethyl acetate (3x- 200 mL). Organics were washed with saturated NH₄Cl_{aq} (2x-100 mL) and brine (1x-100ml), then dried over Na₂SO₄ and concentrated to afford a yellow oil that slowly solidified. The oily solid was recrystallized from

ethyl acetate and hexanes to afford the methylene ester (8.2 g, 44 % yield). ^1H showed the desired compound with some starting material present.

Part C: To a solution of the methylene ester of Part B (3.5 g, 12.5 mmol) and dibromo-diethylether (Lancaster, 1.7 mL, 13.8 mmol) in N,N-dimethylformamide (25 mL) was added 18-Crown-6 (Aldrich, 500 mg, 1.9 mmol), followed by potassium carbonate (Aldrich, 6.0 g, 43.8 mmol). The mixture was heated at 60°C for 4 hours after which more potassium carbonate (1.9 g, 13.7 mmol) was added, and the reaction continued at 60°C for 14 hours. The reaction was worked up by pouring the mixture into stirring 10% HCl_{aq} (200 mL). A gummy solid resulted that was extracted via ethyl acetate (3x- 300 mL). Organics were washed with brine (2x- 200 mL), dried over Na_2SO_4 , and concentrated to afford the methyl ester, 9091-157, as a dark brown oil (2.2 g, 50% yield). ^1H NMR showed the desired compound.


Part D: To a solution of the methyl ester from Part C (2.2 g, 6.3 mmol) in tetrahydrofuran (15 mL) was added potassium trimethylsilonate (Aldrich, 1.9 g, 15.1 mmol). Work up comprised removing the tetrahydrofuran and taking the residue up in H_2O (100 mL). The solution was washed with diethyl ether (50 mL). The aqueous was then cooled to zero°C and 10% HCl_{aq} was slowly added until pH = 3. The acidic mixture was then extracted via ethyl acetate (3x-150 mL). The organics were washed with brine (1x- 100 mL), dried over Na_2SO_4 , and concentrated to afford the carboxylic acid as a dark oil (1.5 g, 70% yield). ^1H NMR showed the desired compound.

Part E: To a solution of the acid of Part D (1.5 g, 4.5 mmol) in dimethylacetamide (10 mL) was added N-methylmorpholine (Aldrich, 1.5 mL, 13.5 mmol), followed by N-hydroxybenzotriazole hydrate (Aldrich, 0.73 g, 5.4 mmol), O-(tetrahydro-2H-pyran-2-yl) hydroxylamine (0.77 g, 6.7 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (Sigma, 1.3 g, 6.7 mmol). The mixture stirred overnight (about 18 hours) and was then stripped of solvent. The residue was taken up in ethyl acetate (300 mL) and water (20 mL). After separation, the organic layer was washed with 5% KHSO₄ aq (1x- 200 mL), saturated potassium carbonate (1x-100 mL), and brine (1x-150 mL). The organic layer was then dried over Na₂SO₄ and concentrated to afford the THP-protected hydroxamate, 9091-161, as a brown foam (1.6 g, 76% yield). ¹H NMR showed the desired compound.

Part F: The THP-protected hydroxamate of Part E (1.6 g, 3.7 mmol) was dissolved in acetonitrile (15 mL) and stirred with 10% HCl_{aq} (20 mL) for 2 hours. The acetonitrile was removed via N₂ stream giving an oil that was crystallized from t-butylmethylether to afford the hydroxamate as a brown solid (0.82 g, 63% yield). ¹H NMR showed the desired compound. HRMS for C₁₃H₂₂N₂O₇S showed M⁺_{found} = 351 (M⁺_{calc} = 351).

Example 49: Preparation of 4-[[4-[[[(3R,5R)-rel-3,5-dimethyl-1-piperidinyl]carbonyl]-1-piperidinyl]sulfonyl]tetrahydro-N-hydroxy-2H-pyran-4-carboxamide and 4-[[4-[[[(3R,5S)-rel-3,5-dimethyl-1-piperidinyl]carbonyl]-1-

piperidinyl)sulfonyl]tetrahydro-N-
hydroxy-2H-pyran-4-carboxamide



Part A: To a solution of N-BOC-isoponic acid (4.0g, 17.4 mmol) in dimethylacetamide (30 mL) were added N-methylmorpholine (Aldrich, 5.7 mL, 52.2 mmol), 3,5-dimethylpiperadine (70% cis, Aldrich, 3.5 mL, 26.1 mmol), N-hydroxybenzotriazole hydrate (Aldrich, 2.8 g, 20.9 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (Sigma, 5.0 g, 26.1 mmol). The mixture was stirred overnight (about 18 hours) at ambient temperature. To work up, the solvent was removed and the residue was taken up in ethyl acetate (300 mL) then washed with 5% KHSO₄ aq (1x 100ml), saturated K₂CO₃ aq (1x 100 mL), and brine (1x 100 mL). After drying over Na₂SO₄, the organic layer was filtered and concentrated to afford the BOC-piperidine as a tan oil that slowly crystallized (5.4 g, 96% yield). ¹H NMR showed the desired compound existing as both the cis- and trans- isomers.

Part B: To a solution of the BOC-piperidine of Part A, (5.4 g, 16.6 mmol) in 1,4-dioxane (20 mL) was added 4 N HCl in 1,4-dioxane (15 mL) and stirred. After 45 minutes, the solvent was removed and the solid was slurried in diethyl ether, filtered, and washed with diethyl ether (2 x 20 mL). Drying

-537-

afforded the amine salt as a white solid (2.6 g, 60% yield). ^1H NMR showed desired product as both cis- and trans- isomers.

Part C: To a cooled solution (zero°C) of the
5 amine salt of Part B (2.2 g, 9.8 mmol) and triethylamine (Aldrich, 3.4 mL, 24.5 mmol) in dichloromethane (50 mL) was slowly added a solution of methane sulfonyl chloride (Aldrich, 1.1 mL, 14.7 mmol) in dichloromethane (25 mL), maintaining the
10 temperature at less than 10°C. After the addition, the ice bath was removed and the reaction stirred 2 hours as it came to ambient temperature. After the disappearance of the starting material, the solvent was stripped and the residue was taken up in ethyl
15 acetate (100 mL) and H_2O (25 mL). Once separated, the organic layer was washed with 5% KHSO_4 aq (3x-50 mL) and brine (1x- 50 mL). The organic layer was then dried over Na_2SO_4 , filtered, and concentrated to afford the sulfonamide as a dried foamy oil (2.5 g,
20 83% yield). ^1H NMR showed the desired product as both the cis- and trans- isomers.

Part D: Oven-dried glassware was charged with the sulfonamide of Part C (2.5 g, 8.3 mmol) and tetrahydrofuran (17 mL) and cooled to -75°C. Lithium
25 bis(trimethylsilyl)amide (Aldrich, 1.0 M in tetrahydrofuran, 16.5 mL, 16.5 mmol) was slowly added, keeping temperature less than -60°C. The reaction was stirred for 30 minutes after the addition and was then charged with a solution of
30 methyl chloroformate (Aldrich, 0.64 mL, 8.3 mmol) in tetrahydrofuran (8 mL), again keeping the temperature at less than -60°C. After stirring for 1 hour at

-75°C, the reaction was quenched with saturated $\text{NH}_4\text{Cl}_{\text{aq}}$ (50 mL), keeping temperature at less than -20°C. The aqueous portion froze into a solid chunk of ice. After warming to 5°C, the mixture was extracted with
5 ethyl acetate (3x- 100 mL). Organics were washed with saturated $\text{NH}_4\text{Cl}_{\text{aq}}$ (2x-50 mL) and brine (1x-50 mL), then dried over Na_2SO_4 and concentrated to afford the methylene ester as a brown oil (2.7 g, 90 % crude yield). ^1H showed the desired compound (cis- and
10 trans-).

Part E- To a solution of the methylene ester of Part D (2.7 g, 7.5 mmol) and dibromo-diethylether (Lancaster, 1.0 mL, 8.2 mmol) in N,N-dimethylformamide (15 mL) was added 18-Crown-6
15 (Aldrich, 200 mg, 1.0 mmol) followed by potassium carbonate (Aldrich, 3.6 g, 26.2 mmol). The mixture was heated at 60°C for 4 hours, after which more potassium carbonate (1.0 g, 7.2 mmol) was added, and the reaction continued at 60°C for 14 hours. The
20 reaction was worked up by pouring the mixture into stirring 10% HCl_{aq} (200 mL). The gummy solid that developed was extracted via ethyl acetate (3x 100 mL). The organics were washed with brine (1x 50 mL), dried over Na_2SO_4 , and filtered to afford the methyl
25 ester as a brown oil (3.4 g, quantitative yield). ^1H NMR showed the desired compound (cis- and trans-).

Part F: To a solution of the methyl ester from Part E (3.0 g, 7.0 mmol) in tetrahydrofuran (20 mL) was added potassium trimethylsilonate (Aldrich, 2.2
30 g, 17.4 mmol). The reaction was stirred overnight (about 18 hours) at ambient temperature. The work up consisted of stripping the tetrahydrofuran and taking the residue up in H_2O (40 mL). The solution was

washed with diethyl ether (30 mL). The aqueous composition was then cooled to 0°C and 10% HCl_{aq} was slowly added until pH = 3. The acidic mixture was then extracted via ethyl acetate (3x-100 mL). The organics were washed with brine (1x- 50 mL), dried over Na₂SO₄, and concentrated to give a brown oil (2.4 g). The oil was crystallized from acetone/diethylether to afford the carboxylic acid as a brown solid (2.2 g, 76% yield). ¹H showed the desired compound as cis- and trans- isomers.

Part G: To a solution of the acid from Part F (2.2 g, 5.3 mmol) in dimethylacetamide (12 mL) were added N-methylmorpholine (Aldrich, 1.7 mL, 15.9 mmol) followed by N-hydroxybenzotriazole hydrate (Aldrich, 0.86 g, 6.4 mmol), O-(tetrahydro-2H-pyran-2-yl) hydroxylamine (0.94 g, 8.0 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (Sigma, 1.5 g, 8.0 mmol). The mixture stirred overnight and was then stripped of solvent. The residue was taken up in ethyl acetate (250 mL) and washed with 5% NaHSO₄ _{aq} (1x- 150 mL), saturated potassium carbonate (1x-150 mL), and brine (1x-150 mL). The organic was then dried over Na₂SO₄ and concentrated to afford the THP-protected hydroxamate as a brown oil (2.2 g, 81% yield). ¹H NMR showed the desired compound as both isomers.

Part H - The THP-protected hydroxamate of Part G (2.2 g, 3.7 mmol) was dissolved in 1,4-dioxane (10 mL) and stirred with 4 N HCl in 1,4-dioxane (15 mL, 60 mmol) for 2 hours. The solvents were stripped and the residue was taken up in acetonitrile (10 mL) and H₂O (10 mL). This solution was filtered and purified via prep RP LC to afford the cis isomer as a white


-540-

solid (0.40 g, 22% yield) and the trans isomer as a white solid (0.20 g, 11%). ^1H NMR showed the desired compounds. HRMS for $\text{C}_{19}\text{H}_{33}\text{N}_3\text{O}_6\text{S}$ showed $M^{+H}_{\text{found}} = 432$ ($M^{+H}_{\text{calc}} = 432$) for both isomers.

5

Example 50: Preparation of N-hydroxy-1-(4-methylphenyl)-4-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide, monohydrochloride

10



Part A: To a mixture of dibromotriphenylphosphorane (Aldrich, 50 g, 118 mmol) in dichlorethane (200 mL) at zero C was added N,N-bis-(2-hydroxyethyl)-P-toluidine (Fluka, 10.5 g, 54 mmol). A slight exotherm was detected. The ice bath was removed and the reaction stirred eighteen hours at ambient temperature. After completion, the solvent was evaporated affording an oily solid that was purified by silica gel (ethyl acetate) to yield the N,N-bis(2-bromoethyl)-p-toluidine as a white solid (12.6 g, 73% yield). ^1H NMR showed the desired compound.

Part B: To a solution of the sulfonamide ester of part D of Example 36 (12.0 g, 31.5 mmol) in N,N-dimethylformamide (63 mL) were added potassium carbonate (Aldrich, 13.0 g, 94.5 mmol), 18-crown-6 (Aldrich, 500 mg, 2.0 mmol), and N,N-bis(2-bromoethyl)-p-toluidine of Part A (10.0 g, 31.5 mmol). The mixture was heated at 80°C for 18 hours after which more potassium carbonate (1.0 g, 7.2 mmol) was added, and the reaction continued at 80°C for 3 hours. The reaction was worked up by removing the solvent by roto-evaporation. The residue was slurried in ethyl acetate and filtered through a Celite® pad. The filtrate was concentrated and the residue crystallized from hot methanol to afford the methyl ester as a yellow solid (9.0 g, 53% yield). ¹H and ¹⁹F NMR showed the desired compound.

Part C: To a solution of the methyl ester of Part B (9.0 g, 16.6 mmol) in tetrahydrofuran (40 mL) was added potassium trimethylsilonate (Aldrich, 6.4 g, 50 mmol). The reaction stirred overnight (about 18 hours) at ambient temperature. Work up comprised stripping the tetrahydrofuran and taking the residue up in H₂O (5 mL), cooling to zero°C, and titrating to pH 7 via 6 N HCl. Solids formed and were collected then washed with diethyl ether. The solid was dried in vacuo to afford the carboxylic acid as a white solid (8.7 g, 100% yield). ¹H and ¹⁹F NMR showed the desired compound.


Part D: To a solution of the carboxylic acid of Part C (8.7 g, 16.5 mmol) in N,N-dimethylformamide (35 mL) was added N-methylmorpholine (Aldrich, 5.4 mL, 49.5 mmol) followed by N-hydroxybenzotriazole hydrate (Aldrich, 2.7 g, 19.8 mmol), O-(tetrahydro-

-542-

2H-pyran-2-yl) hydroxylamine (3.9 g, 33 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (Sigma, 4.7 g, 24.8 mmol). The mixture stirred at 40 C for 4 hours and was then stripped of solvent. The residue was taken up in ethyl acetate/water (350 mL/50 mL). After separation, the aqueous layer was extracted with ethyl acetate (2x-50 mL). The combined organic layers were washed with 5% KHSO₄aq (1x- 30 mL) and brine (1x-30 mL). The organic layer was then dried over Na₂SO₄ and concentrated yielding an orange oily foam that was recrystallized from methanol to afford the THP-protected hydroxamate as a white solid (0.74 g, 64% yield). ¹H and ¹⁹F NMR showed the desired compound.

Part E: The THP-protected hydroxamate of Part D (8.9 g, 14.2 mmol) was wetted with methanol (3.6 mL) and stirred with 4 N HCl in dioxane (36 mL) for one hour. The solvents were evaporated and the oil slurried in diethyl ether to yield a solid which was filtered and dried. This afforded hydroxamate as a white solid (8.0 g, 98% yield). ¹H and ¹⁹F NMR showed the desired compound. HRMS for C₂₅H₃₀F₃N₃O₅S showed M⁺_{found} = 542 (M⁺_{calc} = 542).

Example 51: Preparation of N-hydroxy-1-(4-methylphenyl)-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]-sulfonyl]-4-piperidinecarboxamide, monohydrochloride



Part A: To a solution of N,N-bis(2-hydroxyethyl)-p-toluidine (Fluka, 30 g, 154 mmol) in dichloromethane (100 mL) was slowly added a solution of thionylchloride (Aldrich, 36 mL, 462 mmol) in dichloromethane (300 mL). The reaction was stirred for 2 hours at ambient temperature then heated for 1 hour at reflux. After removing the solvent, the residue was slurried in ethyl acetate. The ethyl acetate was decanted, then the residue was slurried in hexanes giving a solid precipitate. The solid was filtered and washed with hexanes followed by diethyl ether. The solid was dried to afford the N,N-bis(2-dichloroethyl)-p-toluidine monochloride salt as a gray solid (24 g, 58% yield). ¹H showed the desired compound.

Part B: The N,N-bis(2-dichloroethyl)-p-toluidine monochloride salt of Part A was suspended in water (200 mL) and neutralized (pH 7) with saturated sodium bicarbonate. This was then extracted with a mixture of diethyl ether (200 mL) and ethyl acetate (50 mL) which dissolved all solids. The organic layer was washed with brine (1x-100 mL) then dried over Na₂SO₄, filtered and concentrated to

-544-

give the free base amine as a brown oil (6.59 g, 28.39 mmol). The oil was dissolved in 4-methyl-2-pentanone (50 mL) and lithium bromide (Aldrich, 24.66 g, 283.90 mmol) was added. The mixture was heated at
5 reflux for 39 hours. After cooling, the dark mixture was filtered through a Celite pad. After washing the pad, the total filtrate was concentrated. The residue was partitioned between diethyl ether (100 mL) and water (50 mL). The organic layer was washed with
10 water (50ml), brine (50 mL), then dried over Na₂SO₄ and concentrated to afford the dibromo-amine as a brown oil (7.82 g, 86% yield). ¹H showed the desired compound. LCMS showed mixture of dibromo- and monochloro/monobromo compounds.

15 Part C: To a slurry of 4-hydroxypiperadine (Aldrich, 273 g, 2.7 mol) in tetrahydrofuran (1.8 L) was added triethylamine (Aldrich, 402 mL, 2.9 mol), followed by slow addition of a solution of di-tert-butyl-dicarbonate (Aldrich, 600 g, 2.7 mol) in
20 tetrahydrofuran (1.2 L). The temperature was monitored and maintained below 32°C. The mixture was stirred for 4 hours before working up. Work up comprised removing the tetrahydrofuran by roto-evaporation and taking the residue up in ethyl
25 acetate (300 mL). The organic was washed with 5% KHSO₄ aq (2x-1 L) and brine (1x-1 L) then dried over anhydrous Na₂SO₄, filtered, and concentrated to afford a crude brown oil. The oil was crystallized from hexanes providing the N-BOC-4-hydroxypiperidine as a
30 tan solid (515 g, 95% yield). ¹H NMR showed the desired compound.

Part D: To a slurry of the N-BOC-4-hydroxypiperidine of Part C (515 g, 2.6 mol) and

-545-

triethylamine (390 mL, 2.8 mol) in dichloromethane (1.9 L) cooled to zero°C under N₂ was slowly added a solution of methane sulfonyl chloride (Aldrich, 209 mL, 2.7 mol) in dichloromethane (100 mL). After the
5 addition, the ice bath was removed and the reaction stirred 24 hours. Work up comprised removing the solvent by roto-evaporation and taking the residue up in ethyl acetate (2.0 L) then washing with H₂O (2x-1 L), brine (1x- 1 L), and drying over Na₂SO₄. The
10 mixture was filtered and concentrated to afford the piperidine mesylate, SC 80998, as a cream colored solid (662 g, 91% crude yield). ¹H NMR showed the desired compound.

Part E: Oven-dried glassware was charged with 4-
15 trifluoromethoxyphenol (Apollo, 100.0 g, 561 mmol) and N,N-dimethylformamide (1.12 L). After cooling to -5 °C, NaH (Aldrich, 60% oil dispersion, 11.2 g, 281 mmol) was slowly added. The ice bath was removed and the reaction stirred for 1.5 hours. The piperidine
20 mesylate of Part D (78.4 g, 281 mmol) was then added and the reaction was heated to 80 °C for 2 hours. This process was repeated until a total of 2.5 equivalents (1.4 moles) of both the piperidine mesylate and NaH were added. Work up consisted of
25 stripping the solvents by roto-evaporation and taking the residue up in diethyl ether (1 L) and H₂O (400 mL). The layers were separated and the organic layer was washed with H₂O (1x-500 mL) and brine (2x- 500 mL) then dried over Na₂SO₄, filtered, and concentrated to
30 afford the BOC-piperidine, SC 83075, as a crude oil (315 g, 100% crude yield). ¹H NMR showed the desired compound along with the elimination byproduct, 1-tert-butoxycarbonyl-1,2,3,6-tetrahydropyridine.

Part F: To an overhead-stirring solution of 4 N HCl in dioxane (1.4 L, 5.6 mol) was poured the crude oil of the BOC-piperidine of Part E (203 g, 561 mmol). The solvents were then stripped and the
5 residue was slurried in diethyl ether and filtered. The solid was dissolved in H₂O (500 mL) and titrated to pH 10 with saturated potassium carbonate aqueous solution. The aqueous was extracted with dichloromethane (3x-800ml). The organics were
10 combined, dried over Na₂SO₄, filtered and concentrated to afford the piperidine, 10507-054, as a foamy solid (136 g, 93% yield). ¹H NMR showed the desired compound.

Part G: The piperidine of Part F (136 g, 520 mmol) and triethylamine (Aldrich, 110 mL, 781 mmol)
15 were slurried in dichloromethane (1.6 L) and cooled to zero°C. A solution of methane sulfonyl chloride (Aldrich, 60.2 mL, 781 mmol) in dichloromethane (200 mL) was slowly added, maintaining the temperature at
20 less than 10°C. After the addition, the ice bath was removed and the reaction came to ambient temperature. After one hour, the starting material was gone. To work up, the solvent was stripped and the residue was taken up in ethyl acetate (1 L) and H₂O (1 L). Once
25 separated, the aqueous layer was extracted with ethyl acetate (2x- 400 mL). The combined organic layers were washed with 5% KHSO₄ aq (2x-600 mL) and brine (1x-600 mL). The organic layers were then dried over Na₂SO₄, filtered, and concentrated to afford the
30 piperidine mesylate, SC 80766, as a brown solid (139 g, 79% yield). ¹H NMR showed the desired compound.

Part H: Oven-dried glassware was charged with the piperidine mesylate of Part G (92 g, 271 mmol)

-547-

and tetrahydrofuran (600 mL) and cooled to -75°C. Lithium bis(trimethylsilyl)amide (Aldrich, 1.0 M in tetrahydrofuran, 705 mL, 705 mmol) was slowly added, keeping temperature below -60°C. The reaction was
5 stirred for 30 minutes after the addition and was then charged with a solution of methyl chloroformate (Aldrich, 27.4 mL, 325 mmol) in tetrahydrofuran (100 mL) again keeping the temperature below -60°C. After stirring for 1 hour at -75°C, the reaction was
10 quenched with saturated $\text{NH}_4\text{Cl}_{\text{aq}}$, keeping temperature below -20°C. The aqueous phase froze into a solid chunk of ice. After warming to 5°C, the mixture was extracted via ethyl acetate (3x- 200 mL). Organics were washed with saturated $\text{NH}_4\text{Cl}_{\text{aq}}$ (2x-100 mL) and
15 brine (1x-100ml), then dried over Na_2SO_4 and concentrated to afford a tan oil. The oil was crystallized from methanol. The solid was collected and washed with hexanes to afford the methylene ester as a tan solid (78 g, 72%). ^1H NMR showed the desired
20 compound with some starting material present.

Part I: To a mixture of the methylene ester of Part H (4.0 g, 10.0 mmol), potassium carbonate (Aldrich, 4.1 g, 30.0 mmol), and 18-crown-6 (Aldrich, 0.1 g, .04 mmol) in N,N-dimethylformamide (20 mL) was
25 added the dibromo-amine of Part B (3.2 g, 10.0 mmol). The mixture was heated at 80°C for 18 hours, after which more potassium carbonate (1.5 g, 12 mmol) was added, and the reaction continued at 80°C for 14 hours. The reaction was worked up by removing the
30 solvent by roto-evaporation. The residue was slurried in acetonitrile and filtered through a Celite pad. The filtrate was concentrated and the residue was purified on silica gel (ethyl

acetate/hexanes) to afford the methyl ester as an orange solid (2.6 g, 46% yield). ^1H and ^{19}F NMR showed the desired compound.

Part J: To a solution of the methyl ester of
5 Part I (2.1 g, 3.8 mmol) in tetrahydrofuran (10 mL) was added potassium trimethylsilonate (Aldrich, 1.4 g, 11.3 mmol). The reaction was stirred overnight (about 18 hours) at ambient temperature. Work up comprised removing the tetrahydrofuran and taking the
10 residue up in H_2O (5 mL), cooling to zero°C, and titrating to pH 7 with 6 N HCl. Solids formed and were collected then washed with acetonitrile. The solid was dried in vacuo to afford the carboxylic acid, X14137, as a tan solid (1.0 g, 50% yield). ^1H
15 and ^{19}F NMR showed the desired compound.


Part K: To a solution of the carboxylic acid of Part J (1.0 g, 1.8 mmol) in N,N-dimethylformamide (5 mL) was added N-methylmorpholine (Aldrich, 0.6 mL, 5.5 mmol) followed by N-hydroxybenzotriazole hydrate
20 (Aldrich, 0.29 g, 2.2 mmol), O-(tetrahydro-2H-pyran-2-yl) hydroxylamine (0.42 g, 3.6 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (Sigma, 0.52 g, 2.7 mmol). The mixture was stirred at 40 C for 8 hours and was then stripped
25 of solvent. The residue was taken up in ethyl acetate/water (50 mL/10 mL). After separation, the aqueous layer was extracted with ethyl acetate (2x-20 mL). The combined organic layers were washed with 5% NaHSO_4 aq (1x-30 mL) and brine (1x-30 mL). The
30 organic was then dried over Na_2SO_4 and concentrated yielding an orange oily foam that was recrystallized from methanol to afford the THP-protected hydroxamate

-549-

as a white solid (0.74 g, 64% yield). ^1H and ^{19}F NMR showed the desired compound.

Part L: The protected hydroxamate of Part K (0.7 g, 1.1 mmol) was dissolved in methanol (0.5 mL) and stirred with 4 N HCl in dioxane (2.8 mL) for one hour. After which, LC showed no more starting material. The solvents were evaporated and the oil slurried in diethyl ether to yield a solid which was filtered and dried. This afforded the hydroxamate, as a white solid (0.6 g, 92% yield). ^1H and ^{19}F NMR showed the desired compound. HRMS for $\text{C}_{25}\text{H}_{30}\text{F}_3\text{N}_3\text{O}_6\text{S}$ showed $M^{+H}_{\text{found}} = 558$ ($M^{+H}_{\text{calc}} = 558$).

Example 52 : Preparation of tetrahydro-N-hydroxy-4-
[[4-(phenylmethyl)-1-piperazinyl]-
sulfonyl]-2H-pyran-4-carboxamide,
monohydrochloride



20

Part A: To a cooled solution, zero C, of N-benzylpiperazine (Aldrich, 10.0 g, 56.7 mmol) in dichloromethane (75 mL) was added triethylamine (Aldrich, 11.9 mL, 85.1 mmol), followed by the slow addition of a solution of methane sulfonyl chloride (Aldrich, 6.6 mL, 85.1 mmol) in dichloromethane (25 mL). After the addition, the ice bath was removed and the reaction stirred at ambient temperature for

-550-

1.5 hours. Once completed, the solvent was evaporated and the residue was taken up in ethyl acetate (200 mL) and water (100 mL). The phases were separated and the aqueous phase was treated with 1 N NaOH_{aq} (100 mL), then extracted with ethyl acetate (2x-200 mL). The combined organic layers were washed with 5% KHSO₄ _{aq} (2x- 100 mL) and brine (1x- 100 mL). The ethyl acetate was then dried over Na₂SO₄ and concentrated to afford the piperazine mesylate, as an orange solid (13.0 g, 90% yield). ¹H NMR showed the desired compound.

Part B: Oven-dried glassware was charged with the piperazine mesylate of Part A (12.6 g, 50 mmol) and tetrahydrofuran (160 mL) and cooled to -75°C. Lithium bis(trimethylsilyl)amide (Aldrich, 1.0 M in tetrahydrofuran, 165 mL, 165 mmol) was slowly added, keeping temperature at less than -60°C. The reaction was stirred for 30 minutes after the addition and was then charged with a solution of methyl chloroformate (Aldrich, 3.9 mL, 94.5 mmol) in tetrahydrofuran (80 mL) again keeping the temperature at less than -60°C. After stirring for 1 hour at -75°C, the reaction was quenched with saturated NH₄Cl_{aq}, keeping temperature below -20°C. The aqueous portion froze into a solid chunk of ice. After warming to 5°C, the mixture was extracted with ethyl acetate (3x- 200 mL). Organics were washed with saturated NH₄Cl_{aq} (2x-100 mL) and brine (1x-100ml), then dried over Na₂SO₄ and concentrated to afford the methylene ester as a brown oil (19.6 g, quantitative yield). ¹H NMR showed the desired compound.

Part C: To a solution of the methylene ester of Part B (10.0 g, 32.0 mmol), potassium carbonate

(Aldrich, 15.5 g, 112 mmol), and 18-crown-6 (Aldrich, 0.5 g, 2.0 mmol) in dimethylformamide (60 mL) was added dibromo-diethylether (Lancaster, 4.4 mL, 35.3 mmol). The mixture was heated at 60°C for 18 hours, after which more potassium carbonate (1.5 g, 12 mmol) was added, and the reaction continued at 60°C for 4 hours. The reaction was worked up by removing the solvent by roto-evaporation. The residue was taken up in ethyl acetate (250 mL) and water (100 mL). The layers were separated and the organic was washed with water (1x-100 mL) and brine (2x- 100 mL) then dried over Na₂SO₄ and concentrated to afford a black oil. The oil was purified via silica gel (ethyl acetate/hexanes) yielding the methyl ester as a yellow oil (7.3 g, 60% yield). ¹H NMR showed the desired compound.

Part D: To a solution of the methyl ester of Part C (3.5 g, 9.2 mmol) in tetrahydrofuran (20 mL) was added potassium trimethylsilonate (Aldrich, 3.5 g, 27.4 mmol). The reaction was stirred overnight (about 18 hours) at ambient temperature. Work up comprised removing the tetrahydrofuran and taking the residue up in H₂O (8 mL), cooling to zero°C, and titrating to pH 7 with 6 N HCl. Solids formed and were collected and washed with water followed by diethyl ether. The solid was dried in vacuo to afford the carboxylic acid as an off white solid (0.75 g, 22% yield). ¹H NMR showed the desired compound.

Part E: To a solution of the carboxylic acid of Part D (0.75 g, 2.0 mmol) in dimethylacetamide (5 mL) was added N-methylmorpholine (Aldrich, 0.66 mL, 6.0 mmol) followed by N-hydroxybenzotriazole hydrate

-552-


(Aldrich, 0.32 g, 2.4 mmol), O-(tetrahydro-2H-pyran-2-yl) hydroxylamine (0.35 g, 3.0 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (Sigma, 0.57 g, 3.0 mmol). The mixture
5 was stirred at 40 C for thirteen hours and was then stripped of solvent. The residue was purified by reverse phase chromatography (carbon-18, acetonitrile/water). The product was hydrolyzed on the column giving the hydroxamate as a white solid
10 (0.54 g, 57% yield). ^1H and ^{19}F NMR showed the desired compound but as the trifluoroacetic acid salt.

Part F: The hydroxamate of Part E (0.5 g, 1.0 mmol) was dissolved in 1,4-dioxane (3.0 mL) and stirred with 4 N HCl in dioxane (4 mL) for one hour.
15 The solvent volume was reduced in half then diethyl ether was added, providing a solid that was filtered and washed with excess diethyl ether. The solid was dried to afford the hydrochloride salt as a white solid (0.4 g, 100% yield). ^1H NMR showed the desired
20 compound. HRMS for $\text{C}_{17}\text{H}_{25}\text{F}_3\text{N}_3\text{O}_5\text{S}$ showed $M^{+H}_{\text{found}} = 384$ ($M^{+H}_{\text{calc}} = 384$).

Example 53: Preparation of N-hydroxy-1-

(phenylmethyl)-4-[(4-phenyl-1-piperazinyl)sulfonyl]-4-piperidine-
25 carboxamide, bis(trifluoroacetate)

-553-



Part A: To a cooled solution, zero C, of N-phenylpiperazine (Aldrich, 10.0 g, 61.6 mmol) in dichloromethane (75 mL) was added triethylamine (Aldrich, 12.9 mL, 92.4 mmol) followed by the slow addition of a solution of methane sulfonyl chloride (Aldrich, 7.1 mL, 92.4 mmol) in dichloromethane (25 mL). After the addition, the ice bath was removed and the reaction stirred at ambient temperature for 1.5 hours. Once completed, the solvent was evaporated and the residue was taken up in ethyl acetate (200 mL) and water (100 mL). The phases were separated and the aqueous phase was treated with 1 N NaOH_{aq} (100 mL) then extracted with ethyl acetate (2x- 200 mL). The combined organic layers were washed with 5% KHSO₄ _{aq} (2x- 100 mL) and brine (1x- 100 mL). The ethyl acetate was then dried over Na₂SO₄ and concentrated to afford the piperazine mesylate, SC 80658, as solid (13.0 g, 88% yield). ¹H NMR showed the desired compound.

Part B: Oven-dried glassware was charged with the piperazine mesylate of Part A (12.6 g, 52.4 mmol) and tetrahydrofuran (160 mL) and cooled to -75°C. Lithium bis(trimethylsilyl)amide (Aldrich, 1.0 M in tetrahydrofuran, 184 mL, 184 mmol) was slowly added, keeping temperature below -60°C. The reaction was stirred for 30 minutes after the addition and was

then charged with a solution of methyl chloroformate (Aldrich, 4.1 mL, 52.4 mmol) in tetrahydrofuran (80 mL) again keeping the temperature below -60°C. After stirring for 1 hour at -75°C, the reaction was

5 quenched with saturated $\text{NH}_4\text{Cl}_{\text{aq}}$, keeping temperature below -20°C. The aqueous does freeze into a solid chunk of ice. After warming to 5°C, the mixture was extracted via ethyl acetate (3x- 200 mL). Organics were washed with saturated $\text{NH}_4\text{Cl}_{\text{aq}}$ (2x-100 mL) and

10 brine (1x-100ml), then dried over Na_2SO_4 and concentrated to afford the methylene ester, 9091-195, as a brown oil (14.1 g, 90%) ^1H NMR showed the desired compound.

Part C: To a solution of the methylene ester of

15 Part B (4.3 g, 14.4 mmol), potassium carbonate (Aldrich, 6.0 g, 43.2 mmol), and 18-crown-6 (Aldrich, 0.5 g, 2.0 mmol) in dimethylformamide (30 mL) was added N,N-bis(2-chloroethyl)-benzylamine, SC 9275A, (3.5 mL, 15.1 mmol). The mixture was heated at 60°C

20 for 18 hours. The reaction was worked up by removing the solvent by roto-evaporation. The residue was taken up in ethyl acetate (250 mL) and washed with water (1x-100 mL) and brine (2x- 100 mL) then dried over Na_2SO_4 and concentrated to afford an orange

25 solid. The solid was purified via silica gel (ethyl acetate/hexanes) yielding the methyl ester as a yellow solid (5.6 g, 85% yield). ^1H NMR showed the desired compound.

Part D: To a solution of the methyl ester of

30 Part C (2.0 g, 4.4 mmol) in tetrahydrofuran (10 mL) was added potassium trimethylsilonate (Aldrich, 1.7 g, 13.2 mmol). The reaction was stirred overnight (about 18 hours) at ambient temperature. Work up

comprised removing the tetrahydrofuran and taking the residue up in H₂O (8 mL), cooling to zero°C, and titrating to pH 6 with 6 N HCl. Solids formed and were collected, then washed with water followed by diethyl ether. The solid was dried in vacuo to afford the carboxylic acid as an off white solid (1.6 g, 84% yield). ¹H NMR showed the desired compound.


Part E: To a solution of the carboxylic acid of Part D (1.6 g, 3.6 mmol) in dimethylacetamide (8 mL) was added N-methylmorpholine (Aldrich, 1.2 mL, 10.8 mmol) followed by N-hydroxybenzotriazole hydrate (Aldrich, 0.58 g, 4.3 mmol), O-(tetrahydro-2H-pyran-2-yl) hydroxylamine (0.84 g, 7.2 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (Sigma, 1.0 g, 5.4 mmol). The mixture stirred overnight at ambient temperature and was then stripped of solvent. The residue was taken up in ethyl acetate (100 mL) then washed with aqueous saturated sodium bicarbonate (2x-50 mL) and brine (3x-50 mL). The organic was dried over Na₂SO₄ concentrated to afford the tetrahydropyran-protected hydroxamate as an oily foam (1.9 g, 95% yield). ¹H NMR showed the desired compound.

Part F: The tetrahydropyran-protected hydroxamate of Part E (1.6 g, 2.9 mmol) was dissolved in 1,4-dioxane (10.0 mL) and stirred with 4 N HCl in dioxane (12 mL) for 30 minutes. A solid formed that was filtered, washed with diethyl ether and dried. The solid was then purified by reverse phase chromatography (acetonitrile/water) affording the hydroxamate as a tan solid (1.4 g, 70% yield). ¹H

-556-

and ^{19}F NMR showed the desired compound. HRMS for $\text{C}_{23}\text{H}_{30}\text{N}_4\text{O}_4\text{S}$ showed $M^{\text{H}}_{\text{found}} = 439$ ($M^{\text{H}}_{\text{calc}} = 439$).

5 Example 54: Preparation of N-hydroxy-1-(phenylmethyl)-4-[(4-phenyl-1-piperazinyl)sulfonyl]-4-piperidine-carboxamide, dihydrochloride




10

Part A: The hydroxamate of Example 53 was dissolved in water (10 mL) and titrated to pH 10 via 1.0 N NaOH. The mixture was extracted with ethyl acetate (4x-20 mL). The organics were combined, 15 dried over Na_2SO_4 , and concentrated to afford a foamy solid. This solid was dissolved in acetonitrile (5 mL) and then concentrated HCl (1 mL) was dripped in slowly. The mixture was stirred for ten minutes and then was concentrated to a tan oil. The oil was 20 triturated with diethyl ether to form a solid that was dried in vacuo to afford the hydrochloride hydroxamate as a yellow solid (0.82 g, 53 % yield). ^1H and ^{19}F NMR showed the desired compound. HRMS for $\text{C}_{23}\text{H}_{30}\text{N}_4\text{O}_4\text{S}$ showed $M^{\text{H}}_{\text{found}} = 439$ ($M^{\text{H}}_{\text{calc}} = 439$).

25

Example 55: Preparation of 4-[[4-(4-butoxy-3-methylphenyl)-1-piperazinyl]sulfonyl]-

tetrahydro-N-hydroxy-2H-pyran-4-
carboxamide



5

Part A: To a solution of 4-bromo-2-methylphenol (Transworld, 10.0 g, 53.5 mmol) in acetone (105 mL) was added potassium carbonate (Aldrich, 16.6 g, 120 mmol) followed by 1-iodobutane (7.3 mL, 64.2 mmol). The reaction stirred at reflux for 12 hours. After cooling to ambient temperature, the mixture was filtered through a Celite® pad. The filtrate was concentrated giving a yellow oil which was purified on silica gel (ethyl acetate/hexanes) to afford the bromophenyl ether, SC 83965, as a clear oil (10.8 g, 83 % yield). ¹H NMR showed the desired compound. HRMS for C₁₁H₁₅BrO showed M⁺_{found} = 244 (M⁺_{calc} = 244).

Part B: To a solution of the bromophenyl ether of Part A (10.0 g, 41.1 mmol) in toluene (80 mL) was added 1-tert-butylcarbonylpiperazine (Lancaster, 9.2 g, 49.4 mmol) and sodium tert-butoxide (Fluka, 5.5 g, 57.5 mmol). The reaction stirred at ambient temperature for twenty minutes. BINAP (Aldrich, 0.8 g, 1.2 mmol) and tris(dibenzylideneacetone)dipalladium (0) (Aldrich, 0.4 g, 0.4 mmol) were then added and the reaction was stirred at 80°C until the bromide was exhausted. Work up comprised cooling the mixture to ambient temperature, filtering through a Celite® pad,

and concentrating the filtrate. The resulting residue was purified on silica gel (ethyl acetate/hexanes) to afford the BOC-piperazine as a black oil (6.4 g, 44% yield). ¹H NMR showed the
5 desired compound.

Part C: The oil of the BOC-piperazine of Part B (6.4 g, 18.4 mmol) was stirred with 4 N HCl in dioxane 23 mL, 92 mmol) for twenty minutes during which a solid formed. The solid was filtered and
10 washed with diethyl ether and dried affording the phenylpiperazine as an off white solid, which was recrystallized from methanol to yield a white solid (3.6 g, 69% yield). ¹H NMR showed the desired compound.

15 Part D: To a cooled solution, zero C, of the phenylpiperazine of Part C (3.5 g, 12.3 mmol) in dichloromethane (15 mL) was added triethylamine (Aldrich, 4.3 mL, 30.8 mmol), followed by the slow addition of a solution of methane sulfonyl chloride
20 (Aldrich, 1.4 mL, 18.4 mmol) in dichloromethane (10 mL). After the addition, the ice bath was removed and the reaction stirred at ambient temperature for 3 hours. Once completed, the solvent was evaporated and the residue was taken up in ethyl acetate (200
25 mL) and water (100 mL). The phases were separated and the aqueous was treated with 1 N NaOH_{aq} (100 mL) then extracted with ethyl acetate (2x-200 mL). The combined organic layers were washed with water (1x-100 mL) and brine (1x- 100 mL). The ethyl acetate
30 layer was then dried over Na₂SO₄ and concentrated to afford a brown oil that was purified on silica gel (ethyl acetate/hexanes) to give the piperazine

mesylate as a tan solid (3.1 g, 78% yield). ^1H NMR showed the desired compound.

Part E: Oven-dried glassware was charged with the piperazine mesylate of Part D (3.1 g, 9.5 mmol) and tetrahydrofuran (20 mL) and cooled to -75°C . Lithium bis(trimethylsilyl)amide (Aldrich, 1.0 M in tetrahydrofuran, 28.5 mL, 28.5 mmol) was slowly added, keeping temperature below -60°C . The reaction was stirred for 30 minutes after the addition and was then charged with a solution of methyl chloroformate (Aldrich, 0.8 mL, 10 mmol) in tetrahydrofuran (10 mL) again keeping the temperature below -60°C . After stirring for 1 hour at -75°C , the reaction was quenched with saturated $\text{NH}_4\text{Cl}_{\text{aq}}$, keeping temperature less than -20°C . The aqueous phase froze into a solid chunk of ice. After warming to 5°C , the mixture was extracted via ethyl acetate (3x- 100 mL). Organics were washed with saturated $\text{NH}_4\text{Cl}_{\text{aq}}$ (2x-100 mL) and brine (1x-100mL), then dried over Na_2SO_4 and concentrated to give a brown solid that was recrystallized from methanol to afford methylene ester as a yellow solid (1.3 g, 36%) ^1H NMR showed the desired compound.

Part F: To a solution of the methylene ester of Part E (1.3 g, 3.4 mmol), potassium carbonate (Aldrich, 1.4 g, 10.2 mmol), and 18-crown-6 (Aldrich, 0.08 g, 8 mmol) in N,N-dimethylformamide (10 mL) was added dibromo-diethylether (Lancaster, 0.5 mL, 3.6 mmol). The mixture was heated at 60°C for 18 hours and then worked up by removing the solvent by roto-evaporation. The residue was dissolved in ethyl acetate (250 mL) and water (100 mL). The layers were separated and the aqueous was extracted with ethyl

acetate (2x-100 mL). The organics were combined and washed with 5% HCl_{aq} (1x- 50 mL), water (1x-100 mL), and brine (2x- 100 mL) then dried over Na₂SO₄ and concentrated to afford a brown oil that was washed
5 with hexanes to afford the methyl ester as an oil (1.5 g, quantitative yield). ¹H NMR showed the desired compound.

Part G: To a solution of the methyl ester of Part F (1.5 g, 3.3 mmol) in tetrahydrofuran (10 mL)
10 was added potassium trimethylsilonate (Aldrich, 1.3 g, 9.9 mmol). The was stirred overnight (about 18 hours) at ambient temperature. Work up comprised removing the tetrahydrofuran and taking the residue up in H₂O (8 mL). The aqueous was washed with diethyl
15 ether, which resulted in an emulsion. The emulsion was filtered affording a gummy solid that was slurried in acetone to give the carboxylic acid as a white powder (0.71 g, 51% yield). ¹H NMR showed the desired compound.

20 Part H: To a solution of the carboxylic acid of Part G (0.7 g, 1.6 mmol) in N,N-dimethylformamide (5 mL) was added N-methylmorpholine (Aldrich, 0.5 mL, 4.8 mmol), followed by N-hydroxybenzotriazole hydrate (Aldrich, 0.27 g, 2.0 mmol), O-(tetrahydro-
25 2H-pyran-2-yl) hydroxylamine (0.37 g, 3.2 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (Sigma, 0.46 g, 2.4 mmol). The mixture was stirred at 40 C for 8 hours, and was then stripped of solvent. The residue was purified by reverse
30 phase chromatography (carbon-18, acetonitrile/water). After removing the acetonitrile from the desired fractions by roto-evaporation, the aqueous layer was extracted with ethyl acetate (2x-100 mL). The

-561-

organics were dried over Na_2SO_4 and concentrated to afford the tetrahydropyran(THP)-protected hydroxamate as a white solid (0.57 g, 66% yield). ^1H NMR showed the desired compound.

5 Part F: To the tetrahydropyran-protected hydroxamate of Part E (0.6 g, 1.0 mmol) was added methanol (0.2 mL) and 4 N HCl in dioxane (2.5 mL) and the mixture was stirred for one hour. The solvent was stripped and the residue was slurried in diethyl
10 ether to provide a solid that was filtered and washed with excess diethyl ether. The solid was dried to afford the hydroxamate as a white solid (0.12 g, 26% yield). ^1H NMR showed the desired compound. HRMS for $\text{C}_{21}\text{H}_{33}\text{N}_3\text{O}_6\text{S}$ showed $M^{+H}_{\text{found}} = 456$ ($M^{+H}_{\text{calc}} = 456$).

15

Example 56: In Vitro Metalloprotease Inhibition

The compounds prepared in the manner described in the Examples above were assayed for activity by an in vitro assay. Following the
20 procedures of Knight et al., *FEBS Lett.* **296(3)**:263 (1992). Briefly, 4-aminophenylmercuric acetate (APMA) or trypsin-activated MMPs were incubated with various concentrations of the inhibitor compound at room temperature for 5 minutes.

25 More specifically, recombinant human MMP-13 and MMP-1 enzymes were prepared in laboratories of the assignee following usual laboratory procedures. MMP-13 from a full length cDNA clone was expressed as a proenzyme using a baculovirus as discussed in V.A. Luckow, *Insect Cell Expression Technology*, pages 183-
30 218, in Protein Engineering: Principles and Practice, J.L.Cleland et al eds., Wiley-Liss, Inc., (1996). See, also, Luckow et al., *J. Virol.*, **67**:4566-4579

(1993); O'Reilly et al., Baculovirus Expression Vectors: A Laboratory Manual, W.H. Freeman and Company, New York, (1992); and King et al., The Baculovirus Expression System: A Laboratory Guide, Chapman & Hall, London (1992) for further details on use of baculovirus expression systems. The expressed enzyme was purified first over a heparin agarose column and then over a chelating zinc chloride column. The proenzyme was activated by APMA for use in the assay.

MMP-1 expressed in transfected HT-1080 cells was provided by Dr. Harold Welgus of Washington University, St. Louis, MO. The enzyme was also activated using APMA and was then purified over a hydroxamic acid column. Further specifics for preparation and use of these enzymes can be found in the scientific literature describing these enzymes. See, for example, Enzyme Nomenclature, Academic Press, San Diego, Ca (1992) and the citations therein, and Frije et al., J. Biol. Chem., **269**(24): 16766-16773 (1994).

The enzyme substrate is a methoxycoumarin-containing polypeptide having the following sequence:

MCA-ProLeuGlyLeuDpaAlaArgNH₂, wherein MCA is methoxycoumarin and Dpa is 3-(2,4-dinitrophenyl)-L-2,3-diaminopropionyl alanine. This substrate is commercially available from Baychem as product M-1895.

The buffer used for assays contained 100 mM Tris-HCl, 100 mM NaCl, 10 mM CaCl₂ and 0.05 percent polyethyleneglycol (23) lauryl ether at a pH value of 7.5. Assays were carried out at room temperature,

and dimethyl sulfoxide (DMSO) at a final concentration of 1 percent was used to dissolve inhibitor compound.

The assayed inhibitor compound in
5 DMSO/buffer solution was compared to an equal amount of DMSO/buffer with no inhibitor as control using Microfluor™ White Plates (Dynatech). The inhibitor or control solution was maintained in the plate for 10 minutes and the substrate was added to provide a
10 final concentration of 4 μ M.

In the absence of inhibitor activity, a fluorogenic peptide was cleaved at the gly-leu peptide bond, separating the highly fluorogenic peptide from a 2,4-dinitrophenyl quencher, resulting
15 in an increase of fluorescence intensity (excitation at 328 nm/emission at 415 nm). Inhibition was measured as a reduction in fluorescent intensity as a function of inhibitor concentration, using a Perkin Elmer L550 plate reader. The IC₅₀ values were
20 calculated from those values. The results are set forth in the Inhibition Table below, reported in terms of IC₅₀ to three significant figures, where appropriate.

Inhibition Table

5

ED₅₀ nM

Example	MMP-1	MMP-2	MMP-3	MMP-7	MMP-8	MMP-9	MMP-13	MMP-14
1	<10,000	7.0	-	-	-	-	20.0	-
2	4,080	<0.1	94.0	9,000	1.9	1.8	0.3	80.0
3	7,300	<0.1	290	-	3.2	3.0	0.9	47.0
4	<10,000	9.0	400	-	285	37.3	11.4	8,000
5	<10,000	46.4	-	-	-	-	64	-
6	6,400	0.2	50.0	-	1.0	0.7	0.6	73.0
7	5,500	3,000	-	-	-	-	9,000	-
8	6,000	2.0	120	-	3.6	15.8	4.0	55.3
9	<10,000	0.7	186	-	3.0	2.0	0.9	200
10	5,500	0.35	175	-	2.0	18.5	1.4	500
11	<10,000	2.7	2,000	-	8.8	30.0	8.0	900
12	7,000 4,500	0.1 <0.1	42.5	<10,000	0.8	1.1	0.6 0.25 <0.1	80.0
13	1,100	0.2	-	-	-	-	0.7	-
14	<10,000	<0.1 0.4 0.1	250	<10,000	6.7	0.7	<0.1 0.6 0.5	150
15	8,000 3,000	0.15 <0.1	23.5	<10,000	2.4 2.0	0.19	0.6 <0.1	67.5
16	<10,000	1.4	-	-	-	-	8.0	-
17	2,200 3,000	0.1 <0.1	23.5	-	1.9 1.6	0.7	0.1 <0.1	45.4
18	<10,000	0.7	160	-	2.2	1.0	0.8	145
19	2,800	-	30.6	<10,000	2.5	0.5	0.7	32.7
20	-	-	-	-	-	-	-	-
21	-	-	-	-	-	-	-	-
22	-	-	-	-	-	-	-	-
23	-	0.1	-	-	12.0	-	0.4	-
24	-	-	-	-	-	-	-	-
25	-	-	-	-	-	-	-	-
26	<10,000	<10,000	-	-	-	-	2,000	-

-565-


Example	MMP-1	MMP-2	MMP-3	MMP-7	MMP-8	MMP-9	MMP-13	MMP-14
27	<10,000	5,300	-	-	-	-	670	-
28	<10,000	0.3	270	-	4.3	1.8	1.0	360
29	-	-	-	-	-	-	-	-
30	4,000	0.3	28.6	-	1.6	1.0	0.9	45.5
31	<10,000	1.3	100	-	22.0	100	0.8	2,100
32	<10,000	<0.1	-	-	-	-	<0.1	-
33	2,400	4.4	-	-	-	-	22.2	-
34	<10,000	27.0	-	-	-	-	200	-
35	<10,000	<0.1	210	-	14.8	0.6	<0.1	540
36	<10,000	0.2	-	-	-	-	1.4	-
37	<10,000	300	-	-	-	-	190	-
38	<10,000	3.0	66.4	-	136	2.7	0.8	<10,000
39	<10,000	3,700	-	-	-	-	290	-
40	7,300	8.3	-	-	-	-	13.9	-
41	<10,000	5.0	-	-	-	-	3.7	-
42	5,300	0.1	37.3	<10,000	1.6	1.3	0.3	40.0
43	6,000	0.2	165	-	3.4	1.0	1.4	220
44	7,700	0.2	-	-	-	-	0.7	-
45	-	-	-	-	-	-	-	-
46	<10,000	0.25	-	-	-	-	0.6	-
47	<10,000	<0.1	46.9	<10,000	1.8 1.6	0.48	<0.1	83.1
48	370	200	-	-	-	-	42.5	-
49	<10,000	<10,000	-	-	-	-	<10,000	-
50	<10,000	0.4	-	-	-	-	3.0	-
51	<10,000	<0.1	3,500	-	13.0	9.0	0.3 <0.1	9,000
52	<10,000	115	-	-	-	-	195	-
53	3,500	4.4	-	-	-	-	13.5	-
54	6,000	12.8	-	-	-	-	33.0	-
55	<10,000	2.7	-	-	-	-	4.2	-

From the foregoing, it will be observed that numerous modifications and variations can be effected without departing from the true spirit and scope of the novel concepts of the present invention.

- 5 It is to be understood that no limitation with respect to the specific example presented is intended or should be inferred. The disclosure is intended to cover by the appended claims all such modifications as fall within the scope of the claims.

What is Claimed:

1. A compound corresponding in structure to formula II, below, or a pharmaceutically acceptable salt thereof:



wherein

R¹ and R² taken together with the carbon to which they are bonded form a heterocyclo or cycloalkyl group optionally substituted by one, two or three R^x substituents, or R¹ and R² are independently selected from the group consisting of:

hydrido,

an alkyl group, optionally substituted with one, two or three groups independently selected from R^x substituents,

an alkyloxyalkyl group, optionally substituted with one, two or three groups independently selected from R^x substituents,

an alkylthioalkyl group, optionally substituted with one, two or three groups

independently selected from R^x substituents,

an alkenyl group, optionally substituted with one, two or three groups independently selected from R^x substituents,

an alkynyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

an aryl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

an arylalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

an arylalkyloxyalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

an aryloxyalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

an arylthioalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

an arylalkylthioalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

a cycloalkyl or bicycloalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

a cycloalkenyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

a cycloalkylalkyl or bicycloalkylalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

a cycloalkyloxyalkyl or
bicycloalkyloxyalkyl group, optionally substituted
with one, two or three groups independently selected
from R^X substituents,

5 a cycloalkylalkyloxyalkyl or
bicycloalkyloxyalkyl group, optionally substituted
with one, two or three groups independently selected
from R^X substituents,

 a cycloalkylthioalkyl or
10 bicycloalkylthioalkyl group, optionally substituted
with one, two or three groups independently selected
from R^X;

 cycloalkylalkylthioalkyl or
bicycloalkylalkylthioalkyl, optionally substituted
15 with one, two or three groups independently selected
from R^X substituents,

 a heterocyclo group, optionally substituted
with one, two or three groups independently selected
from R^X substituents,

20 a heterocycloalkyl group, optionally
substituted with one, two or three groups
independently selected from R^X substituents

 a heteroaryl group, optionally substituted
with one, two or three groups independently selected
25 from R^X substituents,

 a biarylalkyl group, optionally substituted
with one, two or three groups independently selected
from R^X substituents,

 an arylalkenyl group, optionally
30 substituted with one, two or three groups
independently selected from R^X substituents,

- an arylalkynyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,
- a heterocycloalkylthio group, optionally substituted with one, two or three groups selected independently from R^X substituents,
- a heterocycloalkyloxyalkyl group, optionally substituted with one, two or three groups selected independently from R^X substituents,
- 10 a heteroarylalkenyl group, optionally substituted with one, two or three groups independently selected from R^X substituents, and
- a heteroarylalkyloxyalkyl group, optionally substituted with one, two or three groups
- 15 independently selected from R^X substituents;
- wherein an R^X substituent is selected from the group aminoalkyl, nitro, nitroso, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, alkoxy, aryloxy, heteroaryloxy, alkenyloxy, alkynyloxy, alkoxyaryl,
- 20 alkoxyheteroaryl, alkoxyalkyl, R^C -oxyalkyl, alkoxyalkyl, alkylenedioxy, aryloxyalkyl, perfluoroalkyl, trifluoroalkyl, alkylthio, arylthio, alkyloxycarbonyl, alkyloxycarbonyloxy, aryloxycarbonyl, arylalkyloxycarbonyl,
- 25 alkyloxycarbonyl- R^C -amino, arylalkyloxycarbonyl- R^C -amino, aryloxycarbonyloxy, carboxy, R^{Cd} -aminocarbonyloxy, R^{Cd} -aminocarbonyl, R^{Cd} -aminoalkanoyl, hydroxy- R^C -aminocarbonyl, R^{Cd} -aminosulfonyl, arylsulfonyl(R^C)amino, R^{Cd} -
- 30 aminoalkoxy, R^{Cd} -aminocarbonyl(R^C)amino, trifluoromethylsulfonyl(R^C)amino, heteroarylsulfonyl-

(R^C)amino, alkylsulfonyl, arylsulfonyl(R^C)amino, arylsulfonyl(R^C)aminocarbonyl, alkylsulfonyl(R^C)amino, arylcarbonyl(R^C)-aminosulfonyl, and an alkylsulfonyl(R^C)aminocarbonyl substituent;

- 5 wherein R^C and R^d are independently selected from the group consisting of a hydrido, alkanoyl, arylalkyl, aroyl, bisalkoxyalkyl, alkyl, haloalkyl, perfluoroalkyl, trifluoromethylalkyl, perfluoroalkoxyalkyl, alkoxyalkyl, cycloalkyl, heterocycloalkyl, heterocycloalkylcarbonyl, aryl, heterocyclo, heteroaryl, cycloalkylalkyl, aryloxyalkyl, heteroaryloxyalkyl, heteroarylalkoxyalkyl, heteroarylthioalkyl, arylsulfonyl, alkylsulfonyl, heteroarylsulfonyl, carboxyalkyl, alkoxycarbonylalkyl, aminocarbonyl, 10 alkyliminocarbonyl, aryliminocarbonyl, heterocycloiminocarbonyl, arylthioalkyl, alkylthioalkyl, arylthioalkenyl, alkylthioalkenyl, heteroarylalkyl, haloalkanoyl, hydroxyalkanoyl, thiolalkanoyl, alkenyl, alkynyl, alkoxyalkyl, 20 alkoxycarbonyl, aryloxycarbonyl, aminoalkylcarbonyl, hydroxyalkyl, aminoalkyl, aminoalkylsulfonyl, aminosulfonyl wherein the amino nitrogen is (i) unsubstituted or (ii) independently substituted with one or two R^Y radicals, or the substituents on the amino group taken together with the amino nitrogen form a saturated or partially unsaturated heterocyclo group optionally substituted with one, two or three groups independently selected from R^W substituents or 30 a heteroaryl group optionally substituted with one,

two or three groups independently selected from R^V substituents;

wherein R^Y is selected from the group consisting of an arylalkyl, aryl, heteroaryl, heterocyclo, alkyl, alkynyl, alkenyl, alkoxyalkyl, alkoxyalkyl, substituted or unsubstituted aminoalkyl, alkyloxycarbonyl, arylalkyloxycarbonyl, carboxyalkyl, haloalkyl, alkanoyl, aroyl, substituted or unsubstituted aminoalkanoyl, halo alkanoyl and a hydroxyalkyl group, each of which groups is optionally substituted by one or two groups independently selected from R^U substituents as are the substituents of the substituted aminoalkyl and substituted aminoalkanoyl groups;

wherein R^V is selected from the group consisting of a hydrido, aryl, heteroaryl, heterocyclo, aroyl, alkanoyl, heteroaroyl, halogen, cyano, aldehydo, hydroxy, amino, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, alkoxy, aryloxy, heteroaryloxy, alkenyloxy, alkynyloxy, alkoxyaryl, alkoxyheteroaryl, R^YR^Z-amino, alkoxyalkyl, alkylendioxy, aryloxyalkyl, perfluoroalkyl, trifluoroalkyl, alkylthio, arylthio, alkyloxycarbonyl, alkyloxycarbonyloxy, aryloxycarbonyl, arylalkyloxycarbonyl, arylalkyloxycarbonylamino, aryloxycarbonyloxy, carboxy, R^YR^Z-aminocarbonyloxy, R^YR^Z-aminocarbonyl, R^YR^Z-aminoalkanoyl, hydroxyaminocarbonyl, R^YR^Z-aminosulfonyl, R^YR^Z-aminocarbonyl(R^Y) amino, trifluoromethylsulfonyl(R^Y) amino, heteroarylsulfonyl(R^Y) amino, arylsulfonyl(R^Y) amino, arylsulfonyl(R^Y) -

aminocarbonyl, alkylsulfonyl(RY)amino, arylcarbonyl-(RY)aminosulfonyl, and an alkylsulfonyl(RY)-aminocarbonyl substituent;

wherein R^W is selected from the group
5 consisting of a hydrido, aryl, heteroaryl, heterocyclo, aroyl, alkanoyl, heteroaroyl, halogen, cyano, aldehydo, hydroxy, amino, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, alkoxy, aryloxy, heteroaryloxy, alkenyloxy, alkynyloxy, alkoxyaryl,
10 alkoxyheteroaryl, R^YR^Z -amino, alkoxyalkyl, alkylenedioxy, aryloxyalkyl, perfluoroalkyl, trifluoroalkyl, alkylthio, arylthio, alkyloxycarbonyl, alkyloxycarbonyloxy, aryloxycarbonyl, arylalkyloxycarbonyl,
15 arylalkyloxycarbonylamino, aryloxycarbonyloxy, carboxy, R^YR^Z -aminocarbonyloxy, R^YR^Z -aminocarbonyl, R^YR^Z -aminoalkanoyl, hydroxyaminocarbonyl, R^YR^Z -aminosulfonyl, R^YR^Z -aminocarbonyl(RY)amino, trifluoromethylsulfonyl(RY)amino, heteroarylsulfonyl-
20 (RY)amino, arylsulfonyl(RY)amino, arylsulfonyl(RY)-aminocarbonyl, alkylsulfonyl(RY)amino, arylcarbonyl-(RY)aminosulfonyl, and an alkylsulfonyl(RY)-aminocarbonyl substituent;

R^Z is selected from the group consisting of
25 an arylalkyl, aryl, heteroaryl, heterocyclo, alkyl, alkynyl, alkenyl, alkoxyalkyl, alkoxyalkylalkyl, substituted or unsubstituted aminoalkyl, alkyloxycarbonyl, arylalkyloxycarbonyl, carboxyalkyl, haloalkyl, alkanoyl, aroyl, substituted or
30 unsubstituted aminoalkanoyl, halo alkanoyl and a

hydroxyalkyl group, each of which groups are optionally substituted by one or two R^U substituents;

wherein R^U is selected from the group consisting of an arylalkyl, aryl, heteroaryl, heterocyclo, alkyl, alkynyl, alkenyl, alkoxyalkyl, alkoxyalkyl, substituted or unsubstituted aminoalkyl, alkyloxycarbonyl, arylalkyloxycarbonyl, carboxyalkyl, haloalkyl, alkanoyl, aroyl, substituted or unsubstituted aminoalkanoyl, halo alkanoyl and a hydroxyalkyl group, wherein the substituents of the substituted aminoalkyl and substituted aminoalkanoyl groups are selected from the group consisting of an alkyl, alkenyl, alkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, aryloxycarbonyl and an alkyloxycarbonyl group;

R^{3a} and R^{3b} are independently selected from the group consisting of a hydrido, alkyl, alkenyl, alkynyl, aryl, heterocyclo, heteroaryl, cycloalkyl, and an alkoxyalkyl group, each of which groups is optionally substituted by an -AREY substituent;

in that AREY substituent, A is selected from the group consisting of

- (1) -O-;
- (2) -S-;
- (3) -N(R^e)-;
- (4) -CO-N(R^e) or -N(R^e)-CO-;
- (5) -CO-O- or -O-CO-;
- (6) -O-CO-O-;
- (7) -HC=CH-;
- (8) -NH-CO-NH-;
- (9) -C \equiv C-;
- (10) -NH-CO-O- or -O-CO-NH-;

- (11) -N=N-;
- (12) -NH-NH-;
- (13) -CS-N(R^e)- or -N(R^e)-CS-;
- (14) -CH₂-;
- 5 (15) -O-[(CH₂)₁₋₈]- or -[(CH₂)₁₋₈]O-; and
- (16) -S-CH₂- or -CH₂-S-; or
- (17) A is absent and R is directly connected to R^{3a} or R^{3b}, or both R^{3a} and R^{3b};
- 10 the moiety R is selected from the group consisting of alkyl, alkoxyalkyl, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroparalkyl, heterocycloalkyl, cycloalkylalkyl, cycloalkoxyalkyl, heterocycloalkoxyalkyl, aryloxyalkyl,
- 15 heteroaryloxyalkyl, arylthioalkyl, heteroarylthioalkyl, cycloalkylthioalkyl, and a heterocycloalkylthioalkyl group wherein the aryl, heteroaryl, cycloalkyl or heterocycloalkyl substituent is (i) unsubstituted or (ii) substituted
- 20 with one or two radicals selected from the group consisting of a halo, alkyl, perfluoroalkyl, perfluoroalkoxy, perfluoroalkylthio, trifluoromethylalkyl, amino, alkoxycarbonylalkyl, alkoxy, C₁-C₂-alkylene-dioxy, hydroxycarbonylalkyl,
- 25 hydroxycarbonylalkylamino, nitro, hydroxy, hydroxyalkyl, alkanoylamino, and a alkoxycarbonyl group;
- the group E is selected from the group consisting of
- 30 (1) -CO(R^w)- or -(R^w)CO-;
- (2) -CON(R^e)- or -(R^e)NCO-;

-576-

(3) -CO-;

(4) -SO₂-R^W or -R^WSO₂-;(5) -SO₂-;(6) -N(R^E)-SO₂- or -SO₂-N(R^E)-; or

5 (7) E is absent and R is bonded directly to Y; and

Y is absent or is selected from the group consisting of a hydrido, alkyl, alkoxy, haloalkyl, aryl, aralkyl, cycloalkyl, heteroaryl, hydroxy, nitrile, nitro, aryloxy, aralkoxy, heteroaryloxy, heteroaralkyl, R^Coxyalkyl, perfluoroalkoxy, perfluoroalkylthio, trifluoromethylalkyl, alkenyl, heterocycloalkyl, cycloalkyl, trifluoromethyl, alkoxy carbonyl, and an aminoalkyl group, wherein the

10 aryl, heteroaryl or heterocycloalkyl group is (i) unsubstituted or (ii) substituted with one or two radicals independently selected from the group consisting of an alkanoyl, halo, nitro, nitrile, haloalkyl, alkyl, aralkyl, aryl, alkoxy, and an amino

15 group wherein the amino nitrogen is (i) unsubstituted or (ii) substituted with one or two groups independently selected from hydrido, alkyl, and an aralkyl group;

20

wherein R^E is selected from hydrido, alkyl, alkenyl, alkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, aryloxy carbonyl, alkyloxy carbonyl, R^CR^damino carbonyl, R^CR^daminosulfonyl, R^CR^daminoalkanoyl and R^CR^daminoalkylsulfonyl, and R^C, R^d and R^W are as defined before; or

25

R^{3a} and R^{3b} taken together with the nitrogen atom to which they are bonded form a group -GAREY wherein

G is a N-heterocyclo group;

5 the substituent A is selected from the group consisting of

- (1) -O-;
- (2) -S-;
- (3) -NR^e-;
- 10 (4) -CO-N(R^e) or -N(R^e)-CO-;
- (5) -CO-O- or -O-CO-;
- (6) -O-CO-O-;
- (7) -HC=CH-;
- (8) -NH-CO-NH-;
- 15 (9) -C≡C-;
- (10) -NH-CO-O- or -O-CO-NH-;
- (11) -N=N-;
- (12) -NH-NH-;
- (13) -CS-N(R^e)- or -N(R^e)-CS-;
- 20 (14) -CH₂-;
- (15) -O-[(CH₂)₁₋₈]- or -[(CH₂)₁₋₈]O-; and
- (16) -S-CH₂- or -CH₂-S-; or
- (17) A is absent and R is directly bonded to G;

25 the moiety R is selected from the group consisting of alkyl, alkoxyalkyl, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaralkyl, heterocycloalkyl, cycloalkylalkyl, cycloalkoxyalkyl, heterocycloalkoxyalkyl, aryloxyalkyl, 30 heteroaryloxyalkyl, arylthioalkyl, heteroarylthioalkyl, cycloalkylthioalkyl, and a

heterocycloalkylthioalkyl group wherein the aryl or heteroaryl or cycloalkyl or heterocycloalkyl substituent is (i) unsubstituted or (ii) substituted with one or two radicals selected from the group consisting of a halo, alkyl, perfluoroalkyl, perfluoroalkoxy, perfluoroalkylthio, trifluoromethylalkyl, amino, alkoxycarbonylalkyl, alkoxy, C₁-C₂-alkylene-dioxy, hydroxycarbonylalkyl, hydroxycarbonylalkylamino, nitro, hydroxy, hydroxyalkyl, alkanoylamino, and a alkoxycarbonyl group;

the moiety E is selected from the group consisting of

- (1) -CO(R^W)- or -(R^W)CO-;
- (2) -CONH- or -HNCO-;
- (3) -CO-;
- (4) -SO₂-R^W- or -R^W-SO₂-;
- (5) -SO₂-;
- (6) -NH-SO₂- or -SO₂-NH-; or
- (7) E is absent and R is bonded directly to Y; and

the moiety Y is absent or is selected from the group consisting of a hydrido, alkyl, alkoxy, haloalkyl, aryl, aralkyl, cycloalkyl, heteroaryl, hydroxy, aryloxy, aralkoxy, heteroaryloxy, heteroaralkyl, perfluoroalkoxy, perfluoroalkylthio, trifluoromethylalkyl, alkenyl, heterocycloalkyl, cycloalkyl, trifluoromethyl, alkoxycarbonyl, and a aminoalkyl group, wherein the aryl or heteroaryl or heterocycloalkyl group is (i) unsubstituted or (ii) substituted with one or two radicals independently selected from the group consisting of an alkanoyl,


halo, nitro, aralkyl, aryl, alkoxy, and an amino group wherein the amino nitrogen is (i) unsubstituted or (ii) substituted with one or two groups independently selected from hydrido, alkyl, and an aralkyl group; and

R^{20} is (a) $-O-R^{21}$, where R^{21} is selected from the group consisting of a hydrido, C_1 - C_6 -alkyl, aryl, ar- C_1 - C_6 -alkyl group and a pharmaceutically acceptable cation, (b) $-NH-O-R^{22}$, wherein R^{22} is a selectively removable protecting group such as a 2-tetrahydropyranyl, benzyl, p-methoxybenzyl carbonyl- C_1 - C_6 -alkoxy, trisubstituted silyl group or o-nitrophenyl group, peptide synthesis resin and the like, wherein trisubstituted silyl group is substituted with C_1 - C_6 -alkyl, aryl, or ar- C_1 - C_6 -alkyl, or (c) $-NH-O-R^{14}$, where R^{14} is hydrido, a pharmaceutically acceptable cation or $C(W)R^{15}$ where W is O or S and R^{15} is selected from the group consisting of an C_1 - C_6 -alkyl, aryl, C_1 - C_6 -alkoxy, heteroaryl- C_1 - C_6 -alkyl, C_3 - C_8 -cycloalkyl- C_1 - C_6 -alkyl, aryloxy, ar- C_1 - C_6 -alkoxy, ar- C_1 - C_6 -alkyl, heteroaryl and amino C_1 - C_6 -alkyl group wherein the aminoalkyl nitrogen is (i) unsubstituted or (ii) substituted with one or two substituents independently selected from the group consisting of an C_1 - C_6 -alkyl, aryl, ar- C_1 - C_6 -alkyl, C_3 - C_8 -cycloalkyl- C_1 - C_6 -alkyl, ar- C_1 - C_6 -alkoxycarbonyl, C_1 - C_6 -alkoxycarbonyl, and C_1 - C_6 -alkanoyl radical, or (iii) wherein the amino C_1 - C_6 -alkyl nitrogen and two substituents attached thereto


-580-

form a 5- to 8-membered heterocycle or heteroaryl ring.

2. The compound or salt according to claim 1 wherein R^{21} is hydrido and said compound corresponds in structure to formula I, below.




3. The compound or salt according to claim 1 that corresponds in structure to formula III or formula IIIA, below,




- wherein substituents R^1 , R^2 , R^{20} and -GAREY are as before.

4. The compound or salt according to claim 1 that corresponds in structure to formula IV or formula IVA, below,




- wherein substituents R^1 , R^2 , R^{20} and -AREY are as before.

5. The compound or salt according to claim 1 that corresponds in structure to formula V or formula VA, below,




wherein substituents R^{20} and -AREY are as before.

6. The compound or salt according to claim 1 that corresponds in structure to formula VI or formula VIA, below,





15 wherein substituents R^1 , R^2 , R^{20} and -EY are as before defined, and A is -CH₂-, -O-CH₂-, -CH₂-O-, -S-CH₂- or -CH₂-S-.

7. The compound or salt according to claim 1 that corresponds in structure to formula VII or formula VIIA, below,



wherein substituents R^{20} and -EY are as before defined, and A is $-CH_2-$, $-O-CH_2-$, $-CH_2-O-$, $-S-CH_2-$ or $-CH_2-S-$.

5 8. The compound or salt according to claim
1 that corresponds in structure to formula VIII or
formula VIIIA, below,



10

wherein

R^{3a} , R^{3b} and R^{20} are as defined before; and

m is zero, 1 or 2;

n is zero, 1 or 2;

```
15      p is zero, 1 or 2;
```

the sum of $m + n + p = 1, 2, 3$ or 4 ;

(a) one of X, Y and Z is selected from the group consisting of C(O), NR⁶, O, S, S(O), S(O)₂ and

$\text{NS(O)}_2\text{R}^7$, and the remaining two of X, Y and Z are


20 CR^8R^9 , and $CR^{10}R^{11}$, or

(b) X and Z or Z and Y together constitute a moiety that is selected from the group consisting of $\text{NR}^6\text{C}(\text{O})$, $\text{NR}^6\text{S}(\text{O})$, $\text{NR}^6\text{S}(\text{O})_2$, NR^6S , NR^6O , SS , NR^6NR^6 and $\text{OC}(\text{O})$, with the remaining one of X, Y and Z being

25 CR⁸R⁹, or

-583-

(c) n is zero and X, Y and Z together constitute a moiety selected from the group consisting of



wherein wavy lines are bonds to the atoms
of the depicted ring;

R^6 and $R^{6'}$ are independently selected from the group consisting of hydrido, C_1 - C_6 -alkanoyl, C_6 -aryl- C_1 - C_6 -alkyl, aroyl, bis(C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl)- C_1 - C_6 -alkyl, C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl, C_1 -
 5 C_6 -perfluoroalkyl, C_1 - C_6 -trifluoromethylalkyl, C_1 - C_6 -perfluoroalkoxy- C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl, C_3 - C_6 -cycloalkyl, C_3 - C_8 -heterocycloalkyl, C_3 - C_8 -heterocycloalkylcarbonyl, C_6 -aryl, C_5 - C_6 -heterocyclo, C_5 - C_6 -heteroaryl, C_3 - C_8 -cycloalkyl- C_1 -
 10 C_6 -alkyl, C_6 -aryloxy- C_1 - C_6 -alkyl, heteroaryloxy- C_1 - C_6 -alkyl, heteroaryl- C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl, heteroarylthio- C_1 - C_6 -alkyl, C_6 -arylsulfonyl, C_1 - C_6 -alkylsulfonyl, C_5 - C_6 -heteroarylsulfonyl, carboxy- C_1 - C_6 -alkyl, C_1 - C_4 -alkoxycarbonyl- C_1 - C_6 -alkyl,
 15 aminocarbonyl, C_1 - C_6 -alkyliminocarbonyl, C_6 -aryliminocarbonyl, C_5 - C_6 -heterocycloiminocarbonyl, C_6 -arylthio- C_1 - C_6 -alkyl, C_1 - C_6 -alkylthio- C_1 - C_6 -alkyl, C_6 -arylthio- C_3 - C_6 -alkenyl, C_1 - C_4 -alkylthio- C_3 - C_6 -alkenyl, C_5 - C_6 -heteroaryl- C_1 - C_6 -alkyl, halo- C_1 - C_6 -
 20 alkanoyl, hydroxy- C_1 - C_6 -alkanoyl, thiol- C_1 - C_6 -alkanoyl, C_3 - C_6 -alkenyl, C_3 - C_6 -alkynyl, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_5 -alkoxycarbonyl, aryloxycarbonyl, NR^8R^9 - C_1 - C_5 -alkylcarbonyl, hydroxy- C_1 - C_5 -alkyl, an aminocarbonyl wherein the aminocarbonyl nitrogen is
 25 (i) unsubstituted or (ii) substituted with one or two radicals independently selected from the group consisting of C_1 - C_6 -alkyl, ar- C_1 - C_6 -alkyl, C_3 - C_8 -cycloalkyl and a C_1 - C_6 -alkanoyl group,

hydroxyaminocarbonyl, an aminosulfonyl group wherein the aminosulfonyl nitrogen is (i) unsubstituted or (ii) substituted with one or two radicals independently selected from the group consisting of C₁-C₆-alkyl, ar-C₁-C₆-alkyl, C₃-C₈-cycloalkyl and a C₁-C₆-alkanoyl group, an amino-C₁-C₆-alkylsulfonyl group wherein the amino-C₁-C₆-alkylsulfonyl nitrogen is (i) unsubstituted or (ii) substituted with one or two radicals independently selected from the group consisting of C₁-C₆-alkyl, ar-C₁-C₆-alkyl, C₃-C₈-cycloalkyl and a C₁-C₆-alkanoyl group and an amino-C₁-C₆-alkyl group wherein the aminoalkyl nitrogen is (i) unsubstituted or (ii) substituted with one or two radicals independently selected from the group consisting of C₁-C₆-alkyl, ar-C₁-C₆-alkyl, C₃-C₈-cycloalkyl and a C₁-C₆-alkanoyl group;

R⁷ is selected from the group consisting of a arylalkyl, aryl, heteroaryl, heterocyclo, C₁-C₆-alkyl, C₃-C₆-alkynyl, C₃-C₆-alkenyl, C₁-C₆-carboxyalkyl and a C₁-C₆-hydroxyalkyl group;

R⁸ and R⁹ and R¹⁰ and R¹¹ are independently selected from the group consisting of a hydrido, hydroxy, C₁-C₆-alkyl, aryl, ar-C₁-C₆-alkyl, heteroaryl, heteroar-C₁-C₆-alkyl, C₂-C₆-alkynyl, C₂-C₆-alkenyl, thiol-C₁-C₆-alkyl, C₁-C₆-alkylthio-C₁-C₆-alkyl cycloalkyl, cycloalkyl-C₁-C₆-alkyl, heterocycloalkyl-C₁-C₆-alkyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, aralkoxy-C₁-C₆-alkyl, C₁-C₆-alkoxy-C₁-C₆-alkoxy-C₁-C₆-alkyl, hydroxy-C₁-C₆-alkyl,

hydroxycarbonyl-C₁-C₆-alkyl, hydroxycarbonylar-C₁-C₆-alkyl, aminocarbonyl-C₁-C₆-alkyl, aryloxy-C₁-C₆-alkyl, heteroaryloxy-C₁-C₆-alkyl, arylthio-C₁-C₆-alkyl, heteroarylthio-C₁-C₆-alkyl, the sulfoxide or sulfone of any said thio substituents, perfluoro-C₁-C₆-alkyl, trifluoromethyl-C₁-C₆-alkyl, halo-C₁-C₆-alkyl, alkoxycarbonylamino-C₁-C₆-alkyl and an amino-C₁-C₆-alkyl group wherein the aminoalkyl nitrogen is (i) unsubstituted or (ii) substituted with one or two radicals independently selected from the group consisting of C₁-C₆-alkyl, ar-C₁-C₆-alkyl, cycloalkyl and C₁-C₆-alkanoyl, or wherein R⁸ and R⁹ or R¹⁰ and R¹¹ and the carbon to which they are bonded form a carbonyl group, or wherein R⁸ and R⁹ or R¹⁰ and R¹¹, or R⁸ and R¹⁰ together with the atoms to which they are bonded form a 5- to 8-membered carbocyclic ring, or a 5- to 8-membered heterocyclic ring containing one or two heteroatoms that are nitrogen, oxygen, or sulfur, with the proviso that only one of R⁸ and R⁹ or R¹⁰ and R¹¹ is hydroxy;


R¹² and R^{12'} are independently selected from the group consisting of a hydrido, C₁-C₆-alkyl, aryl, ar-C₁-C₆-alkyl, heteroaryl, heteroaralkyl, C₂-C₆-alkynyl, C₂-C₆-alkenyl, thiol-C₁-C₆-alkyl, cycloalkyl, cycloalkyl-C₁-C₆-alkyl, heterocycloalkyl-C₁-C₆-alkyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, aryloxy-C₁-C₆-alkyl, amino-C₁-C₆-alkyl, C₁-C₆-alkoxy-C₁-C₆-alkoxy-C₁-C₆-alkyl, hydroxy-C₁-C₆-alkyl, hydroxycarbonyl-C₁-

-587-

C_6 -alkyl, hydroxycarbonylar- C_1 - C_6 -alkyl,
 aminocarbonyl- C_1 - C_6 -alkyl, aryloxy- C_1 - C_6 -alkyl,
 heteroaryloxy- C_1 - C_6 -alkyl, C_1 - C_6 -alkylthio- C_1 - C_6 -
 alkyl, arylthio- C_1 - C_6 -alkyl, heteroarylthio- C_1 - C_6 -
 5 alkyl, the sulfoxide or sulfone of any said thio
 substituents, perfluoro- C_1 - C_6 -alkyl, trifluoromethyl-
 C_1 - C_6 -alkyl, halo- C_1 - C_6 -alkyl, alkoxycarbonylamino-
 C_1 - C_6 -alkyl and an amino- C_1 - C_6 -alkyl group wherein
 the aminoalkyl nitrogen is (i) unsubstituted or (ii)
 10 substituted with one or two radicals independently
 selected from the group consisting of C_1 - C_6 -alkyl,
 ar- C_1 - C_6 -alkyl, cycloalkyl and C_1 - C_6 -alkanoyl; and


R^{13} is selected from the group consisting
 of a hydrido, benzyl, phenyl, C_1 - C_6 -alkyl, C_2 - C_6 -
 15 alkynyl, C_2 - C_6 -alkenyl and a C_1 - C_6 -hydroxyalkyl
 group.

9. The compound or salt according to claim
 1 that corresponds in structure to the formula below
 20




10. The compound or salt according to
 claim 1 that corresponds in structure to the formula
 25 below


-588-




11. The compound or salt according to
claim 1 that corresponds in structure to the formula
5 below



12. The compound or salt according to
claim 1 that corresponds in structure to the formula
10 below




13. The compound or salt according to
claim 1 that corresponds in structure to the formula
15 below




14. The compound or salt according to
claim 1 that corresponds in structure to the formula
20 below

-589-




15. The compound or salt according to claim 1 that corresponds in structure to the formula below

5




16. The compound or salt according to claim 1 that corresponds in structure to the formula below

10




17. The compound or salt according to claim 1 that corresponds in structure to the formula below

15




-590-

18. The compound or salt according to claim 1 that corresponds in structure to the formula below



5


19. The compound or salt according to claim 1 that corresponds in structure to the formula below




10

20. A compound corresponding in structure to formulas XI or XIA, below, or a pharmaceutically acceptable salt thereof:

15



XI



XIA

wherein

20

m is zero, 1 or 2;

-591-

n is zero, 1 or 2;


p is zero, 1 or 2;

the sum of $m + n + p = 1, 2, 3$ or 4;


(a) one of X, Y and Z is selected from the
 5 group consisting of $C(O)$, NR^6 , O, S, $S(O)$, $S(O)_2$ and
 $NS(O)_2R^7$, and the remaining two of X, Y and Z are
 CR^8R^9 , and $CR^{10}R^{11}$, or

(b) X and Z or Z and Y together constitute
 a moiety that is selected from the group consisting
 10 of $NR^6C(O)$, $NR^6S(O)$, $NR^6S(O)_2$, NR^6S , NR^6O , SS, NR^6NR^6
 and $OC(O)$, with the remaining one of X, Y and Z being
 CR^8R^9 , or

(c) n is zero and X, Y and Z together
 constitute a moiety selected from the group
 15 consisting of



- 592 -



wherein wavy lines are bonds to the atoms of the depicted ring;

- 5 R⁶ and R^{6'} are independently selected from the group consisting of hydrido, C₁-C₆-alkanoyl, C₆-aryl-C₁-C₆-alkyl, aroyl, bis(C₁-C₆-alkoxy-C₁-C₆-alkyl)-C₁-C₆-alkyl, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-perfluoroalkyl, C₁-C₆-trifluoromethylalkyl, C₁-C₆-perfluoroalkoxy-C₁-C₆-alkyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₃-C₆-cycloalkyl, C₃-C₈-heterocycloalkyl, C₃-C₈-heterocycloalkylcarbonyl, C₆-aryl, C₅-C₆-heterocyclo, C₅-C₆-heteroaryl, C₃-C₈-cycloalkyl-C₁-C₆-alkyl, C₆-aryloxy-C₁-C₆-alkyl, heteroaryloxy-C₁-C₆-alkyl, heteroaryl-C₁-C₆-alkoxy-C₁-C₆-alkyl, heteroarylthio-C₁-C₆-alkyl, C₆-arylsulfonyl, C₁-C₆-alkylsulfonyl, C₅-C₆-heteroarylsulfonyl, carboxy-C₁-C₆-alkyl, C₁-C₄-alkoxycarbonyl-C₁-C₆-alkyl, aminocarbonyl, C₁-C₆-alkyliminocarbonyl, C₆-

aryliminocarbonyl, C₅-C₆-heterocycloiminocarbonyl,
C₆-arylthio-C₁-C₆-alkyl, C₁-C₆-alkylthio-C₁-C₆-alkyl,
C₆-arylthio-C₃-C₆-alkenyl, C₁-C₄-alkylthio-C₃-C₆-
alkenyl, C₅-C₆-heteroaryl-C₁-C₆-alkyl, halo-C₁-C₆-
5 alkanoyl, hydroxy-C₁-C₆-alkanoyl, thiol-C₁-C₆-
alkanoyl, C₃-C₆-alkenyl, C₃-C₆-alkynyl, C₁-C₄-alkoxy-
C₁-C₄-alkyl, C₁-C₅-alkoxycarbonyl, aryloxycarbonyl,
NR⁸R⁹-C₁-C₅-alkylcarbonyl, hydroxy-C₁-C₅-alkyl, an
aminocarbonyl wherein the aminocarbonyl nitrogen is
10 (i) unsubstituted or (ii) substituted with one or two
radicals independently selected from the group
consisting of C₁-C₆-alkyl, ar-C₁-C₆-alkyl, C₃-C₈-
cycloalkyl and a C₁-C₆-alkanoyl group,
hydroxyaminocarbonyl, an aminosulfonyl group wherein
15 the aminosulfonyl nitrogen is (i) unsubstituted or
(ii) substituted with one or two radicals
independently selected from the group consisting of
C₁-C₆-alkyl, ar-C₁-C₆-alkyl, C₃-C₈-cycloalkyl and a
C₁-C₆-alkanoyl group, an amino-C₁-C₆-alkylsulfonyl
20 group wherein the amino-C₁-C₆-alkylsulfonyl nitrogen
is (i) unsubstituted or (ii) substituted with one or
two radicals independently selected from the group
consisting of C₁-C₆-alkyl, ar-C₁-C₆-alkyl, C₃-C₈-
cycloalkyl and a C₁-C₆-alkanoyl group and an amino-
25 C₁-C₆-alkyl group wherein the aminoalkyl nitrogen is
(i) unsubstituted or (ii) substituted with one or two
radicals independently selected from the group
consisting of C₁-C₆-alkyl, ar-C₁-C₆-alkyl, C₃-C₈-
cycloalkyl and a C₁-C₆-alkanoyl group;

R⁷ is selected from the group consisting of a arylalkyl, aryl, heteroaryl, heterocyclo, C₁-C₆-alkyl, C₃-C₆-alkynyl, C₃-C₆-alkenyl, C₁-C₆-carboxyalkyl and a C₁-C₆-hydroxyalkyl group;

5 R⁸ and R⁹ and R¹⁰ and R¹¹ are independently selected from the group consisting of a hydrido, hydroxy, C₁-C₆-alkyl, aryl, ar-C₁-C₆-alkyl, heteroaryl, heteroar-C₁-C₆-alkyl, C₂-C₆-alkynyl, C₂-C₆-alkenyl, thiol-C₁-C₆-alkyl, C₁-C₆-alkylthio-C₁-C₆-
10 alkyl cycloalkyl, cycloalkyl-C₁-C₆-alkyl, heterocycloalkyl-C₁-C₆-alkyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, aralkoxy-C₁-C₆-alkyl, C₁-C₆-alkoxy-C₁-C₆-alkoxy-C₁-C₆-alkyl, hydroxy-C₁-C₆-alkyl, hydroxycarbonyl-C₁-C₆-alkyl, hydroxycarbonylar-C₁-C₆-
15 alkyl, aminocarbonyl-C₁-C₆-alkyl, aryloxy-C₁-C₆-alkyl, heteroaryloxy-C₁-C₆-alkyl, arylthio-C₁-C₆-alkyl, heteroarylthio-C₁-C₆-alkyl, the sulfoxide or sulfone of any said thio substituents, perfluoro-C₁-C₆-alkyl, trifluoromethyl-C₁-C₆-alkyl, halo-C₁-C₆-
20 alkyl, alkoxycarbonylamino-C₁-C₆-alkyl and an amino-C₁-C₆-alkyl group wherein the aminoalkyl nitrogen is (i) unsubstituted or (ii) substituted with one or two radicals independently selected from the group consisting of C₁-C₆-alkyl, ar-C₁-C₆-alkyl, cycloalkyl
25 and C₁-C₆-alkanoyl, or wherein R⁸ and R⁹ or R¹⁰ and R¹¹ and the carbon to which they are bonded form a carbonyl group, or wherein R⁸ and R⁹ or R¹⁰ and R¹¹, or R⁸ and R¹⁰ together with the atoms to which they

are bonded form a 5- to 8-membered carbocyclic ring, or a 5- to 8-membered heterocyclic ring containing one or two heteroatoms that are nitrogen, oxygen, or sulfur, with the proviso that only one of R⁸ and R⁹
5 or R¹⁰ and R¹¹ is hydroxy;

R¹² and R^{12'} are independently selected from the group consisting of a hydrido, C₁-C₆-alkyl, aryl, ar-C₁-C₆-alkyl, heteroaryl, heteroaralkyl, C₂-C₆-alkynyl, C₂-C₆-alkenyl, thiol-C₁-C₆-alkyl,
10 cycloalkyl, cycloalkyl-C₁-C₆-alkyl, heterocycloalkyl-C₁-C₆-alkyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, aryloxy-C₁-C₆-alkyl, amino-C₁-C₆-alkyl, C₁-C₆-alkoxy-C₁-C₆-alkoxy-C₁-C₆-alkyl, hydroxy-C₁-C₆-alkyl, hydroxycarbonyl-C₁-C₆-alkyl, hydroxycarbonylar-C₁-C₆-alkyl,
15 aminocarbonyl-C₁-C₆-alkyl, aryloxy-C₁-C₆-alkyl, heteroaryloxy-C₁-C₆-alkyl, C₁-C₆-alkylthio-C₁-C₆-alkyl, arylthio-C₁-C₆-alkyl, heteroarylthio-C₁-C₆-alkyl, the sulfoxide or sulfone of any said thio substituents, perfluoro-C₁-C₆-alkyl, trifluoromethyl-C₁-C₆-alkyl, halo-C₁-C₆-alkyl, alkoxycarbonylamino-C₁-C₆-alkyl and an amino-C₁-C₆-alkyl group wherein the aminoalkyl nitrogen is (i) unsubstituted or (ii) substituted with one or two radicals independently selected from the group consisting of C₁-C₆-alkyl, ar-C₁-C₆-alkyl, cycloalkyl and C₁-C₆-alkanoyl;
25

R¹³ is selected from the group consisting of a hydrido, benzyl, phenyl, C₁-C₆-alkyl, C₂-C₆-


-596-

alkynyl, C₂-C₆-alkenyl and a C₁-C₆-hydroxyalkyl group; and

- R²⁰ is (a) -O-R²¹, where R²¹ is selected from the group consisting of a hydrido, C₁-C₆-alkyl, aryl, ar-C₁-C₆-alkyl group and a pharmaceutically acceptable cation, (b) -NH-O-R²², wherein R²² is a selectively removable protecting group such as a 2-tetrahydropyranyl, benzyl, p-methoxybenzyl carbonyl-C₁-C₆-alkoxy, trisubstituted silyl group or o-nitrophenyl group, peptide synthesis resin and the like, wherein trisubstituted silyl group is substituted with C₁-C₆-alkyl, aryl, or ar-C₁-C₆-alkyl, or (c) -NH-O-R¹⁴, where R¹⁴ is hydrido, a pharmaceutically acceptable cation or C(W)R¹⁵ where W is O or S and R¹⁵ is selected from the group consisting of an C₁-C₆-alkyl, aryl, C₁-C₆-alkoxy, heteroaryl-C₁-C₆-alkyl, C₃-C₈-cycloalkyl-C₁-C₆-alkyl, aryloxy, ar-C₁-C₆-alkoxy, ar-C₁-C₆-alkyl, heteroaryl and amino C₁-C₆-alkyl group wherein the aminoalkyl nitrogen is (i) unsubstituted or (ii) substituted with one or two substituents independently selected from the group consisting of an C₁-C₆-alkyl, aryl, ar-C₁-C₆-alkyl, C₃-C₈-cycloalkyl-C₁-C₆-alkyl, ar-C₁-C₆-alkoxycarbonyl, C₁-C₆-alkoxycarbonyl, and C₁-C₆-alkanoyl radical, or (iii) wherein the amino C₁-C₆-alkyl nitrogen and two substituents attached thereto form a 5- to 8-membered heterocyclo or heteroaryl ring.


-597-

21. The compound or salt according to claim 20 that corresponds in structure to formula XII or formula XIIA, below,



wherein substituents R^6 , R^{20} , A, and EY are as before defined.

22. The compound or salt according to claim 20 that corresponds in structure to formula XII or formula XIIA, below,




15 wherein substituents R^{20} , A and EY are as before defined.


23. The compound or salt according to claim 20 that corresponds in structure to the formula below

20


-598-



24. The compound or salt according to
claim 20 that corresponds in structure to the formula
5 below




25. The compound or salt according to
10 claim 20 that corresponds in structure to the formula
below




15 26. The compound or salt according to claim
20 that corresponds in structure to the formula below

-599-




27. The compound or salt according to claim 20 that corresponds in structure to the formula below



5


28. The compound or salt according to claim 20 that corresponds in structure to the formula below




10

29. The compound or salt according to claim 20 that corresponds in structure to the formula below

-600-




30. The compound or salt according to claim 20 that corresponds in structure to the formula below



5


31. The compound or salt according to claim 20 that corresponds in structure to the formula below



32. A process for treating a host mammal having a condition associated with pathological matrix metalloprotease (MMP) activity that comprises administering a metalloprotease inhibitor compound or a pharmaceutically acceptable salt thereof in an effective amount to a mammalian host having such a condition, said metalloprotease inhibitor inhibiting the activity of one or more of MMP-2, MMP-9 and MMP-

-601-

13, while exhibiting substantially less inhibitory activity against MMP-1, said compound corresponding in structure to formula I, below,



wherein

R¹ and R² taken together with the carbon to which they are bonded form a heterocyclo or cycloalkyl group optionally substituted by one, two or three R^X substituents, or R¹ and R² are independently selected from the group consisting of:

hydrido,

an alkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

an alkyloxyalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

an alkylthioalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

an alkenyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

an alkynyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

-602-

an aryl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

an arylalkyl group, optionally substituted
5 with one, two or three groups independently selected from R^X substituents,

an arylalkyloxyalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

10 an aryloxyalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

an arylthioalkyl group, optionally substituted with one, two or three groups
15 independently selected from R^X substituents,

an arylalkylthioalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

a cycloalkyl or bicycloalkyl group,
20 optionally substituted with one, two or three groups independently selected from R^X substituents,

a cycloalkenyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

25 a cycloalkylalkyl or bicycloalkylalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

a cycloalkyloxyalkyl or bicycloalkyloxyalkyl group, optionally substituted
30 with one, two or three groups independently selected from R^X substituents,

-603-

a cycloalkylalkyloxyalkyl or bicycloalkyloxyalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

5 a cycloalkylthioalkyl or bicycloalkylthioalkyl group, optionally substituted with one, two or three groups independently selected from R^X;

cycloalkylalkylthioalkyl or
10 bicycloalkylalkylthioalkyl, optionally substituted with one, two or three groups independently selected from R^X substituents,

a heterocyclo group, optionally substituted with one, two or three groups independently selected
15 from R^X substituents,

a heterocycloalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents

a heteroaryl group, optionally substituted
20 with one, two or three groups independently selected from R^X substituents,

a biarylalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

25 an arylalkenyl group, optionally substituted with one, two or three groups independently selected from R^X substituents,

an arylalkynyl group, optionally substituted with one, two or three groups
30 independently selected from R^X substituents,

-604-

a heterocycloalkylthio group, optionally substituted with one, two or three groups selected independently from R^X substituents,

a heterocycloalkyloxyalkyl group,
 5 optionally substituted with one, two or three groups selected independently from R^X substituents,

a heteroarylalkenyl group, optionally substituted with one, two or three groups independently selected from R^X substituents, and

10 a heteroarylalkyloxyalkyl group, optionally substituted with one, two or three groups independently selected from R^X substituents;

wherein an R^X substituent is selected from the group consisting of a hydrido, aryl,
 15 heteroaryl, heterocyclo, aroyl, alkanoyl, heteroaroyl, halogen, cyano, aldehydo, hydroxy, R^{C^d} -amino ($-NR^{C^d}$), R^{C^d} -aminoalkyl, nitro, nitroso, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, alkoxy, aryloxy, heteroaryloxy, alkenyloxy,
 20 alkynyloxy, alkoxyaryl, alkoxyheteroaryl, alkoxyalkyl, R^C -oxyalkyl, alkoxyalkyl, alkylenedioxy, aryloxyalkyl, perfluoroalkyl, trifluoroalkyl, alkylthio, arylthio, alkyloxycarbonyl, alkyloxycarbonyloxy, aryloxycarbonyl,
 25 arylalkyloxycarbonyl, alkyloxycarbonyl- R^C -amino, arylalkyloxycarbonyl- R^C -amino, aryloxycarbonyloxy, carboxy, R^{C^d} -aminocarbonyloxy, R^{C^d} -aminocarbonyl, R^{C^d} -aminoalkanoyl, hydroxy- R^C -aminocarbonyl, R^{C^d} -aminosulfonyl, arylsulfonyl(R^C)amino, R^{C^d} -
 30 aminoalkoxy, R^{C^d} -aminocarbonyl(R^C)amino,

-605-

trifluoromethylsulfonyl(R^C)amino, heteroarylsulfonyl-
(R^C)amino, alkylsulfonyl, arylsulfonyl(R^C)amino,
arylsulfonyl(R^C)aminocarbonyl, alkylsulfonyl-
(R^C)amino, arylcarbonyl(R^C)-aminosulfonyl, and an
5 alkylsulfonyl(R^C)aminocarbonyl substituent;

wherein R^C and R^d are independently
selected from the group consisting of a hydrido,
alkanoyl, arylalkyl, aroyl, bisalkoxyalkyl, alkyl,
haloalkyl, perfluoroalkyl, trifluoromethylalkyl,
10 perfluoroalkoxyalkyl, alkoxyalkyl, cycloalkyl,
heterocycloalkyl, heterocycloalkylcarbonyl, aryl,
heterocyclo, heteroaryl, cycloalkylalkyl,
aryloxyalkyl, heteroaryloxyalkyl,
heteroarylalkoxyalkyl, heteroarylthioalkyl,
15 arylsulfonyl, alkylsulfonyl, heteroarylsulfonyl,
carboxyalkyl, alkoxycarbonylalkyl, aminocarbonyl,
alkyliminocarbonyl, aryliminocarbonyl,
heterocycloiminocarbonyl, arylthioalkyl,
alkylthioalkyl, arylthioalkenyl, alkylthioalkenyl,
20 heteroarylalkyl, haloalkanoyl, hydroxyalkanoyl,
thiolalkanoyl, alkenyl, alkynyl, alkoxyalkyl,
alkoxycarbonyl, aryloxycarbonyl, aminoalkylcarbonyl,
hydroxyalkyl, aminoalkyl, aminoalkylsulfonyl,
aminosulfonyl wherein the amino nitrogen is (i)
25 unsubstituted or (ii) independently substituted with
one or two R^Y radicals, or the substituents on the
amino group taken together with the amino nitrogen
form a saturated or partially unsaturated heterocyclo
group optionally substituted with one, two or three
30 groups independently selected from R^W substituents or
a heteroaryl group optionally substituted with one,

- 606 -

two or three groups independently selected from R^V substituents;

wherein R^Y is selected from the group consisting of an arylalkyl, aryl, heteroaryl, heterocyclo, alkyl, alkynyl, alkenyl, alkoxyalkyl, alkoxyalkyl, substituted or unsubstituted aminoalkyl, alkyloxycarbonyl, arylalkyloxycarbonyl, carboxyalkyl, haloalkyl, alkanoyl, aroyl, substituted or unsubstituted aminoalkanoyl, halo alkanoyl and a hydroxyalkyl group, each of which groups is optionally substituted by one or two groups independently selected from R^U substituents as are the substituents of the substituted aminoalkyl and substituted aminoalkanoyl groups;

wherein R^V is selected from the group consisting of a hydrido, aryl, heteroaryl, heterocyclo, aroyl, alkanoyl, heteroaroyl, halogen, cyano, aldehydo, hydroxy, amino, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, alkoxy, aryloxy, heteroaryloxy, alkenyloxy, alkynyloxy, alkoxyaryl, alkoxyheteroaryl, R^{YR^Z}-amino, alkoxyalkyl, alkylenedioxy, aryloxyalkyl, perfluoroalkyl, trifluoroalkyl, alkylthio, arylthio, alkyloxycarbonyl, alkyloxycarbonyloxy, aryloxycarbonyl, arylalkyloxycarbonyl, arylalkyloxycarbonylamino, aryloxycarbonyloxy, carboxy, R^{YR^Z}-aminocarbonyloxy, R^{YR^Z}-aminocarbonyl, R^{YR^Z}-aminoalkanoyl, hydroxyaminocarbonyl, R^{YR^Z}-aminosulfonyl, R^{YR^Z}-aminocarbonyl(R^Y)amino, trifluoromethylsulfonyl(R^Y)amino, heteroarylsulfonyl(R^Y)amino, arylsulfonyl(R^Y)amino, arylsulfonyl(R^Y)-

-607-

aminocarbonyl, alkylsulfonyl(RY)amino, arylcarbonyl-(RY)aminosulfonyl, and an alkylsulfonyl(RY)-aminocarbonyl substituent;

wherein R^W is selected from the group

5 consisting of a hydrido, aryl, heteroaryl, heterocyclo, aroyl, alkanoyl, heteroaroyl, halogen, cyano, aldehydo, hydroxy, amino, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, alkoxy, aryloxy, heteroaryloxy, alkenyloxy, alkynyloxy, alkoxyaryl,

10 alkoxyheteroaryl, R^YR^Z -amino, alkoxyalkyl, alkylenedioxy, aryloxyalkyl, perfluoroalkyl, trifluoroalkyl, alkylthio, arylthio, alkyloxycarbonyl, alkyloxycarbonyloxy, aryloxycarbonyl, arylalkyloxycarbonyl,

15 arylalkyloxycarbonylamino, aryloxycarbonyloxy, carboxy, R^YR^Z -aminocarbonyloxy, R^YR^Z -aminocarbonyl, R^YR^Z -aminoalkanoyl, hydroxyaminocarbonyl, R^YR^Z -aminosulfonyl, R^YR^Z -aminocarbonyl(RY)amino, trifluoromethylsulfonyl(RY)amino, heteroarylsulfonyl-

20 (RY)amino, arylsulfonyl(RY)amino, arylsulfonyl(RY)-aminocarbonyl, alkylsulfonyl(RY)amino, arylcarbonyl-(RY)aminosulfonyl, and an alkylsulfonyl(RY)-aminocarbonyl substituent;

R^Z is selected from the group consisting of

25 an arylalkyl, aryl, heteroaryl, heterocyclo, alkyl, alkynyl, alkenyl, alkoxyalkyl, alkoxyalkylalkyl, substituted or unsubstituted aminoalkyl, alkyloxycarbonyl, arylalkyloxycarbonyl, carboxyalkyl, haloalkyl, alkanoyl, aroyl, substituted or

30 unsubstituted aminoalkanoyl, halo alkanoyl and a

-608-

hydroxyalkyl group, each of which groups are optionally substituted by one or two R^U substituents;

wherein R^U is selected from the group consisting of an arylalkyl, aryl, heteroaryl, heterocyclo, alkyl, alkynyl, alkenyl, alkoxyalkyl, alkoxyalkyl, substituted or unsubstituted aminoalkyl, alkyloxycarbonyl, arylalkyloxycarbonyl, carboxyalkyl, haloalkyl, alkanoyl, aroyl, substituted or unsubstituted aminoalkanoyl, halo alkanoyl and a hydroxyalkyl group, wherein the substituents of the substituted aminoalkyl and substituted aminoalkanoyl groups are selected from the group consisting of an alkyl, alkenyl, alkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, aryloxycarbonyl and an alkyloxycarbonyl group;

R^{3a} and R^{3b} are independently selected from the group consisting of a hydrido, alkyl, alkenyl, alkynyl, aryl, heterocyclo, heteroaryl, cycloalkyl, and an alkoxyalkyl group, each of which groups is optionally substituted by an -AREY substituent;

in that AREY substituent, A is selected from the group consisting of

- (1) -O-;
- (2) -S-;
- (3) -N(R^e)-;
- (4) -CO-N(R^e) or -N(R^e)-CO-;
- (5) -CO-O- or -O-CO-;
- (6) -O-CO-O-;
- (7) -HC=CH-;
- (8) -NH-CO-NH-;
- (9) -C \equiv C-;
- (10) -NH-CO-O- or -O-CO-NH-;

-609-

- (11) -N=N-;
(12) -NH-NH-;
(13) -CS-N(R^e)- or -N(R^e)-CS-;
(14) -CH₂-;
5 (15) -O-[(CH₂)₁₋₈]- or -[(CH₂)₁₋₈]O-; and
(16) -S-CH₂- or -CH₂-S-; or
(17) A is absent and R is directly

connected to R^{3a} or R^{3b}, or both R^{3a} and R^{3b};

- the moiety R is selected from the group
10 consisting of alkyl, alkoxyalkyl, aryl, heteroaryl,
cycloalkyl, heterocycloalkyl, aralkyl, heteroaralkyl,
heterocycloalkyl, cycloalkylalkyl, cycloalkoxyalkyl,
heterocycloalkoxyalkyl, aryloxyalkyl,
heteroaryloxyalkyl, arylthioalkyl,
15 heteroarylthioalkyl, cycloalkylthioalkyl, and a
heterocycloalkylthioalkyl group wherein the aryl,
heteroaryl, cycloalkyl or heterocycloalkyl
substituent is (i) unsubstituted or (ii) substituted
with one or two radicals selected from the group
20 consisting of a halo, alkyl, perfluoroalkyl,
perfluoroalkoxy, perfluoroalkylthio,
trifluoromethylalkyl, amino, alkoxycarbonylalkyl,
alkoxy, C₁-C₂-alkylene-dioxy, hydroxycarbonylalkyl,
hydroxycarbonylalkylamino, nitro, hydroxy,
25 hydroxyalkyl, alkanoylamino, and a alkoxycarbonyl
group;

the group E is selected from the group
consisting of

- (1) -CO(R^w)- or -(R^w)CO-;
30 (2) -CON(R^e)- or -(R^e)NCO-;
(3) -CO-;

-610-

(4) $-\text{SO}_2-\text{R}^{\text{W}}$ or $-\text{R}^{\text{W}}\text{SO}_2-$;(5) $-\text{SO}_2-$;(6) $-\text{N}(\text{R}^{\text{e}})-\text{SO}_2-$ or $-\text{SO}_2-\text{N}(\text{R}^{\text{e}})-$; or

5 (7) E is absent and R is bonded directly to Y; and

Y is absent or is selected from the group consisting of a hydrido, alkyl, alkoxy, haloalkyl, aryl, aralkyl, cycloalkyl, heteroaryl, hydroxy, nitrile, nitro, aryloxy, aralkoxy, heteroaryloxy, heteroaralkyl, $\text{R}^{\text{C}}\text{oxyalkyl}$, perfluoroalkoxy, perfluoroalkylthio, trifluoromethylalkyl, alkenyl, heterocycloalkyl, cycloalkyl, trifluoromethyl, alkoxy carbonyl, and an aminoalkyl group, wherein the aryl, heteroaryl or heterocycloalkyl group is (i) 10 unsubstituted or (ii) substituted with one or two radicals independently selected from the group consisting of an alkanoyl, halo, nitro, nitrile, haloalkyl, alkyl, aralkyl, aryl, alkoxy, and an amino group wherein the amino nitrogen is (i) unsubstituted 15 or (ii) substituted with one or two groups independently selected from hydrido, alkyl, and an aralkyl group;

wherein R^{e} is selected from hydrido, alkyl, alkenyl, alkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, aryloxycarbonyl, alkyloxycarbonyl, $\text{R}^{\text{C}}\text{R}^{\text{d}}\text{amino carbonyl}$, $\text{R}^{\text{C}}\text{R}^{\text{d}}\text{aminosulfonyl}$, $\text{R}^{\text{C}}\text{R}^{\text{d}}\text{aminoalkanoyl}$ and $\text{R}^{\text{C}}\text{R}^{\text{d}}\text{aminoalkysulfonyl}$, and R^{C} , R^{d} and R^{W} are as defined before; or

$\text{R}^{3\text{a}}$ and $\text{R}^{3\text{b}}$ taken together with the nitrogen atom to which they are bonded form a group -GAREY 30 wherein

-611-

G is a N-heterocyclo group;

the substituent A is selected from the group consisting of

- (1) -O-;
- 5 (2) -S-;
- (3) -NR^e-;
- (4) -CO-N(R^e) or -N(R^e)-CO-;
- (5) -CO-O- or -O-CO-;
- (6) -O-CO-O-;
- 10 (7) -HC=CH-;
- (8) -NH-CO-NH-;
- (9) -C≡C-;
- (10) -NH-CO-O- or -O-CO-NH-;
- (11) -N=N-;
- 15 (12) -NH-NH-;
- (13) -CS-N(R^e)- or -N(R^e)-CS-;
- (14) -CH₂-;
- (15) -O-[(CH₂)₁₋₈]- or -[(CH₂)₁₋₈]O-; and
- (16) -S-CH₂- or -CH₂-S-; or
- 20 (17) A is absent and R is directly bonded to G;

the moiety R is selected from the group consisting of alkyl, alkoxyalkyl, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaralkyl, 25 heterocycloalkyl, cycloalkylalkyl, cycloalkoxyalkyl, heterocycloalkoxyalkyl, aryloxyalkyl, heteroaryloxyalkyl, arylthioalkyl, heteroarylthioalkyl, cycloalkylthioalkyl, and a heterocycloalkylthioalkyl group wherein the aryl or 30 heteroaryl or cycloalkyl or heterocycloalkyl substituent is (i) unsubstituted or (ii) substituted

-612-

with one or two radicals selected from the group consisting of a halo, alkyl, perfluoroalkyl, perfluoroalkoxy, perfluoroalkylthio, trifluoromethylalkyl, amino, alkoxycarbonylalkyl, alkoxy, C₁-C₂-alkylene-dioxy, hydroxycarbonylalkyl, hydroxycarbonylalkylamino, nitro, hydroxy, hydroxyalkyl, alkanoylamino, and a alkoxycarbonyl group;

the moiety E is selected from the group consisting of


- (1) -CO(R^W)- or -(R^W)CO-;
- (2) -CONH- or -HNCO-;
- (3) -CO-;
- (4) -SO₂-R^W- or -R^W-SO₂-;
- (5) -SO₂-;
- (6) -NH-SO₂- or -SO₂-NH-; or
- (7) E is absent and R is bonded directly to Y; and

the moiety Y is absent or is selected from the group consisting of a hydrido, alkyl, alkoxy, haloalkyl, aryl, aralkyl, cycloalkyl, heteroaryl, hydroxy, aryloxy, aralkoxy, heteroaryloxy, heteroaralkyl, perfluoroalkoxy, perfluoroalkylthio, trifluoromethylalkyl, alkenyl, heterocycloalkyl, cycloalkyl, trifluoromethyl, alkoxycarbonyl, and a aminoalkyl group, wherein the aryl or heteroaryl or heterocycloalkyl group is (i) unsubstituted or (ii) substituted with one or two radicals independently selected from the group consisting of an alkanoyl, halo, nitro, aralkyl, aryl, alkoxy, and an amino group wherein the amino nitrogen is (i) unsubstituted or (ii) substituted with one or two groups

-613-


independently selected from hydrido, alkyl, and an aralkyl group.

33. The process according to claim 32
 5 wherein said compound or salt corresponds in structure to formula IIIA, below,




wherein substituents R^1 , R^2 and -GAREY are
 10 as before defined.

34. The process according to claim 32
 wherein said compound or salt corresponds in structure to formula IVA, below,
 15



wherein substituents R^1 , R^2 and -AREN are
 as before defined.

20 35. The process according to claim 32
 wherein said compound or salt corresponds in structure to formula VA, below,




25 wherein the substituent -AREN is as before defined.

-614-

36. The process according to claim 32 wherein said compound or salt corresponds in structure to formula VIA, below,


5



wherein substituents R^1 , R^2 , and -EY are as before defined, and A is -CH₂-, -O-CH₂-, -CH₂-O-, -S-CH₂- or -CH₂-S-.

10

37. The process according to claim 32 wherein said compound or salt corresponds in structure to formula VIIA, below,




15

wherein substituent -EY is as before defined as part of an -AREY or -GAREY group, and A is -CH₂-, -O-CH₂-, -CH₂-O-, -S-CH₂- or -CH₂-S-.

20

38. The process according to claim 32 wherein said compound or salt corresponds in structure to formula VIIIA, below,

-615-



wherein

R^{3a} , R^{3b} and are as defined before; and

5 m is zero, 1 or 2;

n is zero, 1 or 2;

p is zero, 1 or 2;


the sum of $m + n + p = 1, 2, 3$ or 4;

(a) one of X , Y and Z is selected from the
 10 group consisting of $C(O)$, NR^6 , O , S , $S(O)$, $S(O)_2$ and
 $NS(O)_2R^7$, and the remaining two of X , Y and Z are
 CR^8R^9 , and $CR^{10}R^{11}$, or

(b) X and Z or Z and Y together constitute
 a moiety that is selected from the group consisting
 15 of $NR^6C(O)$, $NR^6S(O)$, $NR^6S(O)_2$, NR^6S , NR^6O , SS , NR^6NR^6
 and $OC(O)$, with the remaining one of X , Y and Z being
 CR^8R^9 , or

(c) n is zero and X , Y and Z together
 constitute a moiety selected from the group
 20 consisting of

-616-



5 wherein wavy lines are bonds to the atoms
of the depicted ring;

R^6 and $R^{6'}$ are independently selected from
the group consisting of hydrido, C_1 - C_6 -alkanoyl, C_6 -
aryl- C_1 - C_6 -alkyl, aroyl, bis(C_1 - C_6 -alkoxy- C_1 - C_6 -
10 alkyl)- C_1 - C_6 -alkyl, C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl, C_1 -
 C_6 -perfluoroalkyl, C_1 - C_6 -trifluoromethylalkyl, C_1 - C_6 -

-617-

perfluoroalkoxy-C₁-C₆-alkyl, C₁-C₆-alkoxy-C₁-C₆-
 alkyl, C₃-C₆-cycloalkyl, C₃-C₈-heterocycloalkyl, C₃-
 C₈-heterocycloalkylcarbonyl, C₆-aryl, C₅-C₆-
 heterocyclo, C₅-C₆-heteroaryl, C₃-C₈-cycloalkyl-C₁-
 5 C₆-alkyl, C₆-aryloxy-C₁-C₆-alkyl, heteroaryloxy-C₁-
 C₆-alkyl, heteroaryl-C₁-C₆-alkoxy-C₁-C₆-alkyl,
 heteroarylthio-C₁-C₆-alkyl, C₆-arylsulfonyl, C₁-C₆-
 alkylsulfonyl, C₅-C₆-heteroarylsulfonyl, carboxy-C₁-
 C₆-alkyl, C₁-C₄-alkoxycarbonyl-C₁-C₆-alkyl,
 10 aminocarbonyl, C₁-C₆-alkyliminocarbonyl, C₆-
 aryliminocarbonyl, C₅-C₆-heterocycloiminocarbonyl,
 C₆-arylthio-C₁-C₆-alkyl, C₁-C₆-alkylthio-C₁-C₆-alkyl,
 C₆-arylthio-C₃-C₆-alkenyl, C₁-C₄-alkylthio-C₃-C₆-
 alkenyl, C₅-C₆-heteroaryl-C₁-C₆-alkyl, halo-C₁-C₆-
 15 alkanoyl, hydroxy-C₁-C₆-alkanoyl, thiol-C₁-C₆-
 alkanoyl, C₃-C₆-alkenyl, C₃-C₆-alkynyl, C₁-C₄-alkoxy-
 C₁-C₄-alkyl, C₁-C₅-alkoxycarbonyl, aryloxycarbonyl,
 NR⁸R⁹-C₁-C₅-alkylcarbonyl, hydroxy-C₁-C₅-alkyl, an
 aminocarbonyl wherein the aminocarbonyl nitrogen is
 20 (i) unsubstituted or (ii) substituted with one or two
 radicals independently selected from the group
 consisting of C₁-C₆-alkyl, ar-C₁-C₆-alkyl, C₃-C₈-
 cycloalkyl and a C₁-C₆-alkanoyl group,
 hydroxyaminocarbonyl, an aminosulfonyl group wherein
 25 the aminosulfonyl nitrogen is (i) unsubstituted or
 (ii) substituted with one or two radicals
 independently selected from the group consisting of
 C₁-C₆-alkyl, ar-C₁-C₆-alkyl, C₃-C₈-cycloalkyl and a

-618-

C₁-C₆-alkanoyl group, an amino-C₁-C₆-alkylsulfonyl group wherein the amino-C₁-C₆-alkylsulfonyl nitrogen is (i) unsubstituted or (ii) substituted with one or two radicals independently selected from the group
5 consisting of C₁-C₆-alkyl, ar-C₁-C₆-alkyl, C₃-C₈-cycloalkyl and a C₁-C₆-alkanoyl group and an amino-C₁-C₆-alkyl group wherein the aminoalkyl nitrogen is (i) unsubstituted or (ii) substituted with one or two radicals independently selected from the group
10 consisting of C₁-C₆-alkyl, ar-C₁-C₆-alkyl, C₃-C₈-cycloalkyl and a C₁-C₆-alkanoyl group;

R⁷ is selected from the group consisting of a arylalkyl, aryl, heteroaryl, heterocyclo, C₁-C₆-alkyl, C₃-C₆-alkynyl, C₃-C₆-alkenyl, C₁-C₆-carboxyalkyl and a C₁-C₆-hydroxyalkyl group;
15

R⁸ and R⁹ and R¹⁰ and R¹¹ are independently selected from the group consisting of a hydrido, hydroxy, C₁-C₆-alkyl, aryl, ar-C₁-C₆-alkyl, heteroaryl, heteroar-C₁-C₆-alkyl, C₂-C₆-alkynyl, C₂-C₆-alkenyl, thiol-C₁-C₆-alkyl, C₁-C₆-alkylthio-C₁-C₆-alkyl cycloalkyl, cycloalkyl-C₁-C₆-alkyl, heterocycloalkyl-C₁-C₆-alkyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, aralkoxy-C₁-C₆-alkyl, C₁-C₆-alkoxy-C₁-C₆-alkoxy-C₁-C₆-alkyl, hydroxy-C₁-C₆-alkyl,
20 hydroxycarbonyl-C₁-C₆-alkyl, hydroxycarbonylar-C₁-C₆-alkyl, aminocarbonyl-C₁-C₆-alkyl, aryloxy-C₁-C₆-alkyl, heteroaryloxy-C₁-C₆-alkyl, arylthio-C₁-C₆-alkyl, heteroarylthio-C₁-C₆-alkyl, the sulfoxide or
25

-619-

sulfone of any said thio substituents, perfluoro-C₁-C₆-alkyl, trifluoromethyl-C₁-C₆-alkyl, halo-C₁-C₆-alkyl, alkoxy-carbonylamino-C₁-C₆-alkyl and an amino-C₁-C₆-alkyl group wherein the aminoalkyl nitrogen is

5 (i) unsubstituted or (ii) substituted with one or two radicals independently selected from the group consisting of C₁-C₆-alkyl, ar-C₁-C₆-alkyl, cycloalkyl and C₁-C₆-alkanoyl, or wherein R⁸ and R⁹ or R¹⁰ and R¹¹ and the carbon to which they are bonded form a

10 carbonyl group, or wherein R⁸ and R⁹ or R¹⁰ and R¹¹, or R⁸ and R¹⁰ together with the atoms to which they are bonded form a 5- to 8-membered carbocyclic ring, or a 5- to 8-membered heterocyclic ring containing one or two heteroatoms that are nitrogen, oxygen, or

15 sulfur, with the proviso that only one of R⁸ and R⁹ or R¹⁰ and R¹¹ is hydroxy;

R¹² and R^{12'} are independently selected from the group consisting of a hydrido, C₁-C₆-alkyl, aryl, ar-C₁-C₆-alkyl, heteroaryl, heteroaralkyl, C₂-C₆-alkynyl, C₂-C₆-alkenyl, thiol-C₁-C₆-alkyl,

20 cycloalkyl, cycloalkyl-C₁-C₆-alkyl, heterocycloalkyl-C₁-C₆-alkyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, aryloxy-C₁-C₆-alkyl, amino-C₁-C₆-alkyl, C₁-C₆-alkoxy-C₁-C₆-alkoxy-C₁-C₆-alkyl, hydroxy-C₁-C₆-alkyl, hydroxycarbonyl-C₁-C₆-alkyl, hydroxycarbonylar-C₁-C₆-alkyl,


25 aminocarbonyl-C₁-C₆-alkyl, aryloxy-C₁-C₆-alkyl, heteroaryloxy-C₁-C₆-alkyl, C₁-C₆-alkylthio-C₁-C₆-alkyl, arylthio-C₁-C₆-alkyl, heteroarylthio-C₁-C₆-

-620-

alkyl, the sulfoxide or sulfone of any said thio
 substituents, perfluoro-C₁-C₆-alkyl, trifluoromethyl-
 C₁-C₆-alkyl, halo-C₁-C₆-alkyl, alkoxycarbonylamino-
 C₁-C₆-alkyl and an amino-C₁-C₆-alkyl group wherein
 5 the aminoalkyl nitrogen is (i) unsubstituted or (ii)
 substituted with one or two radicals independently
 selected from the group consisting of C₁-C₆-alkyl,
 ar-C₁-C₆-alkyl, cycloalkyl and C₁-C₆-alkanoyl; and

R¹³ is selected from the group consisting
 10 of a hydrido, benzyl, phenyl, C₁-C₆-alkyl, C₂-C₆-
 alkynyl, C₂-C₆-alkenyl and a C₁-C₆-hydroxyalkyl
 group.

39. The process according to claim 32
 15 wherein said compound or salt corresponds in
 structure to formula XIA, below,




20 wherein the definitions for X, Y, Z, m, n,
 p A, E, Y and R²⁰ are as before defined.

40. The process according to claim 39
 wherein A is absent.

25

-621-

41. The process according to claim 32 wherein said compound or salt corresponds in structure to formula XIIA, below,




5

wherein substituents R^6 , A and EY are as before defined.

42. The process according to claim 41
10 wherein A is absent.

43. The process according to claim 32 wherein said compound or salt corresponds in structure to formula XIIA, below,

15



wherein substituents A and EY are as before defined.

20

44. The process according to claim 43 wherein A is absent.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/03061

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D405/12 A61K31/445 C07D211/66 C07D211/46 C07D309/08
 C07D401/14 C07D409/12 C07D405/14 C07D491/10 A61K31/453
 A61P19/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 38163 A (AMERICAN CYANAMID CO) 3 September 1998 (1998-09-03) cited in the application claim 1	2, 32
X	WO 98 37877 A (AMERICAN CYANAMID CO) 3 September 1998 (1998-09-03) cited in the application claim 1	2, 32
A	GB 1 067 965 A (KENNETH ET AL) 1967 page 1, right-hand column, line 58 - line 60; example 5	
A	WO 97 48368 A (SCHERING CORP) 24 December 1997 (1997-12-24) example 52	
	-/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

24 May 2000

Date of mailing of the international search report

06/06/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

De Jong, B

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/03061

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 336 354 A (HOECHST AG) 11 October 1989 (1989-10-11) example 534 ----	
P,X	EP 0 931 788 A (PFIZER LTD ;PFIZER (US)) 28 July 1999 (1999-07-28) claim 1; examples ----	2,32
P,X	WO 99 29667 A (WHITLOCK GAVIN ALISTAIR ;DACK KEVIN NEIL (GB); PFIZER LTD (GB); PF) 17 June 1999 (1999-06-17) cited in the application claim 1; examples ----	2,32
P,X	WO 99 24399 A (DARWIN DISCOVERY LTD) 20 May 1999 (1999-05-20) cited in the application claim 1; examples ----	2,32
P,X	WO 99 25687 A (CRESCENZO GARY A DE ;MCDONALD JOSEPH J (US); BOEHM TERRI L (US); S) 27 May 1999 (1999-05-27) claim 1 -----	2,32

INTERNATIONAL SEARCH REPORT

international application No.

PCT/US 00/ 03061

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1,3

Present claims 1,3 relate to an extremely large number of possible compounds. It is noted that claim 1 even comprises known compounds having a relatively simple structure such as for instance aminosulfonylacetic acid. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to compounds according to claims 2,4-31.

It should however be noted that the search and the search report can be considered as covering all claimed compounds of the prior art insofar these compounds display a metalloprotease inhibiting activity.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT
Information on patent family members

International Application No
PCT/US 00/03061

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9838163 A	03-09-1998	AU 6168698 A	18-09-1998
		AU 6436898 A	18-09-1998
		BR 9807803 A	22-02-2000
		EP 0973512 A	26-01-2000
		EP 0970046 A	12-01-2000
		NO 994124 A	26-10-1999
		NO 994125 A	26-10-1999
		WO 9837877 A	03-09-1998
WO 9837877 A	03-09-1998	AU 6168698 A	18-09-1998
		AU 6436898 A	18-09-1998
		BR 9807803 A	22-02-2000
		EP 0973512 A	26-01-2000
		EP 0970046 A	12-01-2000
		NO 994124 A	26-10-1999
		NO 994125 A	26-10-1999
		WO 9838163 A	03-09-1998
GB 1067965 A		NONE	
WO 9748368 A	24-12-1997	AU 3567397 A	07-01-1998
		CA 2258752 A	24-12-1997
		CN 1228090 A	08-09-1999
		EP 0912571 A	06-05-1999
EP 0336354 A	11-10-1989	DE 3811777 A	19-10-1989
		AT 113588 T	15-11-1994
		AU 614824 B	12-09-1991
		AU 3254289 A	12-10-1989
		BR 8901645 A	21-11-1989
		DD 283761 A	24-10-1990
		DE 58908577 D	08-12-1994
		DK 169989 A	09-10-1989
		ES 2065348 T	16-02-1995
		HU 51603 A,B	28-05-1990
		JP 2787587 B	20-08-1998
		NZ 228630 A	25-06-1991
		PH 26217 A	01-04-1992
		US 5015284 A	14-05-1991
		ZA 8902495 A	29-11-1989
EP 0931788 A	28-07-1999	JP 11292839 A	26-10-1999
WO 9929667 A	17-06-1999	AU 1230199 A	28-06-1999
WO 9924399 A	20-05-1999	AU 1046999 A	31-05-1999
WO 9925687 A	27-05-1999	AU 1373299 A	07-06-1999